

2014

# Satellite remote sensing of primary production in the Bering Sea

Anugerahanti, P.

Anugerahanti, P. (2014) 'Satellite remote sensing of primary production in the Bering Sea', The Plymouth Student Scientist, 7(2), p. 195-202.

<http://hdl.handle.net/10026.1/14075>

---

The Plymouth Student Scientist  
University of Plymouth

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# Satellite remote sensing of primary production in the Bering Sea

Prima Anugerahanti

*Project Advisor: [Jill Schwartz](#), School of Marine Science & Engineering, Plymouth University, Drake Circus, Plymouth, PL4 8AA*

## Introduction

The Bering Sea covers over 2 million square kilometres of the northernmost region of the Pacific Ocean (NPO, 2008), and is considered to be one of the most productive seas in the world (Walsh et al., 1989). Linking the Pacific and Arctic Oceans, the Bering Sea is almost entirely surrounded by the landmasses of Alaska and Russia. A steep continental slope divides this sea between the expansive continental shelf (<200 m depth) on its eastern waters and the deep basin (>2000 m depth) to the west (Brown et al., 2011).

Sustained with nutrient supplied from the warmer Pacific inflow (Grebmeier et al., 2006), rate of primary productivity in the region is controlled by sea-ice cover, ice-melt stratification, and light condition (Alexander and Niebauer, 1981). During sea-ice formation, brine rejection triggers convective mixing that restores nutrient to surface water (Stabeno et al., 2010). These nutrients are consumed by phytoplankton in spring when freshwater input from melting sea-ice stratifies the surface ocean, leading to higher than average mixed-layer light level and the start of the spring phytoplankton bloom in the marginal ice zone or at the ice edge (Niebauer et al., 2000).

However, sea ice loss also creates additional open water habitat for phytoplankton, whose growth is thought to be light-limited under the sea-ice cover (Hill and Cota, 2005). Rates of CO<sub>2</sub> fixation in the ice-free zone are generally far higher than in sea-ice habitats (Arrigo, 2003), therefore, sea-ice loss potentially leads to a more productive sea. Nevertheless, warming and sea ice loss are likely to affect patterns of primary productivity including the timing and magnitude of the spring bloom (ACIA, 2005). This can also remove the platform for the growth of sea-ice algae, which may act as a seed population for the phytoplankton bloom as the ice-edge moves north each spring (Spindler, 1994).

Satellite ocean colour remote sensing is potentially suitable for quantifying primary productivity in the ocean, worldwide (Hirawake et al., 2012). The simplest productivity model evaluates primary production integrated by depth and time, as a function of sea

surface chlorophyll (e.g. Smith et al., 1982; Eppley et al., 1985) derived from satellite data. A more complex algorithm introduces surface irradiance as a second factor controlling productivity, where depth-integrated production is the product of depth-integrated chlorophyll, daily surface irradiance, and a constant, water-column averaged quantum yield ( $\Psi$ ) for photosynthesis (Morel 1978; Falkowski 1981; Platt 1986; Morel 1991; Behrenfield and Falkowski, 1997).

Studies using satellite remote sensing have revealed an increase in satellite-detectable annual primary production in the Arctic due to the increase in open-water area (Arrigo et al., 2008; Pabi et al., 2008). Conversely satellite-based studies by Brown et al (2011), showed that the seasonal ice pack of the Bering Sea has not contracted since 1979 to 2009, hence the overall primary productivity hasn't increased during those years. In fact the annual primary productivity of the Bering Sea ( $124 \text{ g C m}^{-2} \text{ yr}^{-1}$ ) is below the global mean ( $140 \text{ g C m}^{-2} \text{ yr}^{-1}$ ) (Brown et al., 2011). Proxy records have also indicated a decline in the Bering Sea primary production this century (Schell 2000; Hirons et al., 2001). Similar studies also reported cooling rather than warming (Comiso, 2003) and no change or even an increase in Bering Sea ice extent (Liu et al., 2004; Moore and Laidre, 2006; Parkinson and Cavalieri, 2008; Brown et al., 2011; Brown and Arrigo, 2012). Using the algorithm developed by Pabi et al (2008), a study by Brown and Arrigo (2012) suggest that this phenomenon is caused by the Arctic winter wind bringing frigid air southwards over the past six decades.

Yet, a number of recent studies claim that the Bering Sea is rapidly warming and that its seasonal sea ice cover is retreating (Hunt et al., 2002; Overland and Stabeno, 2004; Grebmeier et al., 2006; Stabeno et al., 2007). According to Hirawake et al (2012) this is caused by the increase in temperature of the Pacific summer water (Shimada et al., 2006) and heat flux in the Bering Strait over the past decade (Mizobata et al., 2010; Woodgate et al., 2010), which accelerates sea-ice reduction.

## **Discussion**

Although it remains uncertain how primary production is affected by sea-ice loss and warming (Grebmeier et al., 2010), Loeng et al (2005) predict an association between an increase in Bering Sea productivity with a longer ice-free growing season, which has been observed and modelled in the adjacent Arctic Ocean. However, the increase of satellite-derived primary productivity in the Bering Sea has been questioned by Lee et al. (2012) because of the overestimation of chlorophyll-a concentration due to CDOM because it affects the absorption of short wavelength. Nonetheless this increase in primary productivity has been observed during summer 2002 to 2010, using absorption based approaches from satellite remote sensing (Hirawake et al., 2012).

Despite being a suitable tool to quantify primary production by deriving chlorophyll-a concentration, there is an unresolved problem associated with the subsurface chlorophyll maximum (SCM) detection. Investigation by Arrigo and van Dijken (2011) on the effect of the SCM on the estimation of annual primary production from satellite imagery showed that when neglecting the SCM a 7.6% underestimation of primary production results. In addition, a study by Pabi et al (2008) shows that an estimation of 10% error is found in the annual net primary production in the Arctic Ocean is present when neglecting the SCM.

Another drawback in the estimation of primary production using satellite ocean colour is the accuracy of chlorophyll-a estimation used in algorithms. Coloured dissolved organic

matter (CDOM) from the terrestrial in the Arctic (Gue`guen et al., 2007) strongly affects the absorption of short wavelengths, about 400 nm, which results in an overestimation of chlorophyll-a concentration. This effect has pronounced seasonality as a consequence of photobleaching along with additional flow of Alaskan Coastal Waters towards the Bering Sea, and creation of CDOM from the degradation of phytoplankton (Matsuoka et al., 2011). Non-algal particles (NAP) and CDOM have similar effects on chlorophyll-a estimation and change seasonally (Matsuoka et al., 2011). Another error contribution comes from more highly packaged pigments related to large cells, such as diatoms (Cota et al., 2004).

As an improvement of the vertically generalised product model, (VGPM; Behrenfield and Falkowski, 1997), an absorption-based primary productivity model (ABPM) was developed for the southern ocean (Hirawake et al., 2011). The ABPM uses the light absorption coefficient of phytoplankton, determined from the optical density of particles from in-situ data, and applied to MODIS data by trapezoidal integration at the wavelength of MODIS bands (412-55nm) (Hirawake et al., 2012) thus it is possible to reduce inputs of uncertainties in satellite chlorophyll-a data from SCM, CDOM, and NAP, and photosynthetic rate as a SST. The VGPM model is a function of satellite-derived chlorophyll-a and SST:

$$PP_{eu} = P_{opt}^b \times \frac{0.66125 \times E_o}{E_o + 4.1} \times Z_{eu} \times C_{surf} \times D_{irr}$$

where  $C_{surf}$  is sea surface chlorophyll-a concentration,  $Z_{eu}$  is the euphotic depth,  $P_{opt}^b$  is the maximum photosynthetic rate at depth, and  $D_{irr}$  is the photoperiod. Marra et al. (2007) showed that near-surface primary productivity has a linear and invariant relationship with a phytoplankton absorption coefficient,  $\bar{a}_{ph}(0-)$  rather than with chlorophyll-a concentration. Thus,  $P_{opt}^b \times C_{surf}$  was replaced by:

$$PP_{eu} = f[\bar{a}_{ph}(0-)] \times \frac{0.66125 \times E_o}{E_o + 4.1} \times Z_{eu} \times D_{irr}$$

In this study, it is assumed that chlorophyll-a concentration is vertically homogenous, and similar to the  $C_{surf}$  in the context of this model (Hirawake et al., 2012). Hirawake et al. (2012) uses MODIS-Aqua daily averaged, with 4km resolution, from 2002-2010 to quantify the primary productivity. Using this model, the level of primary productivity obtained is less than half that of the estimation from VGPM. High productivity that is found in VGPM ( $>10\ 000\ \text{mg C m}^{-2}\ \text{day}^{-1}$ ) is not found using ABPM apart from in estuaries and river mouths, where in-situ data was not collected. The high difference between  $PP_{eu}$  in ABPM and VGPM might be caused by the overestimation of the chlorophyll-a concentration by VGPM, related to high absorption in blue wavelength by CDOM and NAP (Hirawake et al, 2012). However this study does not taken account of the sea-ice cover, winds, and the Bering Sea environment.

On the other hand, it can be assumed from the decline of benthic biomass and oxygen utilization (Grebmeier et al., 2006), as well as from a reduction in transport through the Aleutian Passes providing limited nutrients to the continental shelf (Schumaker and Alexander, 1999) that primary productivity at the Bering Sea has declined. Over decadal time-scales, such as from the 1970s-1990s, a decrease in primary productivity has also

been suggested by Lee et al., (2007; 2012) The decline is also supported by the proxy record (Schell 2000; Hirons et al., 2001). Recent satellite-based studies by Brown and Arrigo (2012) shows that net primary productivity at the Bering Sea, apart from the Chirikov Basin, which due to sea ice loss has increased by 30% of its net primary productivity, has not increased.

The study by Brown and Arrigo (2012) used satellite reanalysis data of sea ice concentration, sea surface temperature (SST), atmospheric forcing, and ocean colour, coupled to the primary productivity algorithm of Arrigo et al. (2008) and modified by Pabi et al. (2008), to both characterise environmental change and quantify net primary production (NPP) in the Bering Sea. Although this primary production algorithm is adapted for the Southern Ocean, this algorithm calculates the rate of primary production as a function of diurnal changes in spectral downwelling irradiance, sea surface temperature, (°C) and chlorophyll-a concentration ( $\text{mg m}^{-3}$ ). Horizontal distributions of chlorophyll-a were determined from satellite derived chlorophyll-a concentration. This algorithm assumes that within the upper mixed layer, chlorophyll a concentration is uniform and decreases exponentially with depths (Arrigo et al., 2008). The relationship between the chlorophyll-a,  $Chl(0)$ , on the surface, derived from satellite data, and its reduction is obtained from chlorophyll-a profiles from the Southern Ocean (Arrigo et al., 2000), where chlorophyll-a at depth,  $z$  is:

$$Chl(z) = Chl(0)\exp[(0.033(z - Mixed\ Layer\ Depth))]$$

To derive the daily primary productivity at 100m depth in time,  $t$ , hourly in 24 hours, net biomass-specific phytoplankton growth rate,  $G(z,t)$  along with phytoplankton chlorophyll-a to carbon ratio,  $C/Chl$ , is integrated with depth:

$$PP = \int_{z=0}^{100} \int_{t=0}^{t=24} Chl(z) \frac{C}{Chl} G(z, t) dt dz$$

The net biomass is calculated at each depth and each hour as a function of the temperature dependent upper limit to net growth and a light limitation term. This term is calculated using the Beer-Lambert Law and the ratio of photosynthetically usable radiation (PUR) and the spectral photoacclimation parameter (Arriago and Sullivan, 1994; Arrigo et al., 2008; Pabi et al., 2008).

In this study, the near shore chlorophyll-a pixels were replaced by the daily mean of valid pixels within the same region of interest (Brown et al., 2011), which might cause an underestimate in the overall result. Nevertheless, this study relied upon primary production algorithms with inherited errors for SeaWiFS (Arrigo et al., 2008; Pabi et al., 2008; Brown and Arrigo, 2012) and does not taken account of the error from CDOM and SCM. Moreover, this algorithm was initially developed for the Southern Ocean rather than the Arctic Ocean, where the chlorophyll-a decreasing relationship equation is obtained from Southern Ocean dataset, not from the Arctic Ocean. The input data for the algorithm also has lower resolution, since it uses 9km resolution for both SeaWiFS and MODIS-Aqua. This will not capture better spatial variability in a very productive place, with varied topography. Additional errors may arise from the use of weekly averaging of the input chlorophyll datasets because time-scales of phytoplankton growth is short (less than 10 weeks) in high latitudes (Racault et al., 2012), as well as the use of baseline correction to join chlorophyll datasets from different sensors

(SeaWiFS and MODIS-Aqua). This is because SeaWiFS dataset is only available from the years 1998 through 2007, therefore to fill the data gap for the year 2008 to 2010, Brown and Arrigo (2012) join two different sensors, SeaWiFS and MODIS-Aqua. Furthermore, an analysis of the mean chlorophyll-a concentration in Arctic waters between 2003 and 2007 where these two datasets are available, shows that SeaWiFS-derived chlorophyll-a concentration exceeds those from MODIS-Aqua by ~ 2.6%.

The sum of studies to date suggests that a more accurate estimation of primary productivity would be obtained using ABPM along with physical oceanography data, such as sea-ice loss, wind speed, and temperature, and nutrient analysis. This could also establish whether the trends of increasing primary productivity and warming of the sea that are calculated using the model are real.

### **Conclusion**

Two different models were used to estimate the primary productivity at the Bering Sea. The first model, by Hirawake et al. (2012) is an improvement on VGPM. This model utilised an absorption coefficient from in-situ phytoplankton samples and applied it to MODIS bands in order to calculate primary productivity, rather than just surface chlorophyll-a concentration. By applying this model, CDOM and NAP effect is minimised. This could be seen where using ABPM, the productivity estimated is less than half of the VGPM. Using this approach the net primary productivity at the Bering Sea has increased over the year of 2002 -2010. However, in this study, the authors do not provide any link between physical forcing and the Bering Sea environment. Conversely, using an algorithm by Arrigo et al (2008) then modified by Pabi et al. (2008), primary productivity has decreased at the Bering Sea from 1998-2010. However, the algorithm was initially developed for the Southern Ocean and uses two different sensors (SeaWiFS and MODIS-Aqua), which introduces baseline errors when joining two sensors together. Additional errors may also arise from weekly averaging of the input chlorophyll datasets, and they were derived at lower resolutions than the model by Hirawake et al. (2012). This primary productivity algorithm has also neglected the uncertainties contributed by CDOM, SCM, and NAP. Nonetheless, this result is coupled with sea ice loss, air temperature, and wind stress, and spans a longer time period. If the author uses ABPM, this study could produce more accurate results and confirm the trends between sea-ice loss and an increase in primary productivity.

### **Reference**

- ACIA (Arctic Climate Impact Assessment). 2005. Arctic Climate Impact Assessment, scientific report. Cambridge University Press, Cambridge, UK
- Alexander, V., and Niebauer, H. J. 1981. Oceanography of the eastern Bering Sea ice-edge zone in spring. *Limnology and Oceanography*, 26: 1111-1125.
- Arrigo, K. R. 2003. Primary production in sea ice. In *Sea Ice-an Introduction to Its Physics, Chemistry, Biology, and Geology*, pp. 143-183. Ed. by D. N. Thomas, and G. S. Dieckmann. Blackwell Science, Oxford, UK.
- Arrigo, K. R., and van Dijken, G. L. 2011. Secular trends in Arctic Ocean net primary production. *Journal of Geophysical Research*, 116: C09011.
- Arrigo, K. R., DiTullio, G. R., Dunbar, R. B., Lizotte, M. P., Robinson, D. H., Van Woert, M., and Worthen, D. L. (2000), Phytoplankton taxonomic variability and nutrient utilization and primary production in the Ross Sea, *J. Geophys. Res.*, 105, 8827-8846, doi:10.1029/1998JC000289

- Arrigo, K. R., van Dijken, G. L., and Bushinsky, S. 2008. Primary production in the Southern Ocean, 1997-2006. *Journal of Geophysical Research*, 113: C08004.
- Beaugrand, G., Reid, P.C., Ibañez, F., Alistair Lindley, J. and Edwards, M. 2002. Reorganization of North Atlantic marine copepod biodiversity and climate, *Science*, 296: 1692-1694
- Behrenfeld, M. J., and Falkowski, P. G. 1997. Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnology and Oceanography*, 42: 1-20.
- Brown, Z. W., van Dijken, G. L., and Arrigo, K. R. 2011. A reassessment of primary production and environmental change in the Bering Sea. *Journal of Geophysical Research*, 116: C08014.
- Brown, Z. W., and Arrigo, K. R. 2012. Contrasting trends in sea ice and primary production in the Bering Sea and Arctic Ocean. - *ICES Journal of Marine Science*, 69: 1180-1193.
- Cavaleri, D., C. Parkinson, P. Gloersen, and H. J. Zwally. 2008. Sea ice concentrations from Nimbus-7 SMMR and DMSP SSM/I passive microwave data, Jan. 1998 to Dec. 2009, <http://nsidc.org/data/nsidc?0051.html>, Natl. Snow and Ice Data Cent., Boulder, Colo.
- Comiso, J. C. 2003. Warming trends in the Arctic from clear sky satellite observations, *J. Clim.*, 16: 3498-3510, doi:10.1175/1520-0442(2003)016<\$3498:WTITAF\$>\$2.0.CO;2.
- Eppley, R.W., Stewart, E., Abbot, M.R., and Heyman, V 1985. Estimating ocean primary production from satellite chlorophyll, introduction to regional differences and statistics for the Southern California Bight. *J. Plankton Res.* 7: 57-70.
- Falkowski, P.G. 1981. Light-shade adaptation and assimilation numbers. *J. Plankton Res.* 3: 203-216
- Grebmeier, J. M., J. E. Overland, S. E. Moore, E. V. Farley, E. C. Carmack, L. W. Cooper, K. E. Frey, J. H. Helle, F. A. McLaughlin, and S. L. McNutt . 2006. A major ecosystem shift in the northern Bering Sea, *Science*, 311: 1461-1464, doi:10.1126/science.1121365
- Grebmeier, J. M., S. E. Moore, J. E. Overland, K. E. Frey, and R. Gradinger (2010), Biological response to recent Pacific Arctic sea ice retreats, *Eos Trans. AGU*, 91, 161-168, doi:10.1029/2010EO180001.
- Hare, C. E., K. Leblanc, G. R. DiTullio, R. M. Kudela, Y. Zhang, P. A. Lee, S. Riseman, and D. A. Hutchins (2007), Consequences of increased temperature and CO<sub>2</sub> for phytoplankton community structure in the Bering Sea, *Mar. Ecol. Prog. Ser.*, 352, 9-16, doi:10.3354/meps07182.
- Hill, V., and Cota, G. 2005. Spatial patterns of primary production on the shelf, slope and basin of the western Arctic in 2002. *Deep Sea Research II*, 52: 3344-3354.
- Hirawake, T., Shinmyo, K., Fujiwara, A., and Saitoh, S. 2012. Satellite remote sensing of primary productivity in the Bering and Chukchi Seas using an absorption-based approach. - *ICES Journal of Marine Science*, 69: 1194-1204.
- Hirons, A. C., D. M. Schell, and B. P. Finney. 2001. Temporal records of d<sup>13</sup>C and d<sup>15</sup>N in North Pacific pinnipeds: Inferences regarding environmental change and diet, *Oecologia*, 129: 591-601
- Hunt, G. L., Jr., P. Stabeno, G. Walters, E. Sinclair, R. D. Brodeur, J. M. Napp, and N. A. Bond (2002), Climate change and control of the southeastern Bering Sea pelagic ecosystem, *Deep Sea Res., Part II*, 49, 5821-5853, doi:10.1016/S0967-0645(02)00321-1.
- Liu, J., J. A. Curry, and D. G. Martinson. 2004. Interpretation of recent Arctic sea ice variability, *Geophys. Res. Lett.*, 31, L02205, doi:10.1029/2003GL018732.

- Lee, S. H., T. E. Whitledge, and S. Kang (2007), Recent carbon and nitrogen uptake rates of phytoplankton in Bering Strait and the Chukchi Sea, *Cont. Shelf Res.*, 27, 2231-2249, doi:10.1016/j.csr.2007.05.009.
- Lee, S. H., Joo, H. M., Yun, M. S., and Whitledge, T. E. 2012. Recent phytoplankton productivity of the northern Bering Sea during early summer in 2007. *Polar Biology*, 35: 83-98
- Loeng, H., K. Brander, E. Carmack, S. Denisenko, K. Drinkwater, B. Hansen, K. Kovacs, P. Livingston, F. McLaughlin, and E. Sakshaug. 2005. Marine systems, in *Arctic Climate Impact Assessment*, pp. 453-538, Cambridge Univ. Press, Cambridge, U.K.
- Marra, J., Trees, C. C., and O'Reilly, J. E. 2007. Phytoplankton pigment absorption: a strong predictor of primary productivity in the surface ocean. *Deep Sea Research I*, 54: 155-163.
- Matsuoka, A., Hill, V., Huot, Y., Babin, M., and Bricaud, A. 2011. Seasonal variability in the light absorption properties of western Arctic waters: parameterization of the individual components of absorption for ocean color applications. *Journal of Geophysical Research*, 116: C02007
- Mizobata, K., Shimada, K., Woodgate, R., Saitoh, S., and Wang, J. 2010. Estimation of heat flux through the eastern Bering Strait. *Journal of Oceanography*, 66: 405-424.
- Moore, S. E., and K. L. Laidre. 2006. Trends in sea ice cover within habitats used by bowhead whales in the Western Arctic, *Ecol. Appl.*, 16, 932-944, doi:10.1890/1051-0761(2006)016[0932:TISICW]2.0.CO;2.
- Morel, A. 1978. Available, usable, and stored radiant energy in relation to marine photosynthesis. *Deep-Sea Res.* 25: 673-688.
- Morel, A. 1991. Light and marine photosynthesis: A spectral model with geochemical and climatological implications. *Prog. Oceanogr.* 26: 263-306.
- Niebauer, H. J. 1988. Effects of El Niño? Southern Oscillation and North Pacific weather patterns on interannual variability in the subarctic Bering Sea, *J. Geophys. Res.*, 93, 5051-5068, doi:10.1029/JC093iC05p05051.1877, doi:10.1016/0278-4343(94)00097-7.
- Niebauer, H. J., N. A. Bond, L. P. Yakunin, and V. V. Plotnikov. 1999. An update on the climatology and sea ice of the Bering Sea, in *Dynamics of the Bering Sea*, edited by T. R. Loughlin and K. Ohtani, pp. 29-60, Univ. of Alaska Sea Grant, Fairbanks, Alaska.
- NPO, 2008. Bering Sea. [<http://www.pmel.noaa.gov/np/pages/seas/bseamap.html>]. Accessed on: 11 December 2013.
- Overland, J. E., and P. J. Stabeno. 2004. Is the climate of the Bering Sea warming and affecting the ecosystem?, *Eos Trans. AGU*, 85, 309-316, doi:10.1029/2004EO330001.
- O'Reilly, J., Maritorena, S., Mitchell, B., Siegel, D., Carder, K., Garver, S., Kahru, M., et al. 1998. Ocean color chlorophyll algorithms for SeaWiFS. *Journal of Geophysical Research*, 103: 24937-24953.
- Pabi, S., van Dijken, G. L. and Arrigo, K. R. 2008. Primary production in the Arctic Ocean, 1998-2006, *J. Geophys. Res.*, 113, C08005, doi:10.1029/2007JC004578.
- Platt, T. 1986. Primary production of the ocean water column as a function of surface light intensity. Algorithms for remote sensing. *Deep-Sea Res.* 33: 149-163.
- Schell, D. M. 2000. Declining carrying capacity in the Bering Sea: Isotopic evidence from whale baleen, *Limnol. Oceanogr.*, 45, 459-462, doi:10.4319/lo.2000.45.2.0459.
- Smith, R.C., Eppley, R.W., and Baker, K.S. 1982. Correlation of primary production as measured aboard ship in southern California coastal waters and as estimated from satellite chlorophyll images. *Mar. Biol.* 66: 281-288



- Springer, A. M., and C. P. McRoy. 1993. The paradox of pelagic food webs in the northern Bering Sea? Part III. Patterns of primary production, Cont. Shelf Res., 13, 575-599, doi:10.1016/0278-4343(93)90095-F.
- Stabeno, P. J., J. D. Schumacher, and K. Ohtani. 1999. The physical oceanography of the Bering Sea, in *Dynamics of the Bering Sea*, edited by T. R. Loughlin and K. Ohtani, pp. 1-28, Univ. of Alaska Sea Grant, Fairbanks, Alaska.
- Stabeno, P. J., N. A. Bond, N. B. Kachel, S. A. Salo, and J. D. Schumacher (2001), On the temporal variability of the physical environment over the south-eastern Bering Sea, *Fish. Oceanogr.*, 10, 81-98, doi:10.1046/j.1365-2419.2001.00157.x.
- Stabeno, P. J., Napp, J., Mordy, C., and Whitley, T. 2010. Factors influencing physical structure and lower trophic levels of the eastern Bering Sea shelf in 2005: sea ice, tides, and winds. *Progress in Oceanography*, 85: 180-196.
- Walsh, J. J., and C. P. McRoy. 1986. Ecosystem analysis in the southeastern Bering Sea, *Cont. Shelf Res.*, 5, 259-288, doi:10.1016/0278-4343(86)90018-X.
- Walsh, J. J., et al. (1989), Carbon and nitrogen cycling within the Bering/Chukchi Seas: Source regions for organic matter effecting AOU demands of the Arctic Ocean, *Prog. Oceanogr.*, 22, 277-359, doi:10.1016/0079-6611(89)90006-2
- Woodgate, R. A., Weingartner, T., and Lindsay, R. 2010. The 2007 Bering Strait oceanic heat flux and anomalous Arctic sea-ice retreat. *Geophysical Research Letters*, 37: L01602.