

2024-03

# Planktonic Marine Fungi: A Review

Peng, X

<https://pearl.plymouth.ac.uk/handle/10026.1/22490>

---

10.1029/2023jg007887

Journal of Geophysical Research: Biogeosciences

American Geophysical Union (AGU)

---

*All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.*

# JGR Biogeosciences



## REVIEW ARTICLE

10.1029/2023JG007887

### Key Points:

- Planktonic marine fungi play key roles in the cycling of carbon, nitrogen, phosphorus, sulfur, and metals in the ocean
- There is a large potential for discovering novel lineages and functions of planktonic marine fungi, particularly in the open ocean
- The ecological roles of planktonic marine fungi should be studied by combining multi-omics and biochemical tools

### Correspondence to:

X. Peng,  
xpeng@seoe.sc.edu

### Citation:

Peng, X., Amend, A. S., Baltar, F., Blanco-Bercial, L., Breyer, E., Burgaud, G., et al. (2024). Planktonic marine fungi: A review. *Journal of Geophysical Research: Biogeosciences*, 129, e2023JG007887. <https://doi.org/10.1029/2023JG007887>







Received 1 NOV 2023

Accepted 12 FEB 2024

### Author Contributions:

**Conceptualization:** Xuefeng Peng  
**Formal analysis:** Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe, Virginia P. Edgcomb, Hans-Peter Grossart, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Judith van Bleijswijk, Allison K. Walker, Syrena Whitner  
**Funding acquisition:** Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe, Virginia P. Edgcomb, Hans-Peter Grossart, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Judith van Bleijswijk, Allison K. Walker, Syrena Whitner  
**Investigation:** Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe,

## Planktonic Marine Fungi: A Review

Xuefeng Peng<sup>1</sup> , Anthony S. Amend<sup>2</sup>, Federico Baltar<sup>3</sup> , Leocadio Blanco-Bercial<sup>4</sup> , Eva Breyer<sup>3</sup>, Gaëtan Burgaud<sup>5,6</sup>, Michael Cunliffe<sup>7,8</sup>, Virginia P. Edgcomb<sup>9</sup> , Hans-Peter Grossart<sup>10,11</sup> , Paraskevi Mara<sup>9</sup>, Hossein Masigol<sup>10</sup>, Ka-Lai Pang<sup>12</sup>, Alice Retter<sup>10</sup>, Cordelia Roberts<sup>7,8</sup>, Judith van Bleijswijk<sup>13</sup>, Allison K. Walker<sup>14</sup> , and Syrena Whitner<sup>2</sup>

<sup>1</sup>School of Earth, Ocean and Environment, University of South Carolina, Columbia, SC, USA, <sup>2</sup>Pacific Biosciences Research Center, University of Hawai'i at Mānoa, Honolulu, HI, USA, <sup>3</sup>Department of Functional and Evolutionary Ecology, University of Vienna, Vienna, Austria, <sup>4</sup>Bermuda Institute of Ocean Sciences, Arizona State University, St. George's, Bermuda, <sup>5</sup>Laboratoire Universitaire de Biodiversité et Écologie Microbienne, INRAE, University of Brest, Plouzané, France, <sup>6</sup>Institut Universitaire de France, Paris, France, <sup>7</sup>School of Biological and Marine Sciences, University of Plymouth, Plymouth, UK, <sup>8</sup>Marine Biological Association, The Laboratory, Plymouth, UK, <sup>9</sup>Woods Hole Oceanographic Institution, Woods Hole, MA, USA, <sup>10</sup>Plankton and Microbial Ecology, Leibniz Institute for Freshwater Ecology and Inland Fisheries (IGB), Neuglobsow, Germany, <sup>11</sup>Potsdam University Institute of Biochemistry and Biology, Potsdam, Germany, <sup>12</sup>Institute of Marine Biology and Centre of Excellence for the Oceans, National Taiwan Ocean University, Keelung, Taiwan, <sup>13</sup>Department of Marine Microbiology and Biogeochemistry, NIOZ Royal Netherlands Institute for Sea Research, Texel, The Netherlands, <sup>14</sup>Department of Biology, Acadia University, Wolfville, NS, Canada

**Abstract** Fungi in marine ecosystems play crucial roles as saprotrophs, parasites, and pathogens. The definition of marine fungi has evolved over the past century. Currently, “marine fungi” are defined as any fungi recovered repeatedly from marine habitats that are able to grow and/or sporulate in marine environments, form symbiotic relationships with other marine organisms, adapt and evolve at the genetic level, or are active metabolically in marine environments. While there are a number of recent reviews synthesizing our knowledge derived from over a century of research on marine fungi, this review article focuses on the state of knowledge on planktonic marine fungi from the coastal and open ocean, defined as fungi that are in suspension or attached to particles, substrates or in association with hosts in the pelagic zone of the ocean, and their roles in remineralization of organic matter and major biogeochemical cycles. This review differs from previous ones by focusing on biogeochemical impacts of planktonic marine fungi and methodological considerations for investigating their diversity and ecological functions. Importantly, we point out gaps in our knowledge and the potential methodological biases that might have contributed to these gaps. Finally, we highlight recommendations that will facilitate future studies of marine fungi. This article first provides a brief overview of the diversity of planktonic marine fungi, followed by a discussion of the biogeochemical impacts of planktonic marine fungi, and a wide range of methods that can be used to study marine fungi.

**Plain Language Summary** Fungi are an important yet often overlooked member of marine food webs, especially in the water column. This review highlights the important ecological roles fungi play in the ocean water column, such as the recycling of dead organic matter and acting as parasites of microalgae in the ocean. We provide a summary of state-of-the-art methods for studying planktonic marine fungi and making discoveries in the future.

## 1. Introduction

Fungi in marine ecosystems play crucial roles as saprotrophs, parasites, pathogens, commensals, and symbionts. The definition of marine fungi has evolved over the past century, partially because taxonomic overlap can often be seen with some terrestrial fungi. Currently, “marine fungi” are defined as any fungi recovered repeatedly from marine habitats that are able to grow and/or sporulate in marine environments, form symbiotic relationships with other marine organisms, adapt and evolve at the genetic level, or are active metabolically in marine environments (Pang et al., 2016). Driven by rising interests from multiple disciplines in the past decade, a number of reviews, books, and perspective articles have provided synthesis of our knowledge derived from over a century of research on marine fungi (Amend et al., 2019; Breyer & Baltar, 2023; Burgaud et al., 2022; Cunliffe, 2023; Gladfelter et al., 2019; Gonçalves, Esteves, & Alves, 2022; Grossart et al., 2019; Hassett et al., 2019; Jones & Pang, 2012; Kempken, 2023; Raghukumar, 2017a; Rédou et al., 2016; Richards et al., 2012; Sen et al., 2022). Planktonic marine fungi refer to those (in the form of spores, yeasts, mycelia, sporangia or other fungal propagules) that are

© 2024. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Virginia P. Edgcomb, Hans-Peter Grossart, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Judith van Bleijswijk, Allison K. Walker, Syrena Whitner

**Methodology:** Xuefeng Peng

**Project administration:** Xuefeng Peng

**Resources:** Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe, Virginia P. Edgcomb, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Judith van Bleijswijk, Allison K. Walker, Syrena Whitner

**Supervision:** Xuefeng Peng

**Validation:** Xuefeng Peng

**Visualization:** Xuefeng Peng, Anthony S. Amend, Leocadio Blanco-Bercial, Gaëtan Burgaud, Hans-Peter Grossart, Hossein Masigol, Alice Retter, Syrena Whitner

**Writing – original draft:** Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe, Virginia P. Edgcomb, Hans-Peter Grossart, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Syrena Whitner

**Writing – review & editing:**

Xuefeng Peng, Anthony S. Amend, Federico Baltar, Leocadio Blanco-Bercial, Eva Breyer, Gaëtan Burgaud, Michael Cunliffe, Virginia P. Edgcomb, Hans-Peter Grossart, Paraskevi Mara, Hossein Masigol, Ka-Lai Pang, Alice Retter, Cordelia Roberts, Judith van Bleijswijk, Allison K. Walker, Syrena Whitner

found to be active in suspension or attached to particles, substrates or hosts in the ocean (G. Wang et al., 2012). This review builds upon previous reviews by focusing on the roles planktonic marine fungi play in ocean biogeochemistry and by providing summaries of state-of-the-art methods for investigating the diversity, function, and activity of marine fungi. Finally, we highlight recommendations that will facilitate future studies of marine fungi.

This article first provides a brief overview of the diversity of planktonic marine fungi, followed by a discussion of the ecological roles and biogeochemical impacts of planktonic marine fungi, and a wide range of methods that can be used to study marine fungi.

## 2. Diversity of Planktonic Marine Fungi


### 2.1. Diversity of Planktonic Marine Fungi in Coastal Waters

Coastal regions constitute areas where land masses meet the ocean and vary from ocean and inland systems in terms of precipitation patterns, humidity, food web dynamics, and input of organic matter (Ketchum, 1972). Coastal regions are highly dynamic, complex, and productive systems that host about 40% of the human population within a range of 100 km from the shoreline (Baztan et al., 2015). Near-shore marine ecosystems are not as stable as their open ocean counterparts and can vary in input of terrestrial organic matter, sediment, nutrients, and pollutants, as well as changes in salinity due to evapotranspiration and freshwater mixing (Clipson et al., 2005; Ward et al., 2017).

Coastal planktonic fungi typically constitute a mixture of marine and terrestrial lineages (Christmas et al., 2023). There are currently 1,898 described species of marine fungi (<https://www.marinefungi.org/>, 12 September 2023; Calabon et al., 2023), but their actual diversity is estimated to be much higher, with some studies stating that only around 1% of marine fungi have been identified (Gladfelter et al., 2019; Jones, 2011). In coastal waters, planktonic fungi are commonly found on aggregates or detritus associated (Gutiérrez et al., 2010, 2020). Moreover, they can be associated with other eukaryotes or phytoplankton as pathogens or symbionts (Grossart et al., 2016; Gutiérrez et al., 2010; Raghukumar, 2017a; Richards et al., 2015; Sen et al., 2022).

Fungi can successfully thrive in aquatic ecosystems due to their unique physiological and biochemical traits (Sridhar, 2020). Seasonal changes, in for example, water temperature, pH, and chlorophyll-a, are mainly responsible for driving fungal community composition and for observed uneven distribution across coastal waters and time (Priest et al., 2021). Highest diversities are found closest to the shoreline, and decreased toward the open ocean (Debeljak & Baltar, 2023; Duan et al., 2021; Z. Gao et al., 2010; Gutiérrez et al., 2010; Sen et al., 2021). Close to shore, proximity and connectivity to estuaries seem to be determining factors in observed high diversity, probably due to directly linked input of fungi from the inflowing rivers (Debeljak & Baltar, 2023; Taylor & Cunliffe, 2016). Moreover, eutrophication has an impact on the assembly and composition of the most abundant fungal taxa along a eutrophication gradient. After environmental disturbance, however, assembly mechanisms and adaptation strategies differed between abundant and rare fungal taxa (H. Zhao et al., 2023). This study also highlighted the essential role of rare taxa in fungal community structure, stability, and diversity in coastal waters. Additionally, the authors emphasized the important biogeochemical role planktonic fungi play in the cycling of carbon, nitrogen, and phosphorus in coastal ecosystems, predominantly via their extracellular enzyme activities. The fungal communities recovered from coastal waters were mostly dominated by members of Dikarya (Ascomycota and Basidiomycota), including genera such as *Cystobasidium*, *Rhodotorula*, *Aspergillus*, *Penicillium*, and *Cladosporium*, as well as some early diverging lineages including Chytridiomycota (Pham et al., 2021; Sen et al., 2021).

For a worldwide meta-analysis of fungal 18S rRNA gene (V4 region) amplicon data from the water column exclusive to coastal regions, we retrieved information from the publicly available MetaPR2 database (Vaultot et al., 2022; Figures 1a and 1b). Sample numbers were significantly different between the southern ( $N = 99$ ) and the northern hemisphere ( $N = 730$ ) (Chi-squared test,  $p < 0.001$ ), indicating that coastal habitats south of the equator are still largely understudied (Figure 1c). In the southern hemisphere, coastal fungal communities were mainly composed of Basidiomycota (72.3%), Ascomycota (14.1%), and unclassified fungi (10.6%), whereas Ascomycota (36.6%), Chytridiomycota (33.4%), and Basidiomycota (19.3%) dominated the community in the northern hemisphere (Figures 1a and 1b). The limited number of southern hemisphere samples were primarily collected during active phytoplankton blooms (e.g., Trefault et al., 2021), so there is a seasonal bias in the existing



**Figure 1.** Global distribution of fungi and samples across coastal areas depicting their composition at phylum level, as well as oceanic regions retrieved from the MetaPR2 database (a). Fungal community composition on phylum level (colored according to a) between oceanic regions showing the dominant phyla and dominant families in parenthesis (rel. abundance >0.3) (b), where the bar colors correspond to the legend shown in the upper part of panel (a). Distribution of the number of samples between oceanic regions indicating potential under-sampling of specific coastal regions (c). Note that the average salinity of the Baltic Sea (7.4 practical salinity units) is lower than the oceanic average (Janssen et al., 1999).

data set. Interestingly, relative abundances of chytrids correlated with increasing temperatures, especially in the coastal regions of the North Pacific Ocean (Spearman's Rho = 0.54,  $p < 0.01$ ) and North Atlantic Ocean (Spearman's Rho = 0.17,  $p < 0.01$ ). In the North Atlantic Ocean, Sordariomycetes were more abundant in colder water temperatures (Spearman's Rho =  $-0.12$ ,  $p < 0.05$ ). In the Southern Ocean, where water temperatures were generally lowest (mean  $T = 0.8 \pm 5.6^\circ\text{C}$ ), the relative abundance of fungi was higher in warmer regions. The opposite was true for Cystobasidiomycetes (Spearman's Rho =  $-0.42$ ,  $p < 0.01$ ) and Tremellomycetes (Spearman's Rho =  $-0.30$ ,  $p < 0.01$ ), of which the relative abundance was higher at lower temperatures.

In conclusion, members of the phylum Chytridiomycota often are one of the dominating fungal lineages in coastal regions worldwide, which is consistent with findings in Debeljak and Baltar (2023). Yet, Chytridiomycota are understudied, hence they often contribute to the fungal "dark matter" (Grossart et al., 2016). Partly, this is due to the usage of primer pairs known to have a taxonomic bias toward Dikarya (Tedersoo et al., 2015), and to the incompleteness of publicly available databases often failing to classify early diverging fungal lineages to any ecologically meaningful taxonomic level. In addition, coastal waters in the southern hemisphere are predicted to harbor a larger number of undescribed species, suggesting a potentially high and untapped fungal diversity in these understudied regions.

## 2.2. Diversity of Planktonic Marine Fungi in the Open Ocean

There are fewer studies focusing on planktonic fungi in the open ocean compared to coastal regions, and even fewer have targeted the mesopelagic and below. This discrepancy can be attributed to logistical challenges associated with reaching the open ocean and the assumption that the open ocean offers fewer substrates for fungi to proliferate. Nearly all cultivation-based studies of marine fungi collected inocula from coastal and estuarine environments (Jones & Pang, 2012). One exception is an early cultivation-based study that collected samples in the Indian Ocean along the 60°E meridian from 12°N to 41°S at depths of 0–2,000 m (Fell, 1967). This study found that the density of yeast cells in the Indian Ocean was low ( $0\text{--}513 \text{ cells L}^{-1}$ ), and *Rhodotorula* (Basidiomycota) and *Candida* (Ascomycota) were cosmopolitan yeast genera that were frequently cultivated (Fell, 1967). The tendency to cultivate yeasts from the open ocean is consistent with their apparent dominance (Bass et al., 2007), which is attributed to their large surface-to-volume ratio and flexible physiology (Kutty & Philip, 2008). Currently there are no known early diverging fungi cultivated from the open ocean.

Despite the widespread application of metabarcoding surveys of marine microbial communities, the number of studies that examined open ocean fungal diversity remains very small. A review by Breyer and Baltar (2023) provided an up-to-date and comprehensive summary of metabarcoding surveys of planktonic fungi. Of the 26 studies summarized by Breyer and Baltar (2023), only 11 included samples from the open ocean (Bass et al., 2007; Debeljak & Baltar, 2023; Hassett et al., 2017, 2019; Jeffries et al., 2016; W. Li et al., 2019; Morales et al., 2019; Orsi et al., 2022; Peng & Valentine, 2021; Tisthammer et al., 2016; X. Wang et al., 2014). Most of these studies focused on one region of the global ocean during one field campaign, while TARA Oceans was the primary source of fungal diversity data providing larger coverage of the global ocean.

While Ascomycota and Basidiomycota consistently account for over two thirds (often  $>80\%$ ) of the fungal community in the open ocean detected using molecular approaches, there appears to be a systematic difference between studies depending on the primer set used for metabarcoding. The relative abundance of early diverging fungi, particularly Chytridiomycota and Mucoromycota, was usually very low ( $<1\%$ ) when a primer set targeting the ITS2 region was used (e.g., W. Li et al., 2019; Peng & Valentine, 2021), but when the V4 region of the 18S rRNA gene or the ITS1 region was the target, the relative abundance of early diverging fungi can reach  $>10\%$  in the open ocean (Debeljak & Baltar, 2023; Jeffries et al., 2016; Orsi et al., 2022). Regardless of the type of primer used, the relative abundance of unclassified fungal taxa, which potentially represent novel lineages, was the highest in the open ocean when compared to coastal and estuarine environments (Debeljak & Baltar, 2023; W. Li et al., 2019; Peng & Valentine, 2021).

While meta-analysis of existing data sets has generated insights into patterns of fungal diversity in the open ocean (e.g., Debeljak & Baltar, 2023), most of the large-scale surveys (e.g., TARA Oceans) were not designed to target fungi, so the primers used were typically universal 18S rRNA gene primers that result in small numbers (e.g., tens to hundreds) of fungal reads per sample due to the low abundance of fungal cells in open ocean microbial communities and potential primer biases. The sequencing depth in these previous studies was likely insufficient for estimating the diversity of planktonic marine fungi. On the other hand, effects of airborne and/or human skin

associated fungi contamination during sampling collection and handling can be relatively large unless appropriate controls are in place to minimize contaminations (Marcelino et al., 2019). Regardless of which region of the rRNA gene is targeted (18S, internal transcribed spacer [ITS], or 28S), using primers specifically designed to maximize the coverage of fungi (e.g., Banos et al., 2018; Tedersoo et al., 2015; Wurzbacher et al., 2019) will be crucial to advance our understanding of fungal diversity in the open ocean.

### 3. Biogeochemical Impacts of Planktonic Marine Fungi

#### 3.1. Carbon Cycling

##### 3.1.1. Particle-Associated Fungi

Particulate organic matter (POM) is a central component of the biological carbon pump (BCP), which sequesters atmospheric carbon dioxide and facilitates its transport to the deep ocean (Falkowski et al., 1998). As POM sinks, its molecular composition transforms via heterotrophic degradation, supplying bioavailable nutrients to a broad microbial community (Azam et al., 1983; Worden et al., 2015). Until recently, bacteria and archaea were considered the sole biological decomposers of POM and drivers of the “microbial loop” (Baumas et al., 2021; Duret et al., 2019; Leu et al., 2022; Ollison et al., 2021). However, recent investigations in regions characterized by coastal upwelling demonstrate the prevalence, and at times the dominance, of saprotrophic fungi (and/or fungi-like organisms such as Labyrinthulomycetes) closely associated with POM, herein referred to as “particle-associated fungi” (Duret et al., 2020; Gutiérrez et al., 2011).

Particle-associated fungi are distinct from free-living fungi and are characterized by their attachment to various particles found in the marine environment. These particles encompass a range of substrates such as detritus (i.e., microbial necromass, fecal pellets) and microplastics (Y. Yang et al., 2020). Particle-associated fungal communities can be sampled using a marine snow catcher (Duret et al., 2020) or by using size-fractionated filtration (Peng & Valentine, 2021), but most studies of marine fungi use only a single filter size, which would capture both free-living and particle-associated communities (Morales et al., 2019). Particle-associated fungi are generally dominated by Ascomycota (primarily Dothideomycetes) and Basidiomycota (primarily Microbotryomycetes and Exobasidiomycetes) across the majority of depths and in both coastal and pelagic regions (Duret et al., 2020; Peng & Valentine, 2021).

Analogous to their terrestrial and freshwater counterparts, marine fungi possess an arsenal of extracellular enzymes for metabolizing biopolymers, including chitinases, laccases, cellulases, amylases, and pectinases (Bonugli-Santos et al., 2010; Kamei et al., 2008; Pointing & Hyde, 2000; Y. Wang et al., 2016). Additionally, marine fungi have demonstrated their ability to process algal derived polysaccharides and POM (Cunliffe et al., 2017; Y. Wang et al., 2016). Fungal roles in lipid, amino acid, and carbohydrate metabolism, are depth-dependent, as reflected in the distribution of fungal carbohydrate-active enzymes (CAZymes) spanning surface waters to the seafloor (Baltar et al., 2021; Cunliffe et al., 2017; Orsi et al., 2013, 2022). Glycosyl hydrolases involved with chitin degradation have been identified (Christmas & Cunliffe, 2020; Orsi et al., 2022), suggesting that fungi may have a role in degrading non-phytoplankton-derived POM such as those derived from copepods which form an important vector for carbon flux in the mesopelagic (Bradford-Grieve et al., 2001; Kobari et al., 2008). During periods of heightened productivity, particularly in coastal upwelling zones characterized by phytoplankton blooms, the abundance of marine fungi correlates positively with the production of hydrolytic enzymes (Gutiérrez et al., 2011). Dikarya fungi effectively assimilate <sup>13</sup>C-labeled algal transparent exopolymer particles (TEP) or substances (TEP or EPS) (Cunliffe et al., 2017; Orsi et al., 2022), providing direct evidence that saprotrophic fungi in marine environments use algal-derived organic matter. Recent evidence based on the comparison of the transcriptomic profiles of fungal CAZymes (Baltar et al., 2021; Orsi et al., 2022) and peptidases (Breyer et al., 2022) to the prokaryotic ones (Z. Zhao et al., 2020) indicate that oceanic prokaryotes and fungi might work together to degrade oceanic organic matter (Orsi et al., 2022).

Although the presence of specific fungal hydrolytic enzymes retrieved from the water column suggests the preference for a particle-associated lifestyle, studies of fungal community composition and metabolic activity on marine POM directly are scarce. Moreover, it remains uncertain whether marine fungi contribute to the formation and stabilization of particles or they delay and reduce aggregation of particles in the water column. The former has been shown in marine sediments (Damare et al., 2008), while the latter has been demonstrated in freshwater environments (Klawonn et al., 2021; Klawonn, Van den Wyngaert, et al., 2023). The potential for planktonic

marine fungi to control organic matter fluxes in the ocean are critically important to consider when examining efficiency and microbial controls of the BCP.


### 3.1.2. Microalgae-Associated Fungi

Microscopic algae and photosynthetic bacteria, more commonly referred to as phytoplankton, are the primary producers that support marine ecosystems globally. Marine phytoplankton such as diatoms and dinoflagellates, including widespread bloom-forming taxa, may become infected by fungal parasites, particularly by the zoosporic Chytridiomycota (“chytrids”) (Gutiérrez et al., 2016; Hassett & Gradinger, 2016; Lepelletier et al., 2014; Scholz et al., 2016; Sparrow, 1969). Interactions between phytoplankton and parasitic fungi influence the health and dynamics of phytoplankton populations directly, through parasitism, and indirectly by impacting the flow of organic matter within and among microbial communities (Grossart et al., 2019; Sime-Ngando, 2012; Van den Wyngaert et al., 2022). Despite their ecological importance, parasitic, microalgae-associated marine fungi remain understudied in comparison to other microbial groups, especially with regard to implications for ocean biogeochemistry (Ilicic & Grossart, 2022).

The diversity and distribution of chytrids is constrained by both biotic and abiotic factors including presence of putative host species, temperature, pH, light intensity, and salinity (Duan et al., 2018; Gutiérrez et al., 2016; Ibelings et al., 2004; Taylor & Cunliffe, 2016). Chytrids demonstrate strong seasonality in coastal environments where variations in water column nutrient composition, oxygen levels, and microbial composition are tightly synchronized with the timing of seasonal phytoplankton blooms (Hassett et al., 2019; Taylor & Cunliffe, 2016). Chytrid-microalgae interactions are generally considered “host-specific,” with several studies utilizing cross-infection experiments to determine generalist versus specialist tendencies (Kagami et al., 2021; Reñé et al., 2023; Van den Wyngaert et al., 2022). However, there are comparatively few studies where host range is examined empirically, in situ within marine systems (Gleason et al., 2011; Kagami, de Bruin, et al., 2007; Kagami, von Elert, et al., 2007). Potential host specificity can be deduced through various means, including analyzing the proportion of attached and detached zoospores in relation to the phytoplankton community composition. For instance, the strong correlation between large, colonial diatoms, specifically *Thalassiosira* and *Skeletonema*, and attached fungal sporangia in the upwelling ecosystem off central Chile demonstrated the seasonality and host preferences of chytrids (Gutiérrez et al., 2016). Subsequently, a shift in diatom community composition, dominated by the genus *Chaetoceros*, coincided with decreased attached sporangia and increased free-swimming chytrid zoospores (Gutiérrez et al., 2016).

During phytoplankton infection, chytrids liberate often inedible biomolecules, including their host cells, to heterotrophic grazers via osmotrophic, extracellular degradation, a concept incorporated into the “mycoloop” (Kagami et al., 2014). In the mycoloop, chytrid zoospores, rich in essential fatty acids, also serve as a high-quality food source for zooplankton like *Daphnia* in freshwater systems (Kagami, de Bruin, et al., 2007; Kagami, von Elert, et al., 2007). Moreover, chytrid infections may facilitate wider ecosystem functioning upon release of nutrients in surface waters after the lysis of host cells (Kagami et al., 2006). Recent work in lakes using both the *Synedra-Zygophlyctis* model system and field samples demonstrated that chytrid infections reduced the formation of aggregates, in part by promoting bacterial colonization of parasitized phytoplankton cells (Klawonn, Van den Wyngaert, et al., 2023). Whether such interactions occur in the ocean remains unknown.

A timely question is if, and how, climate change will impact chytrid-phytoplankton interactions. Ocean warming and ocean acidification directly alter phytoplankton physiology, size structure, and community composition (Basu & Mackey, 2018; Sommer et al., 2012), and they may also indirectly affect marine phytoplankton by promoting chytrid infections. For example, increased light penetration resulting from reduced sea ice coverage in Arctic waters may lead to photoinhibition and physiological stress in diatoms, creating favorable conditions for disease proliferation (Hassett & Gradinger, 2016). Kiliyas et al. (2020) demonstrated that chytrids were positively correlated with the extent of sea ice melting, implying that under periods of prolonged light exposure, due to a reduction in seasonal sea ice coverage, the rate of fungal infection may increase. It is unclear if these patterns extend to other marine systems, but they suggest the possibility that rates of chytrid infection may increase in the future. Given the role of phytoplankton at the base of most marine food chains, this might impact food quality and abundance at higher trophic levels, the composition of photosynthetically derived organic matter released, and even overall global nutrient cycling (Rasconi et al., 2020; Sánchez Barranco et al., 2020; Senga et al., 2018).



**Figure 2.** Literature review of 13 studies analyzing the Plastisphere on marine plastic debris with a focus on fungal communities (updated from Wright et al., 2020). Study design highlights whether the study was conducted in a laboratory or in the field (yellow), the different types of plastics, that is, conventional or biodegradable (green), and whether the plastics were naturally collected from the sea (collection) or introduced by the researchers and collected either at a single time point or at a series of time points (blue). Numbers indicate the first and last days of incubation, with numbers in brackets indicating the number of points included in the time series. Fungal community characterization (purple) indicates whether the Plastisphere was analyzed via 18S or internal transcribed spacer (ITS) rRNA high throughput sequencing or by a culture-based approach, and whether microscopy was used to visualize the biofilm. The controls (orange) highlight those studies that compare the Plastisphere with the microbial community of the surrounding seawater (either bulk or between 0.2 and 3  $\mu\text{m}$ ) or particulate organic matter (>3  $\mu\text{m}$ ), or biofilms that develop on inert surfaces (e.g., glass, rock, wood, etc.). Meaning of polymer acronyms: PE, Polyethylene; PP, Polypropylene; PET, Polyethylene terephthalate; PS, Polystyrene; HDPE, High-density polyethylene; LDPE, Low-density polyethylene; SAN, Styrene-Acrylonitrile; PA, Polyamide; PU, Polyurethane; CA, Cellulose Acetate; PVC, Polyvinyl chloride; PLA, Polylactic acid; PCL, Polycaprolactone; PHBV, Poly(3-hydroxybutyrate-co-3-hydroxyvalerate). PLA has been enclosed in brackets due to its biodegradability under specific conditions. \* The publication by Zettler et al. (2013) was included, even though it was not focused on fungi, because it represents the first publication indicating the presence of microeukaryotes on marine plastic debris.

### 3.1.3. Plastic-Associated Fungi

Fragments of plastic waste that find their way into the oceans swiftly become enveloped by organic matter, giving rise to an “ecocorona,” a distinctive layered structure that modifies the hydrophobic properties of the plastic surfaces and facilitates the adherence and establishment of microbial communities (Latva et al., 2022; Wright et al., 2020; Y. Yang et al., 2020). These communities, referred to as the “plastisphere” (Zettler et al., 2013), promptly establish intricate biofilms, typically within minutes to hours (Latva et al., 2022), reaching a mature state within 15–30 days (Cheng et al., 2021; Kirstein et al., 2018; Odobel et al., 2021). Extensive research in the field of prokaryotic communities' colonization dynamics has shown that over large spatial scales environmental factors, rather than the type of polymer, play a pivotal role in shaping the composition of the plastisphere (Wright et al., 2021). However, different types of polymers that were locally incubated (Zettler et al., 2013) or harvested from the same field location (Vaksmas et al., 2021), showed different microbial communities.

Here we have reviewed primary research articles dedicated to elucidating fungal communities associated with marine plastics (Figure 2), and we acknowledge another recent review highlighting the potential role marine fungi play in plastic degradation (Zeghal et al., 2021). Planktonic marine fungi associated with plastics do not necessarily degrade plastics, and only a few fungal lineages have been shown to degrade plastics. *Zalerion maritimum* (Paço et al., 2017), *Alternaria alternata* FBI (R. Gao et al., 2022) and *Rhodotorula mucilaginosa* (Vaksmas et al., 2023) have been identified to degrade polyethylene (PE) and *Cladosporium halotolerans* 6UPA1 was shown to degrade polyurethane (K. Zhang et al., 2022). Our investigation revealed that the study of plastic-



associated fungi, primarily conducted through a metabarcoding approach, constitutes an emerging field, as evidenced by 6 out of 13 relevant studies published between 2020 and 2023. For those that included a comparable seawater control, plastic-associated fungal communities were significantly distinct from fungal communities in the surrounding bulk seawater. Interestingly, a specific study delved into fungal communities inhabiting marine sediments and associated plastics, employing a culture-based methodology (Florio Furno et al., 2022). This investigation yielded findings consistent with earlier research, demonstrating disparities in species richness between plastic samples and sediment or water samples. Among the selected research articles featuring an inert control material such as wood or glass, three studies (Kettner et al., 2017, 2019; Kirstein et al., 2018) reported notable distinctions between the control and plastic substrates. However, in one study (Oberbeckmann et al., 2016) specifically examining fungal communities colonizing polyethylene terephthalate (PET), no significant differences were observed when compared to those colonizing glass substrates. These findings imply that geographical locations and inherent environmental variables may exert a stronger influence on fungal colonization than the type of plastic material.

Most published studies indicate that the polymeric composition of plastics does not exert significant influence on fungal operational taxonomic unit (OTU)/amplicon sequence variant (ASV) richness or community composition (Kettner et al., 2017, 2019; Lacerda et al., 2020, 2022; Sérvulo et al., 2023). Only one study detected differences in community composition, of fungal communities on polystyrene and PET, which contrasted with those on polyurethane (S. Yang et al., 2022). A recent study comparing fungal colonization on conventional versus biodegradable plastics underscores the significant impact of immersion duration on fungal colonization (Philippe et al., 2023). This is supported by temporal variability observed in fungal communities (De Tender et al., 2017). However, these results diverge from those of Sérvulo et al. (2023), who reported stable communities over a 1-year immersion period. In terms of taxonomic diversity, Ascomycota dominate plastic-associated fungal communities, where other fungal phyla are also present (De Tender et al., 2017; Kettner et al., 2017; Lacerda et al., 2020, 2022; Philippe et al., 2023; Sérvulo et al., 2023; Zeghal et al., 2021), which aligns with the general patterns in pelagic marine fungal diversity (Jones et al., 2019). Moreover, plastic-associated Chytridiomycota exhibit higher abundance during the autumn and winter seasons (Sérvulo et al., 2023) or at higher latitudes (Kettner et al., 2017; Kirstein et al., 2018).

These findings reveal emerging trends and distribution patterns of plastic-associated fungi. However, it is crucial to interpret these preliminary insights with caution due to the diverse array of plastic formats (plates, sheets, pellets, etc.) used in the highlighted studies, putative variations due to the presence of additives in the plastic matrices, distinct environmental conditions, and differences in immersion durations across these investigations. This collective variability raises significant questions about the development of a comprehensive understanding of fungal colonization of plastics. The study of plastic-associated fungi is further complicated by interactions with bacterial and protist communities, as well as the influence of stochastic processes during initial colonization stages. This underscores the need for standardized research to gain additional insights into whether plastics genuinely constitute a unique habitat for fungi.

### 3.1.4. Fungi at the Ocean-Atmosphere Interface

The ocean-atmosphere interface (or sea surface microlayer, SML) covers approximately 71% of the Earth's surface and is the site at which all substances entering or leaving the ocean must pass through (Engel et al., 2017). The combined physical, chemical, and biological properties of the SML impact ocean-atmosphere exchange processes, including those that are part of the marine carbon cycle (Hardy, 1982). Ocean-atmosphere CO<sub>2</sub> flux is globally important, as about 30% of anthropogenic CO<sub>2</sub> emissions since the beginning of the industrial revolution was taken up by the ocean (Gruber et al., 2019).

Microbial life in the SML, often referred to as the neuston, can be sampled using the mesh screen sampler (Agogué et al., 2004), and bacteria in the neuston community (i.e., bacterioneuston) are relatively well-studied (Cunliffe et al., 2008; Franklin et al., 2005; Wurl & Holmes, 2008). Neuston communities can modify air-sea gas transfer directly, either through the degradation of specific gases (e.g., methane, carbon monoxide) (Conrad & Seiler, 1988; Frost, 1999; Upstill-Goddard et al., 2003) or general microbial activity impacting net CO<sub>2</sub> and O<sub>2</sub> flux (Reinthal et al., 2008), or indirectly by creating and modifying SML materials that impact the physicochemical properties of the SML, such as surfactants, that impact gas exchange (Kurata et al., 2016). Biological surfactants on the surface of the Atlantic Ocean caused up to a 32% reduction of air-sea CO<sub>2</sub> exchange likely

through turbulence repression (Kurata et al., 2016). Enhanced microbial activity in the SML, particularly associated with slicks, can also modify the physicochemical properties of the “cool ocean surface skin” (Wurl et al., 2018), which also likely impacts CO<sub>2</sub> exchange (Watson et al., 2020).

Fungi in the SML (i.e., myconeuston) are so far very poorly studied. Environmental DNA (eDNA)-based assessment of myconeuston diversity has been conducted in coastal marine waters in the Western English Channel (Taylor & Cunliffe, 2014) and in the open waters of the Mediterranean Sea (Zäncker et al., 2021), with both studies showing that fungal diversity in the SML is distinct from the underlying water column. In the Mediterranean Sea study, particularly for samples collected from the Ionian Sea, fungi were a major eukaryote group in the SML, with the dominant fungal taxon in the study belonging to the genus *Cladosporium*. Other studies have shown, using stable isotope probing, that marine *Cladosporium* actively degrade phytoplankton-produced high-molecular-weight polysaccharides that form gels (Cunliffe et al., 2017), indicating that myconeuston may have a role in processing SML organic material (Zäncker et al., 2021) and therefore impact the physicochemical composition of the ocean-atmosphere interface. These two studies are based on eDNA assessments of myconeuston diversity and therefore only show the presence of fungi in these locations. Currently, we have no information on general (e.g., respiration) or specific (e.g., polysaccharide processing) fungal activity in the SML. Thus we are unable to establish the significance of myconeuston on ocean-atmosphere exchange.

### 3.2. Nitrogen Cycling

Nitrogen (N) is the primary limiting nutrient in many parts of the ocean (Gruber, 2008). Many planktonic microorganisms possess extremely high affinity for fixed inorganic N such as ammonium and nitrate (Martens-Habbena et al., 2009; Mulholland & Lomas, 2008). Because fungal cells typically have a smaller surface-to-volume ratio, planktonic fungi may be at disadvantage when competing with prokaryotes for dissolved inorganic nitrogen such as ammonium and nitrate. On the other hand, aquatic fungi demonstrate highly plastic C:N ratios in their cellular biomass (Danger & Chauvet, 2013; Danger et al., 2016), which may allow them to better survive in ocean biomes where nitrogen is the limiting nutrient (Moore et al., 2013). Additionally, fungi can fulfill their N requirements by degrading organic matter (Hodge & Fitter, 2010), which may be an important source of fixed N for oceanic fungi. A recent study, assigning fungal peptidase-like sequences in the TARA oceans metagenomes and metatranscriptomes, found that fungi express genes responsible for protein degradation throughout the water column in the world's ocean (Breyer et al., 2022). As marine bacteria also assimilate dissolved organic nitrogen (Bronk & Glibert, 1993; Korth et al., 2012), there is potentially competition between marine fungi and bacteria for dissolved organic nitrogen, but this hypothesis still requires experimental verification. In sum, marine fungi play important roles in the recycling and assimilation of fixed organic nitrogen.

Besides their roles in fixed nitrogen assimilation and remineralization, fungi could also play an important role in the dissimilatory cycling of nitrogen in the ocean. Specifically, many fungal species can perform incomplete denitrification, the dissimilatory reduction of nitrate sequentially to nitrite, nitric oxide, and nitrous oxide (Maeda et al., 2015). While fungal denitrification is an important process in soil (H. Chen et al., 2014; Laughlin & Stevens, 2002), this process has been reported in marine environments (Lazo-Murphy et al., 2022; Su et al., 2021; Wankel et al., 2017), including the eastern tropical North Pacific oxygen minimum zone (Peng & Valentine, 2021). On one hand this suggests fungi make a small (up to 10%) but significant contribution to fixed nitrogen loss from the ocean water columns. On the other hand, because nitrous oxide (N<sub>2</sub>O) is a potent ozone-depleting greenhouse gas (Ravishankara et al., 2009) and no fungi have been identified to reduce N<sub>2</sub>O to dinitrogen gas (Shoun & Fushinobu, 2016), fungi may make a disproportionately large contribution to N<sub>2</sub>O emission from the ocean, especially as fungal denitrification appeared to be less sensitive to oxygen inhibition (Peng & Valentine, 2021; Phillips et al., 2016). The apparent lack of sensitivity to oxygen of fungal N<sub>2</sub>O production in a number of tested strains (Phillips et al., 2016; Zuo et al., 2023) suggests that if these strains were present and active in the oxygenated surface ocean, they have the potential to contribute to N<sub>2</sub>O efflux from the ocean.

The diagnostic gene for fungal denitrification is an unconventional nitric oxide reducing cytochrome P450 (*P450<sub>nor</sub>*) likely acquired by fungi via horizontal gene transfer from Actinobacteria (W. Chen et al., 2014; Shoun & Tanimoto, 1991). Although fungal N<sub>2</sub>O production has been demonstrated by many strains (Jirut, 2015; Maeda et al., 2015), a recent study using comparative genomics analyzed >700 fungal genomes and challenged the paradigm that anaerobic respiration is responsible for N<sub>2</sub>O production by fungi. By identifying

genes associated with secondary metabolisms in the vicinity of *P450nor* in a subset (up to a third) of the analyzed fungal genomes, Higgins et al. (2018) suggested that secondary metabolism might be responsible for fungal N<sub>2</sub>O production. However, the fungal genomes analyzed by Higgins and colleagues were nearly exclusively from terrestrial environments. To understand the mechanisms behind fungal denitrification in the ocean, there is an urgent need to conduct studies using fungal strains isolated from the ocean, particularly oxygen minimum zones.

### 3.3. Other Elemental Cycling

#### 3.3.1. Sulfur

Marine fungi participate in sulfur (S) cycling primarily via sulfate assimilation to S-bearing amino acids (Morales et al., 2019; Sen et al., 2021). Nonetheless, extracellular mineralization of organic S (e.g., algal sulfate esters) by pelagic fungi (Ascomycota and Basidiomycota), such as the hydrolysis of sulfate esters present in algal cell walls, was recently reported in studies of fungal isolates from the Atlantic Ocean and the Antarctic Peninsula (Salazar Alekseyeva et al., 2022). The activity (hydrolysis rate and half-saturation constant) of observed enzymes was temperature- and species-dependent, and agrees with previous studies that described fungal sulfatases as thermo-sensitive enzymes with an optimal temperature of 25°C (Korban et al., 2017). The impact of temperature on S-related enzymes suggests that global ocean warming may alter (in some cases increase) the contribution of pelagic fungi to marine S cycling. Other fungal S-hydrolyzing enzymes that can contribute to mineralization of S-bearing compounds in the ocean include putative metallopeptidases that cleave or modify dimethylsulfoniopropionate (DMSP) to dimethyl sulfide (DMS) (Todd et al., 2009). DMSP is an abundant organosulfur compound in the marine environment (i.e., produced by marine bacteria and phytoplankton (Yoch, 2002) and plays a major role in the global S cycle and in marine food webs (Mahajan et al., 2015; Teng et al., 2021)). Marine Ascomycota taxa are known to hydrolyze DMSP (Bacic & Yoch, 1998) using homologs to the *dddP* gene that modify DMSP to DMS and were likely acquired via horizontal gene transfer from bacteria (Todd et al., 2009).

The role of pelagic fungi in S cycling and S mineralization may extend to the breakdown of polycyclic organosulfur hydrocarbons found in oil spills or anthropogenic S compounds that enter marine waters as detergents, pesticides, or pharmaceuticals. Assimilation of complex organosulfur hydrocarbons by fungi depends on desulphurization (breakdown) of the C–S bond (Linder, 2018). Terrestrial basidiomycetes and ascomycetes are reported to desulphurize polycyclic organosulfur compounds using aromatic peroxygenases and cytochrome P450 monooxygenases (e.g., Piontek et al., 2013); however, polycyclic organosulfur degradation by pelagic fungi is not yet fully understood. Oil spills are shown to result in marine sediment eukaryotic communities almost exclusively dominated by fungal taxa (Bik et al., 2012). Oil spills also change the structure and function of pelagic fungal communities, increasing the abundance of presently unclassified fungi within the mycoplankton that have an uncharacterized role in degradation of hydrocarbons (Neethu et al., 2019). Nonetheless, the overall ability of pelagic fungi to degrade hydrocarbons is related to the toxicity of hydrocarbons and the ability of specific fungal lineages to tolerate this toxicity. In the case of oil spills, the abundance of *Candida* and *Rhodotorula* decreased (Neethu et al., 2019).

Pelagic marine fungi may also contribute to S cycling through the production of S-bearing secondary metabolites; however, this is still an area of active research. Production of secondary metabolites can enhance survival and is a strategy utilized by many microorganisms, in particular those with particle-associated lifestyles (e.g., Geller-McGrath et al., 2023). Marine fungi can produce a suite of different S-bearing secondary metabolites with antimicrobial and cytotoxic properties (e.g., Julianti et al., 2022; Liu et al., 2022). These S-bearing secondary metabolites were isolated primarily from deep-sea fungi and taxa that inhabit sediments. Whether these compounds are also produced by pelagic fungi remains to be investigated, however it is likely we will find that they play a role in competition with co-colonizing bacteria on sinking particles. Such S-bearing metabolites could allow fungi to couple sulfur cycling with defense mechanisms against prokaryotes and other fungi that colonize particles.

#### 3.3.2. Phosphorus

Inorganic phosphorus (Pi) is an important macronutrient utilized by marine microorganisms for the synthesis of macromolecules (DNA, RNA, proteins) and under conditions where it is in limited supply, the effect on marine

ecosystems can be substantial (F. Zhang et al., 2022). Our knowledge of the role of pelagic fungi in cycling of phosphorus (P) derives to a great extent from Tara Ocean metagenome and metatranscriptome analyses that show the presence and expression of fungal genes encoding proteases and peptidases involved in protein cleavage, tight coupling of protein and carbohydrate degradation, and the likely preference of fungi for a particle-associated lifestyle (Baltar et al., 2021; Breyer et al., 2022). Breyer et al. (2022) found that phosphorus availability was one important controlling factor of fungal peptidase gene expression. Overall, most non-marine fungi regulate the uptake and mobilization of Pi using the phosphate-responsive signaling pathway (PHO) which is activated upon Pi-deprivation (Vila et al., 2022 and references therein). Pelagic and coastal waters, however, are not Pi-limited (Karl & Björkman, 2015). This implies that the PHO pathway in pelagic fungi (if present) might not be active considering that its activation depends on Pi-limitation. Indeed, Breyer et al. (2022) suggested that pelagic fungi could potentially cover their needs for Pi and participate in P mineralization by recycling proteins using proteases and peptidases. Expression levels of fungal proteases were found to be regulated by P, N, and Fe availability while the abundance of fungal peptidases in detected metagenomes was significantly correlated with temperature, O<sub>2</sub>, Fe and net primary production (Breyer et al., 2022). To determine whether pelagic fungi can only participate in P mineralization via protein recycling or whether additional mechanisms are involved will require laboratory experiments using additional fungal isolates from pelagic settings.

### 3.3.3. Iron and Manganese

Micronutrients like iron (Fe<sup>3+</sup>) are important for the growth of microeukaryotes including fungi because they are co-factors in enzymes and because they are critical for production of energy-rich molecules such as NADPH and ATP. Because of its low solubility in oxic waters, Fe<sup>3+</sup> is found at concentrations <1 nM in the marine water column (Street & Paytan, 2005). Little is known about the role of pelagic fungi in the cycling of micronutrients like iron (Fe), and the mechanism(s) for how they overcome low Fe<sup>3+</sup> solubility require further investigation. Metagenome and metatranscriptome surveys from 68 Tara Ocean stations showed that the ability of fungi in the 3–2,000 μm size range to express CAZymes was correlated with Fe availability (Baltar et al., 2021). This suggests that, at least for pelagic fungi with saprophytic feeding modes (on particles), pelagic fungi require mechanisms for Fe<sup>3+</sup> acquisition. Laboratory studies showed certain ascomycetous isolates from the coast of West India produce siderophores that facilitate solubilization and sequestration of Fe<sup>3+</sup> from the surrounding environment (Baakza et al., 2004; Vala et al., 2006). Additionally, under aerobic conditions, many strains of marine fungi were reported to produce hydroxamate-type siderophores during iron limitation, suggesting they can compete with bacteria for Fe in the environment (Holinsworth & Martin, 2009). The types and structure of most fungal siderophores, as well as the mechanisms involved in transportation of siderophore-bound Fe<sup>3+</sup> into the fungal cell still need to be elucidated (Holinsworth & Martin, 2009).

Manganese is a micronutrient required for the water-splitting complex of photosystem II in photosynthetic organisms, and for many other biological activities (van Hulst et al., 2017 and references therein). Marine fungi are suggested to mineralize dissolved Mn(II) that exists in nM concentrations in the open ocean (e.g., Tebo et al., 2005). Marine fungi enhance the oxidization rates of Mn(II) to Mn oxides in the water column when dissolved Mn from the ocean's surface or from phytoplankton decay is abundant (Sunda & Huntsman, 1994; Sutherland et al., 2018; Tebo et al., 2005). Likewise, experimental studies showed that filamentous Ascomycota can oxidize Mn(II) at their hyphal tips using extracellular superoxide produced during cell differentiation (Hansel et al., 2012; Tang et al., 2013). Aside from hyphal-associated Mn(II) oxidation, analyses of fungal secretomes (the set of biomolecules produced by an organism and secreted into the extracellular environment) documented the capacity of various filamentous Ascomycota to oxidize Mn(II). This capacity is dictated by species-specific Cu-dependent (e.g., tyrosinase) and FAD-dependent (e.g., glucosemethanol-choline oxidoreductases) enzymes (Zeiner et al., 2021). Further, the capacity of these Ascomycota to oxidize Mn(II) varied with secretome age (decreased enzymatic Mn(II) oxidative capacity after 21 days) (Zeiner et al., 2021). These findings are intriguing and require a thorough examination of the role of pelagic fungi in Mn(II) cycling in the water column. Fungal biomineralization of Mn(II) may also support the biogeochemical cycling of other essential micronutrients in the water column (e.g., selenium; Rosenfeld et al., 2020). Finally, this capability may have technological applications where generation of sustainable electrochemical materials produced from fungal Mn biomineralization may have advantages (Q. Li et al., 2016).

## 4. Methods for Studying Planktonic Marine Fungi

### 4.1. Cultivation-Based Methods

Recent advancements in sequencing techniques have improved our comprehension of the overall diversity of planktonic marine fungi, and isolation of fungi provides an avenue for investigating their ecophysiology, yielding key insights into their functional roles and adaptations within the marine environment (Pang et al., 2020; Velez et al., 2015). The isolation and preservation of marine fungi in global culture collections not only helps to maintain the representative biodiversity of an ecosystem at a particular moment but also carries significant potential for long-term research, particularly in the context of climate change's effects on organisms. It is conceivable that in the coming decades, we will be able to unveil profound genetic and ecophysiological adaptations through the study of isolates preserved in culture collections (V. Kumar et al., 2021). The labor-intensive and time-consuming process of isolating marine fungi, involving collection, isolation, identification, and preservation, must persist. We seize here the opportunity to recognize Jan Kohlmeyer and Brigitte Volkmann-Kohlmeyer's substantial contributions as leading collectors and identifiers of marine fungi in the twentieth and early twenty-first centuries, with an herbarium housing over 25,000 specimens (Cunliffe, 2023; Pang et al., 2023).

Among cultivated marine fungi, members of Ascomycota clearly dominate, representing 77% of cultured marine fungal species, followed by Basidiomycota (~11%), Microsporidia (~7%), and Chytridiomycota (~2%), the remaining species represented by other basal-fungal lineages. However, it's worth noting that this documented count falls far short of the estimated 10,000 marine fungal species (Jones, 2011), a discrepancy underscoring the vast fungal diversity residing in the oceans, yet to be comprehensively isolated and cataloged. This section provides a brief overview of culture methods for planktonic marine fungi, and the approaches well-suited for studying their physiology.

For sampling of seawater, small sterile bottles, tubes, bags or buckets (H. Zhao et al., 2023) are common containers that have been used for near shore locations. Van Dorn bottles can also be used for sampling after proper washing before use (Kimura et al., 1999). Niskin bottles fitted in a rosette arrangement with a conductivity, temperature and depth device are suitable at offshore sites and samples can be taken at different depths from the top 1 m (Duan et al., 2018) to hundreds of meters (Z. Gao et al., 2010; Peng & Valentine, 2021) and even exceeding 1,000 m (Breyer et al., 2023). To collect particle-associated communities, particles can be separated using devices such as the Marine Snow Catcher (Riley et al., 2012) or by using size-fractionated filtration (Peng & Valentine, 2021). Neuston samples (air-sea interface samples) can be collected by a mesh screen sampler (Cunliffe et al., 2011). It is essential to ensure the sterility of sampling equipment and consumables throughout the collection process, with constant vigilance for potential contamination risks. Managing these risks necessitates conducting checks at different stages of the process, such as deploying sealed bottles/tubes/bags/buckets containing sterile seawater on-site to detect any potential contamination. Aerosol samples at sea can be taken at the time of seawater sampling, using common air samplers such as quartz fiber or glass fiber filter-based devices for culture-independent analysis (Fröhlich-Nowoisky et al., 2012), and six-stage Andersen impactor (Yu et al., 2013) or Burkard sampler (Mescioglou et al., 2021) for culture-based examination. This allows for the assessment of culturable fungal diversity in the sampled air, providing a basis for comparison with the fungal diversity in the seawater at the same time of sampling.

Samples should be handled immediately either on board or transported to the laboratory at a low temperature (4°C) for culture as soon as possible. Seawater samples can be serially diluted (Pham et al., 2021) if there is a high sediment load in coastal waters, or membrane filtered (G. Wang et al., 2012) and the residue resuspended in a smaller volume of sterile seawater for oceanic water samples with the relatively lower number of fungal propagules (Vrijmoed, 2000). The diluent/suspension can be spread plated onto agar plates or inoculated into liquid media in flasks or microplates. Ideally, sinking particles can be separated from the seawater using particle interceptor traps (Fontanez et al., 2015) or filtered (Bochdansky et al., 2017) in order to compare fungal isolates that are in suspension to those attached to particles. Some common media suitable for the isolation of planktonic marine fungi include Sabouraud dextrose agar (Odds, 1991), malt extract agar (MEA) (Bridson & Brecker, 1970), potato dextrose agar (Beever & Bollard, 1970), Czapek Dox agar (Waksman, 1927), cornmeal agar (Hazen & Reed, 1960), glucose-yeast extract agar (Evans & Niven, 1951) and a medium containing glucose, yeast extract, peptone and starch (Rao, 1977) to list a few (all supplemented with sea salts matching the salinity of the sampling site). Original culture media can also be designed in order to mimic the in situ conditions as much as possible to

select for marine fungi (e.g., Álvarez-Barragán et al., 2023; Panno et al., 2013). Some studies used 1/5 strength of the media so as to imitate environmental concentrations (Pang et al., 2020). Antibiotics such as chloramphenicol, streptomycin sulfate and penicillin may be added to the media to inhibit bacterial growth. However, some studies may opt not to use antibiotics in order to promote fungi-bacteria interactions, ultimately enhancing culturability, as already demonstrated on deep-sea samples (Rédou et al., 2015). The inoculated media can be incubated at different temperatures and/or light regimes, always mimicking the natural conditions as much as possible. The plates should be checked daily and colonies of different types (e.g., color, pigment on agar, mycelial density, branching) be subcultured onto fresh agar plates (same media) as pure cultures for further identification based on morphology or molecular analysis. For liquid media inoculated with seawater samples, aliquots can be spread plated onto agar media to isolate colonies of different morphologies for identification as mentioned above. Rose Bengal may also be added to the media to slow down fast-growing species for the isolation of slow-growing species (Ottow, 1972). Few Mucoromycota and related taxa have been reported from the marine environment (Calabon et al., 2023), and might be sensitive to salinity (Johnson & Sparrow, 1961).

The dilution-to-extinction technique can also be used to culture slow-growing marine fungi (Overy et al., 2019). Seawater samples, either undiluted or filtered and resuspended in sterile seawater, are inoculated into wells of a microplate. A series of dilutions (e.g., 2-fold) is made with a liquid medium (mentioned above), preferably with a reduced strength, along the rows/columns of the microplates. Dilutions with the highest frequency of single colonies are plated out (Collado et al., 2007).

Direct plating (spread plating, streaking) of samples and baiting of pine pollen, snake skin, other keratinous substrates, or live hosts are common methods used to isolate saprobic zoosporic true fungi (e.g., Van den Wyngaert et al., 2022). Seawater samples can be directly plated/streaked onto Emerson's 1/4 YpSs agar medium (S. F. Chen & Chien, 1998). For the baiting method, seawater samples are seeded with sterilized pine pollen, and/or snake skin fragments supplemented with antibiotics (Guo et al., 2023). Infected pollens or snake skin fragments are picked by a loop and streaked onto Emerson's 1/4 YpSs agar medium or the seawater with zoospores is spread plated onto the medium. Colonies with large sporangia on the agar medium after incubation (both plating or baiting methods) are picked up under a stereomicroscope, subcultured by streaking, purified and maintained on the same medium. The colony morphology of zoosporic true fungi and Labyrinthulomycetes is very similar rendering isolation of the former group very difficult.

Zoosporic fungi that are parasitic require co-culturing with the hosts. Seawater samples can be directly (or serially diluted) inoculated into the culture medium (e.g., Guillard's medium (f/2-Si), F/2 medium, Jaworski's medium) in multi-well plates with the enrichment of monocultures of diatoms (Scholz et al., 2017) or dinoflagellates (Fernández-Valero et al., 2022). Cultures are obtained after a few subcultures into axenic hosts (Fernández-Valero et al., 2022).

Recent mid/high-throughput culturing approaches (M/HTC) may also be implemented to enhance culturability and the number of marine fungal isolates. M/HTC methods enhance the recovery and isolation of the broadest possible representation of in situ fungi based on the fact that a large number of parameter combinations can be tested at once (temperatures, salinities, pH, oxygen concentrations, different substrates and different concentrations of each). Possible combinations can be constrained based on knowledge of the environment (e.g., concentrations and co-distribution of nutrients and substrates such as nitrate, ammonium, phosphate) to select a reasonable number of conditions to test for isolation of environmentally relevant fungi. Laser nephelometry (BMC Labtech) has already been used as an M/HTC approach. Laser nephelometry measures light scattered by particles (unicellular and/or filamentous cells) in 96 to 384-wells and was recently used to generate >150 isolates from lower oceanic crust samples (Quemener et al., 2021). Mini/micro-satellite primed-PCR amplification can be used as a strategy to select unique fungal isolates from such a large collection generated by the M/HTC approach (Rédou et al., 2015).

Long term preservation of fungi without genetic change allows further studies in basic and applied research (Nakagiri, 2012). Planktonic marine fungi can be preserved through culture transfer, drying and freezing (Nakagiri & Jones, 2000). Freeze- or liquid-drying are suitable methods for planktonic marine fungi which produce abundant spores. Freeze-drying (in the presence of a lyoprotectant) was found feasible for long term preservation of filamentous fungi (Tan et al., 2007) and yeasts (Bond, 2007). Filamentous fungi and yeasts can be cryopreserved for years to decades at  $-80^{\circ}\text{C}$  while zoosporic true fungi can be kept in liquid nitrogen ( $-196^{\circ}\text{C}$ ). Examples of cryoprotectants prepared in seawater include 10% glycerol, 10% dimethylsulfoxide (DMSO), and

10% glycerol + 5% trehalose (Nakagiri & Jones, 2000). Cryopreservation in 10% glycerol or DMSO as cryoprotectants was found to be suitable for chytrids (Gleason et al., 2007). For parasitic species, diatoms/dinoflagellates infected with chytrids can be submerged in a mixture of 10% glycerol and 5% trehalose (Nakagiri, 2012). As part of this process, when new species are obtained and described, isolates are expected to be deposited in at least two internationally recognized culture collections (Seifert & Rossman, 2010).

#### 4.2. Methods to Study the Physiology of Planktonic Marine Fungi

Physiology of planktonic marine fungi refers to the responses (metabolism, growth, reproduction, spore germination, death) under their immediate biological (symbiosis), chemical (salinity, pH, nutrition, interference competition, pollution) and physical (temperature, ultraviolet irradiation) surroundings (adapted from Walker & White, 2017). Fungal growth has been the main attribute to assess in response to different environmental factors.

Plate assay based on colony diameter or liquid assay based on biomass or absorbance are common methods for assessing growth of filamentous fungi and yeasts. For a plate assay, actively growing mycelia (on an agar plug/block) are inoculated onto the surface of the assay agar (i.e., with supplementation of different organic or inorganic nutrients, salinities, pHs or pollutants and/or incubated under different physical conditions such as temperature). Colony diameter (average of two perpendicular diameters) represents growth of the fungi. For some enzyme assays such as cellulose degradation (e.g., carboxy-methyl cellulose), dyes are used to stain the agar after incubation (Masigol et al., 2021; Pointing et al., 1998). For detection of chitinase activity, a basal medium composed of colloidal chitin and the dye Bromocresol Purple containing crab shell flakes can be used (Masigol et al., 2021), and the appearance of purple to black area around the mycelia represents a positive reaction. However, some fungi produce abundant aerial mycelia and so this method does not reflect the true intensity of growth. This method is also not suitable for fungi with yeast-like growth.

The growth response of three planktonic fungi (*Scheffersomyces spartinae*, *Rhodotorula sphaerocarpa*, *Sarocladium kiliense*) under different temperatures and salinities was investigated using a liquid culture method (Breyer et al., 2023). Spores, yeasts (in suspension) and fragmented mycelia in suspension or as an agar plug/block are inoculated into a liquid assay medium in flasks. For yeast species, optical density of subsamples is measured periodically at 660 nm, which reflects growth (Heitger & Baltar, 2023; Salazar Alekseyeva et al., 2022). Filamentous fungi do not produce homogenous growth in liquid media. Therefore, total mycelial biomass after incubation is collected by filtration, dried and weighed for measuring growth.

Different environmental factors exert an interactive effect on fungal growth (Pang et al., 2020). It is often impractical to examine the effect of all possible combinations of physicochemical conditions (with replication) on fungal growth. Pang et al. (2020) studied the physiological growth of filamentous fungi isolated from substrates collected at a marine shallow-water hydrothermal vent site in Taiwan under the combined effect of temperature, salinity and pH using microtitre plates. Spores/fragmented mycelia in suspension are inoculated into wells of a 96-well microtitre plate prefilled with the liquid assay medium. Due to the much smaller volume of medium required (200  $\mu$ L) in each well, the combined effects of multiple factors with replications on fungal growth can be assessed by optical density at 630 nm (Langvad, 1999). This method is also applicable to zoosporic true fungi (Guo et al., 2023).

Sporulation can be used to study reproduction success of planktonic marine fungi under environmental pressure, especially those attached to a substrate. In this approach, fungi (mainly hyphomycetes) are either plug- or point-inoculated onto an assay agar medium (e.g., MEA; Damare et al., 2008). After incubation, agar plugs are retrieved from the colony, and spores are dislodged from each of the plugs by shaking with a solution made of 0.02% Tween 80 and glass beads in a sterile tube and counted by a hemocytometer (Byrne & Gareth Jones, 1975). The concentrations of spores from each agar plug are expressed as number of spores per cm square of colony. For saprobic zoosporic true fungi, isolates are spread-plated or streaked onto an Emerson's 1/4 YpSs agar medium. After incubation, seawater is flooded onto the agar plate, which is then incubated for up to an hour (Guo et al., 2023). Samples are taken from the overlaying seawater and stained/fixed with lactophenol cotton blue. The number of zoospores is counted, and the result is expressed as number of zoospores per number of colonies produced on the agar plate. This method can apply to parasitic species as number of zoospores normalized to the number of diatom or dinoflagellate cells.

Spore germination is the key life cycle stage determining successful colonization of substrates in the pelagic zone of the ocean. A spore suspension is prepared by flooding the top of the colony of the fungi (mostly hyphomycetes) with a solution of 0.02% Tween 80 and shaking to dislodge the spores. This spore suspension can be inoculated into the assay medium in wells of a microtiter plate. Samples are taken from the wells after incubation and the percentage of germinated spores are counted (Damare et al., 2008). Fungal spores/conidia are considered germinated when the length of the germ tube equaled or exceeded the largest dimension of the original spores/conidia (Nguyen Van Long et al., 2017). Following this, germination kinetics, expressed as the % of germination as a function of time, can then be determined.


The BIOLOG platform not only provides rapid identification and characterization of filamentous fungi and yeasts, it can also be applied to substrate utilization and metabolic profiling of planktonic marine fungi under the influence of physical and chemical variables (Breyer et al., 2023; Chou et al., 2022). A suspension with spores or mycelial fragments is inoculated into the 96 wells of the microplate, each with a different substrate (e.g., BIOLOG FF MicroPlate™). Optical density is measured daily at 490 nm for substrate utilization (reduction of iodinitrotetrazolium by NADH from colorless to purple) and 750 nm for mycelial growth (turbidity). On a similar basis, Mid/High-Throughput Devices like Laser Nephelometry or oCelloScope (Harirchi et al., 2023) can also be employed to evaluate fungal growth parameters such as lag time and maximal growth rate using for example, 96-wells microplates. The combination of Laser Nephelometry and OcelloScope provides robust data comprising both quantitative information (number of fungal particles in each well) and qualitative data (microscopic images of each well), thus allowing in-depth analysis of the growth potential of numerous fungal isolates simultaneously.

### 4.3. Microscopy

During the twentieth century, microscopy observations of marine fungi were limited, and when conducted, protocols were based on phase-contrast optics or non-specific stains (Kohlmeyer & Kohlmeyer, 1979), which presented a challenge when attempting to quantify natural populations. Planktonic yeast cells are typically round shaped with a diameter as small as  $\sim 1 \mu\text{m}$ , so it is difficult to distinguish them from many bacterial cells (Figure 3a). Filamentous fungi have distinct morphology compared to most other planktonic microbes (Figure 3b), but they are typically expected to be attached to particles, which makes it challenging to distinguish fungal filaments (Figure 3c). Later in the twentieth century, one of the most widely used methods in mycology and in the medical community was based on the affinity of wheat germ agglutinin (WGA) for chitin. WGA is a lectin with a high affinity for N-acetylglucosamine residues, which, in turn, constitute the monomers of the polymer chitin. This approach has then been utilized independently to detect fungi in both light and electron microscopy (Meyberg, 1988 and references therein). The addition of fluorescein isothiocyanate (FITC) as a conjugate to the WGA (WGA-FITC), enabled the enhancement of fungi detection and quantification by utilizing the fluorescence of the conjugate (Meyberg, 1988). Originally developed for plant-fungal interactions, this method was quickly integrated into the study of aquatic fungi (Montgomery et al., 1990).


Despite the high specificity of the previous method, the most popular procedure, however, has been another staining method, used in the clinical field to detect fungal infections (Harrington & Hageage, 2003). This method involves the staining of chitin using Calcofluor White (CFW) (Damare & Raghukumar, 2008; Rasconi et al., 2009). CFW has a high affinity for the  $\beta 1-3$  and  $\beta 1-4$  polysaccharides present in chitin, thus effectively staining the fungal wall. Additionally, CFW can be excited under UV light, which is one of the commonly used wavelengths in aquatic microbiology microscopy, resulting in a deep, bright blue fluorescence (Figure 4). While some adjustments may be necessary depending on the samples, such as optimizing staining intensity and adjusting pH to minimize background noise, CFW has become one of the most used methods in aquatic mycology. Its popularity might be then due to its cost-effectiveness and relatively straightforward application (e.g., Frenken et al., 2016; Gutiérrez et al., 2011; Rasconi et al., 2012). However, CFW is not entirely free of some undesired side effects. One of the main limitations is its lack of specificity for chitin, as it also stains other common organic compounds present in the aquatic environment, such as cellulose, which is widespread in both planktonic and benthic realms. As a result, while it successfully stains fungi, it may also stain other organisms like dinoflagellates, diatoms (these also contain chitin; Durkin et al., 2009), ecdysozoans, or plant and algal debris, among others. This necessitates careful examination to confirm that the stained bodies are indeed fungi, whether they are yeasts or hyphae, by observing fungi-specific characteristics. Another drawback of using CFW is that its excitation and emission spectra overlap with those of a common microbiological stain, DAPI, which specifically binds to DNA and is frequently used for characterizing and counting bacteria, protists, and filamentous fungi.





**Figure 3.** Micrographs of *Rhodotorula sphaeroacarpa* ETNP2018 (a), *Exophiala* sp. ETNP 2018 (b), and particles from seawater collected from the coast of South Carolina (c). Images were generated by FlowCam 8000. Numbers below each image represent area-based diameters in  $\mu\text{m}$  (Kydd et al., 2018).

To address these undesired consequences, alternative approaches have been developed. Researchers have sought new staining methods or modifications to improve specificity and reduce overlap with other stains, thus enhancing the accuracy of fungal identification and quantification in aquatic environments. One protocol involves the use of fluorescence in situ hybridization (FISH) (Baschien et al., 2008) or catalyzed reporter deposition fluorescence in situ hybridization (CARD-FISH) (Jobard et al., 2010). These methods utilize labeled probes that specifically bind to certain DNA regions of the target community. The probes are conjugated with fluorescent dyes that can accommodate different wavelengths, allowing for the avoidance of overlap with DAPI or any other stain. While many FISH probes are non-specific and contain mismatches, recent efforts have been made to design FISH probes at a phylum and OTU-level (Priest et al., 2021). Other challenges using FISH/CARD-FISH include the effective permeabilization of the fungal and cell walls, which is necessary to enable the probes to reach their targets. In aquatic samples, the presence of TEP in the filters can obstruct the path of the probes, or even interfere with the binding of common dyes like DAPI (Bochdanský et al., 2017). Consequently, appropriate pre-treatments are required to achieve the desired results and ensure the accurate staining of the fungal community using these



**Figure 4.** Epifluorescence microscopy photographs of planktonic fungi (1,000 $\times$ ; Olympus IX-83; UV Wide filter cube) stained with calcofluor white.

methods. Despite these challenges, FISH and CARD-FISH represent valuable alternatives for enhancing the specificity and reliability of detecting fungi in the ocean.

Chitin-binding domains (CBD) in bacterial chitinases have been used in conjugation with fluorescein isothiocyanate (CBD-FITC) as an improved staining technique for aquatic fungi (Wurzbacher & Grossart, 2012). This method offers several advantages over previous approaches. CBD-FITC is highly specific for chitin, making it superior to Calcofluor White (CFW) or WGA-FITC in terms of specificity. Unlike CFW, CBD-FITC does not stain other compounds containing sialic acid or N-acetylglucosamine residues, ensuring accurate detection of chitin-containing structures in fungi. CBD-FITC also eliminates the need for pre-treatments to create pores in cell walls, as required in FISH and CARD-FISH. This simplifies the staining process and reduces the potential for sample alteration during preparation. Another characteristic of CBD-FITC (shared with WGA-FITC) is its compatibility with other stains like DAPI and FISH. Researchers can combine the chitin-specific CBD-FITC staining with other staining techniques, allowing for a comprehensive understanding of the aquatic fungal community, and facilitating better quantification and characterization of the organisms present. Unfortunately, CBD-FITC is no longer available commercially (Klawonn, Dunker, et al., 2023), but laboratories equipped for protein purification could perform FITC labeling in-house with commercially available conjugation kits.

The approach of using commercial conjugates can be further expanded to include various fluorescent probes with different emission wavelengths. By utilizing a wide range of fluorescent probes, researchers can combine multiple staining procedures to avoid overlapping emission wavelengths and achieve a more comprehensive analysis of the aquatic fungal community. However, it is essential to acknowledge that incorporating multiple methods does increase the complexity of the staining procedures. The order in which staining or probing is performed becomes critical to avoid interference and to obtain accurate results. Careful optimization and testing are required to ensure compatibility and successful accommodation of various staining techniques, such as WGA-FITC, CARD-FISH, and DAPI (Biancalana et al., 2017). Marine fungi, like their terrestrial counterparts, exhibit autofluorescence under various wavelengths, a trait influenced by growth stage, species, and substrate availability; hence researchers need to be cautious when interpreting fluorescence-dependent fungal images (Breyer et al., 2021).

#### 4.4. Quantification of Marine Fungal Biomass and Abundance

The quantification of biomass is a fundamental prerequisite to quantify microbial contributions to marine biogeochemical cycles. While marine fungi are commonly encountered in DNA-based studies spanning global oceans (Amend et al., 2019; Breyer & Baltar, 2023; Morales et al., 2019) there remains a significant lack of comprehensive information concerning their abundances and biomass. Nonetheless, this enigmatic kingdom can exhibit biomass levels that surpass prokaryotes during phytoplankton blooms (Gutiérrez et al., 2011; Hassett et al., 2019) and associated with bathypelagic marine snow (Bochdansky et al., 2017).

Common techniques to quantify the contribution of fungi to marine microbial biomass include the use of fluorescence microscopy (FISH; Calcofluor or Wheatgerm Agglutinin staining) or the use of biomarkers (e.g., fatty


acids) as indicator for fungal biomass. Traditionally, quantitative polymerase chain reaction (qPCR) has also been used to estimate fungal abundances in environmental samples based on gene copy numbers (Taylor & Cunliffe, 2016; X. Wang et al., 2014; Y. Wang et al., 2018, 2019) but the results are not as accurate when compared to other methods (Smith & Osborn, 2009). Instead, fluorescent probes, which specifically bind to group-specific fungal rRNA genes, have been used to enumerate fungal cells and quantify biomass in the North Sea (Priest et al., 2021), which has been discussed in Section 4.3. Fluorescent staining of marine fungi can be combined with other instruments such as a flow cytometer to achieve high-throughput enumeration of natural freshwater populations (e.g., Klawonn, Dunker, et al., 2023). Another approach involves measuring the concentration of ergosterol as a proxy for fungal biomass (Gessner, 2020), with a focus on Ascomycota and Basidiomycota (Dikarya). The ergosterol extraction method was recently adapted to marine fungi to allow for the quantification of fungal biomass in the oligotrophic (low productive) regions (Salazar Alekseyeva et al., 2022). However, the ergosterol-based method falls short when dealing with Chytridiomycota taxa (Gutiérrez et al., 2020), which is important to consider during phytoplankton blooms.

Given the limitations associated with different methods, future studies aiming to comprehensively assess planktonic marine fungi would benefit from employing a combination of the aforementioned methods. Such a holistic approach has been performed in a recent study estimating the biomass of fungi in the open ocean water column across a wide range of latitudes and productivity regimes (Breyer et al., submitted). This study revealed an overall good agreement among these different techniques to estimate fungal biomass, while also indicating fungi as relevant contributors to open ocean microbial biomass. More comparative studies in similar and contrasting environments will be crucial in unraveling the mysteries of mycoplankton and their significance within the intricate complexity of marine ecosystems.

#### 4.5. Metabarcoding

Molecular techniques such as metabarcoding offer an alternative approach for comprehensively evaluating marine fungal communities. The present era of “omics” has introduced the capability to produce large-scale evaluations of marine fungal diversity by focusing on taxonomically informative regions of the rRNA gene. A comprehensive literature review was conducted to assess various studies regarding their objectives and metabarcoding methodologies employed for characterizing planktonic marine fungal communities, as illustrated in Figure 5. Our analysis identified that, out of the 18 studies reviewed, 16 employed a single genetic marker (18S, ITS [ITS1 and/or ITS2], or 28S rRNA genes), with 12 focusing on the ITS1 and/or ITS2 genetic marker, 3 on the 18S, and 1 on the 28S. Additionally, some studies adopted dual genetic markers, namely, the 18S and 28S (Hassett et al., 2017) or the 18S and ITS (Sen et al., 2021). Based on this literature search, the primers utilized for the amplification of genetic markers appear highly conserved within studies. Primers ITS1F/ITS2 and ITS3/ITS4 are often employed for the amplification of ITS1 and ITS2, respectively. In the case of the 18S region, primers FF390/FR1, also referred to as nu-SSU-1333/nu-SSU-1647, are used to target the V7/V8 region, while the primers Euk1391f/EukBr are employed to amplify the highly variable V9 region of small-subunit ribosomal RNA genes.

Primer pairs are acknowledged for their selective amplification within a target community (Stoeck et al., 2006). For instance, the primers ITS1F/ITS2 exhibit a bias toward specific fungal groups, notably Mucoromycota, Chytridiomycota (Orsi et al., 2022) and also completely miss Microsporidia (Tedersoo & Lindahl, 2016) as part of the Opisthosporidia phylum (based on the phylogeny proposed by Naranjo-Ortiz & Gabaldón, 2019). Furthermore, both ITS1F/ITS2 and ITS3/ITS4 primers have been demonstrated to be less efficient compared to alternative primers, both in terms of amplification efficiency and the identification of OTUs from diverse phyla (Beeck et al., 2014). Despite the possibility of targeting the entire ITS region in metabarcoding studies using ITS1F/ITS4 primers (Duan et al., 2018, 2021), bias may still arise due to discrimination against longer PCR products (Ihrmark et al., 2012). 18S rRNA primers designed for fungal specificity also differ highly in their fungal coverage rate of higher and basal fungal lineages, as well as in their tendency for co-amplifying non-fungal eukaryotic sequences. For instance, the commonly utilized primer pair FF390/FR1 (nu-SSU-1333/nu-SSU-1647) has recently been demonstrated to co-amplify non-fungal eukaryotic groups such as Stramenopiles, Alveolata, Rhizaria, and Telonema (Banos et al., 2020; Prévost-Bouré et al., 2011). These non-fungal eukaryotes are both abundant and diverse in marine environments, raising concerns regarding the suitability of these primers for aquatic samples although blocking primers were designed to reduce co-amplification (Banos et al., 2020).



**Figure 5.** Literature review of 18 studies analyzing planktonic fungal communities using a metabarcoding approach within the period spanning from 2017 to 2023. The reference and sampling site section highlights details regarding the reference source and the locations of seawater sampling. The study design section specifies whether samples were collected at a single time point or at a series of time points (orange), including information about the duration of the time series and the number of samples analyzed (enclosed in brackets), the sequencing methodology employed (yellow), the targeted genetic marker (green), including the primers used, and whether the bioinformatic analysis was based on operational taxonomic units (OTUs) or amplicon sequence variants (blue). Finally, the community characterization section presents the different fungal phyla detected, based on the phylogeny proposed by Naranjo-Ortiz and Gabaldón (2019), (purple), along with the percentage representation of fungal OTUs when available. The asterisks (\*) indicate low-level detections.

Every primer pair employed carries its own set of advantages and disadvantages. However, the results obtained remain valuable, provided that the authors openly acknowledge the limitations of the approach employed and weigh the results with scientific integrity. The diversity described in these studies represents a partial picture of the total fungal diversity. The ITS region was recommended by the Fungal Barcoding Consortium as a universal barcode for fungi due to its capacity for species-level resolution across a wide spectrum of fungal taxa (Schoch et al., 2012). However, this region does have the drawback of lacking a discernible phylogenetic signal, resulting in the identification of many OTUs only at the kingdom or phylum level due to the scarcity of reference sequences or inadequately annotated sequences (Nilsson et al., 2016). As discussed in Section 2 above on the diversity of planktonic marine fungi, the current ITS databases (e.g., UNITE) include few sequences sampled from the ocean, rendering many marine (both coastal and open ocean) fungal ITS OTUs unclassified even at the phylum level (Peng & Valentine, 2021). To address this challenge, Banos et al. (2018) proposed a solution by incorporating a segment of the 18S region as a phylogenetic marker during ITS amplification. This approach enables phylogeny-based classification of fungal sequences and subsequently enhances the assignment to lower taxonomic levels, particularly for unknown or poorly annotated taxa. Alternatively, the 28S region can achieve higher taxonomic resolution than the 18S region and allow phylogenetic reconstruction (Xu et al., 2016). Rising interests in the 28S region have led to efforts such as the Fungal 28S Ribosomal RNA (LSU) RefSeq Targeted Loci Project that will likely facilitate taxonomic classifications in future studies (Robbertse, 2023). The entire rRNA region can be sequenced to provide maximum phylogenetic resolution and accuracy (Heeger et al., 2018; Runnel et al., 2022; Tedersoo et al., 2018), but this approach has not been adopted broadly due to cost and technical challenges such as obtaining high-molecular-weight DNA from environmental samples.

This comprehensive literature review underscores the prevalence of Ascomycota, Basidiomycota, and Chytridiomycota, although other fungal taxonomic phyla are also detected, albeit at lower abundances (Mucoromycota, Glomeromycota, Opisthosporidia, etc.). This trend is consistently observed across studies, whether based on OTUs or ASVs, and in both single time point and time series investigations (Figure 5). Notably, in polar marine environments, there is an intriguing departure from this pattern, with Chytridiomycota appearing to dominate, suggesting a unique ecological niche for these fungi (Burgaud et al., 2022).

Metabarcoding also provides the potential to categorize fungal OTUs/ASVs into ecological roles (e.g., animal pathogens, plant pathogens, wood saprotrophs) within three trophic modes (pathotroph, symbiotroph, and saprotroph) (Nguyen et al., 2016). Recent application of this tool to coastal water samples in the Western Pacific Ocean suggests that planktonic fungi significantly influence biogeochemical cycles and food webs through their multi-trophic nutrition (W. Li et al., 2019). Despite some limitations, such as underrepresentation of both the more evolved groups (e.g., *Aspergillus*, *Cladosporium*) and the basal fungal lineages (e.g., Chytridiomycota, Opisthosporidia), this approach facilitates the generation of hypotheses regarding the ecological roles of marine fungi. Furthermore, network analyses based on weighted topological overlaps (Banos et al., 2020) can reveal co-occurrence patterns among fungal OTUs/ASVs, allowing the inference of specific lifestyles like saprotrophy, antagonism, parasitism, etc., thus providing an additional layer of information.

A recent investigation into planktonic marine fungi in Chinese coastal waters, utilizing metabarcoding and metatranscriptomics, has revealed contrasting results from the two methods. ITS2 metabarcoding failed to detect early diverging fungal lineages that were abundant in the metatranscriptomic data set (M. Wang et al., 2023). This observation aligns with findings from a prior investigation that focused on oceanic oxygen minimum zones and employed metabarcoding and metagenomics. In this earlier study, a comparable pattern emerged, wherein the metagenomic data set revealed approximately one-third of early diverging fungal phyla that were not detected by metabarcoding (Peng & Valentine, 2021). This underscores the imperative need for employing integrated approaches, such as metabarcoding in conjunction with metagenomics and/or metatranscriptomics, to provide an accurate depiction of the community structure of planktonic marine fungi.

#### 4.6. Omics Methods

The application and integration of genomics, transcriptomics, proteomics, and metabolomics in marine microbiology in the past two decades have brought about significant advances (Kim, 2016). However, the application of these modern approaches to study marine fungi has been limited. For example, the 1,000 fungal genome project has brought the entire mycological community into the genomics age (Spatafora et al., 2017), but fewer than 10 of the sequenced genomes were of marine origin. The number of individual genomics studies on marine fungi is very small, and most published studies using genomics to study marine fungi are motivated by potential biotechnological applications of marine fungi (Ameen et al., 2021; Kempken, 2023; A. Kumar et al., 2018; Xue et al., 2022). Given that nearly 1,900 marine fungal species have been described (Calabon et al., 2023), there needs to be a community effort to increase the number of genomic studies of marine fungi.

Phylogenomic investigation of marine fungi can help elucidate evolutionary relationships between marine fungi and terrestrial fungi, as there is still a large uncertainty whether fungi originated in the sea or on land (Raghukumar, 2017b). A recent comparative genomics study of 15 red yeast *Rhodotorula* species isolated from different environments showed that the oceanic strain hosts the smallest of the 15 genomes and yet fully conserves core metabolic pathways, suggesting the adaptation of *Rhodotorula* to the oligotrophic ocean (Lane et al., 2023). A genomic characterization of *Emericellopsis atlantica*, wood-associated *Amylocarpus encephaloides* and algae-associated *Calycina marina* showed that these marine fungi have a generalist lifestyle and can degrade multiple types of marine biomass while possessing a large repertoire of CAZymes (Hagestad et al., 2021).

A challenge for studying microbial eukaryotes stems from the added complexity of eukaryotic genome assembly and annotation compared to relatively straightforward prokaryotic genomes that can typically be performed using an unsupervised, self-trained bioinformatic pipeline (Hyatt et al., 2010). Eukaryotic genome annotation usually requires supervision, and given the presence of variable number of repeated regions that require masking, introns, and exons, is best complemented with transcriptomic and proteomic evidence to improve model accuracies (Stanke & Waack, 2003; Yandell & Ence, 2012). Facing these challenges, transcriptome sequencing has been adopted as a complementary approach to investigate the functional diversity of marine microbial eukaryotes (Marine Microbial Eukaryotic Transcriptome Sequencing Project, “MMETSP”) (Keeling et al., 2014). Like the 1,000 fungal genome project, marine fungi are poorly represented by the MMETSP despite the large number of cultivated fungi from marine environments. Future studies using transcriptomics can be designed to study physiological adaptation of marine fungi to environmental changes. However, transcriptomic profiling is ideally performed with reference genomes, which are still rare for marine fungi. To avoid this “chicken-and-egg” problem, both genomics and transcriptomics should be incorporated in experimental designs.

When applied to natural assemblages of microorganisms, genomics and transcriptomic sequencing takes the “meta” form and can reveal the diversity, function, and activity of marine fungi. However, the low relative abundance of fungal DNA (as low as less than 0.02%, e.g., Peng & Valentine, 2021) and RNA (~1% in seawater) (Kolody et al., 2019) poses a major challenge. Theoretically, increasing the metagenome sequencing depth to at least 10 billion reads (up to 250 bp per read) per sample could recover millions of fungal reads, but even at today's reduced cost of high-throughput sequencing, the cost of performing such an experiment would be prohibitive. Most up-to-date studies use a sequencing depth up to hundreds of millions of reads per sample (Lan et al., 2022), making it challenging to reconstruct fungal metagenome-assembled genomes (MAGs), which have been achieved in environmental samples with relatively low diversity and high fungal abundance (Peng et al., 2021; West et al., 2018). Despite these limitations, metagenomics have been used to assess fungal diversity in the ocean (Hassett et al., 2020; Morales et al., 2019; Peng & Valentine, 2021), as it has the advantage of avoiding biases associated with PCR, a required step of metabarcoding methods. The analysis of a metagenome data set from the eastern tropical North Pacific oxygen minimum zone revealed that early diverging fungi accounted for about one third of the fungal community at locations where metabarcoding of the ITS2 region detected no more than 5% of early diverging fungi (Peng & Valentine, 2021). The accuracy of classifying metagenomic reads depends upon the quality of the reference database, so it is critical to increase the number of marine fungal genomes which will serve as references.

Although current metagenome data sets typically do not have sufficient sequencing depth to enable the recovery of fungal MAGs, many fungal contigs are included in the general metagenomes assemblies and can be separated using tools such as EukRep (West et al., 2018). Moreover, eukaryotic metatranscriptome libraries constructed by targeting the polyadenylated tails of mature transcripts selecting for mainly protein-coding messenger RNA are another important method to investigate the function and activity of marine fungi. This approach has enabled the discovery of fungal transcripts coding for cell division and hydrolases involved in lipid, carbohydrate, and protein degradation in the deep biosphere (Orsi et al., 2013; Quemener et al., 2020), as well as the ocean water columns (Orsi et al., 2022; M. Wang et al., 2023). Additionally, there is a large potential for discovery in publicly available marine metagenomes and metatranscriptomes, most of which were primarily explored targeting bacteria and archaea. For example, analyses of the TARA Oceans data sets revealed a widespread utilization of different types of enzymes that hydrolyze carbohydrates and proteins by pelagic fungi, which potentially occupy a deeper niche than marine bacteria (Baltar et al., 2021; Breyer et al., 2022; Christmas & Cunliffe, 2020).

Proteomic analysis has been used to study the enzymatic profile of marine fungi, especially the secreted proteins (secretome) (Pilgaard et al., 2019). Proteomics are typically performed to complement genomic and/or transcriptomic analysis of isolated fungal strains, and provide an additional line of evidence for fungal metabolisms such as polysaccharide depolymerization (Pilgaard et al., 2019) and secondary metabolite production (Kramer et al., 2015). In contrast, untargeted metaproteomics remains highly challenging not only because of the lack of reference genomes and transcriptomes of planktonic marine fungi, but the low concentration of proteins in the water column makes sample acquisition highly demanding, which requires filtration of at least tens to hundreds of liters of seawater (Saito et al., 2019).

Metabolomics measure a group metabolites from organisms and has been a staple tool for systems biology for decades (Kell, 2004). Application of metabolomics in fungal research, including the subfield of marine fungi, has largely focused on the measurements of secondary metabolites and usually driven by the interests in natural product discovery (G. Li et al., 2022; Stuart et al., 2020). Metabolic profiling of marine fungi captures snapshots of fungal metabolites under different growth conditions. Dissolved metabolites in the aqueous phase are the most common type of metabolites analyzed using modern instruments such as high-performance liquid chromatography-mass spectrometry (G. Li et al., 2022). Very few metabolomic studies have been performed using planktonic marine fungi as subjects. Metabolic profiling of the marine fungus *Emericellopsis cladophorae* demonstrated its ability to produce antimicrobial and anti-inflammatory compounds (Gonçalves, Hilário, et al., 2022).

#### 4.7. Stable Isotope-Based Methods

Stable isotopes have been applied broadly in biogeochemistry, microbial ecology, and oceanography to study the microbial transformation of elements from molecular to ecosystem levels. The tracing of added stable isotopes in incubations is often combined with other techniques to provide novel insights. In this section we highlight three

stable isotope-based methods: secondary ion mass spectrometry (SIMS), stable isotope probing, and biogeochemical rate measurements.

The incorporation of substrates labeled with stable isotopes (e.g.,  $^{13}\text{C}$  and  $^{15}\text{N}$ ) by individual cells in mono- or co-cultures or in a natural assemblage of microbial community can be visualized using SIMS, or nanoSIMS that can focus the primary ion beam down to 50 nm, enabling the quantification of stable isotopes with extremely high spatial resolution (Mayali, 2020). This technique quantitatively determines the number of both the rare (e.g.,  $^{15}\text{N}$ ) and abundant isotopes (e.g.,  $^{14}\text{N}$ ), as well as their spatial distribution, within the cell. Very few studies have applied nanoSIMS to the study of marine or aquatic fungi, partially due to limited instrument availability and the high cost associated with the analysis. A recent study incubated  $^{13}\text{C}$ -labeled polyethylene with mono-cultures of *R. mucilaginosa* isolated from seawater and used nanoSIMS to trace the incorporation of plastic-derived carbon into individual cells in addition to polyethylene degradation rates (Vaksmas et al., 2023). They found that *R. mucilaginosa* incorporated UV-treated  $^{13}\text{C}$ -polyethylene, which suggests the red yeast could contribute to plastic degradation in marine environments. NanoSIMS has also been used to demonstrate that in a model system of freshwater diatom and chytrids, fungi derived ~100% of their carbon content from the diatom (Klawonn et al., 2021). Klawonn et al. (2021) also showed that in a complex microbial community, unidentified taxa can be identified using FISH at the phylum or class level. Given the versatility of stable isotope labeled substrates, future experiments including nanoSIMS as a method can reveal novel biochemical pathways of planktonic marine fungi, as well as the roles fungi play in marine food webs.

While nanoSIMS offers a visual approach to quantify the incorporation of stable isotope-labeled substrates by cells, DNA stable isotope probing (DNA-SIP) reveals the identity and functional potential of the microorganisms responsible for the metabolism of added substrates, which is particularly useful in a complex microbial community of which most members are uncultivated. In marine microbial communities, saprotrophic fungi and bacteria play similar roles, such as in biomass degradation. DNA-SIP allows the identification of both fungal and bacterial lineages responsible for biomass degradation. The DNA of microorganisms that incorporated labeled substrate (e.g.,  $^{13}\text{C}$ -labeled polysaccharide) can be separated from the DNA without stable isotope labels by ultracentrifugation (Neufeld et al., 2007). Separated  $^{13}\text{C}$ -labeled DNA can be used for metabarcoding and metagenomic analysis as discussed above, but they represent the microbial community that incorporated the added substrate. Seawater incubations with  $^{13}\text{C}$ -labeled diatom biomass have demonstrated that *Malassezia* and *Cladosporium* assimilate microalgae-derived biomass in the ocean water column (Cunliffe et al., 2017; Orsi et al., 2022). The design of DNA-SIP experiments should consider ways to minimize potential cross feeding, which refers to the incorporation of stable isotope labels by cells that did not directly metabolize stable isotope-labeled substrate (e.g., by feeding on other cells that directly metabolize the labeled substrate).

Incubation with stable isotopes is a common method used to determine the transformation rates of chemical elements in any environment on earth, including in the ocean (Glibert et al., 2019). When stable isotope incubations are applied to a natural assemblage of microorganisms, the overall rates of elemental transformation mediated by microbial enzymes and abiotic processes are measured (e.g., nitrification by both archaea and bacteria in the ocean; Peng et al., 2015). The contribution of fungi to a biogeochemical process, such as denitrification, can be constrained by applying antimicrobial compounds that inhibit specific microbial groups in stable isotope incubations (Peng & Valentine, 2021). For example, chloramphenicol, a broad-spectrum antibiotic commonly used to isolate fungi from the environment (see Section 4.1), inhibits both bacteria and archaea (Yunis, 1988). On the other hand, antifungal agents such as cycloheximide could be used to inhibit fungal activities. In theory, if the antimicrobial compound achieved 100% specific and effective inhibition of the targeted groups (e.g., bacteria and archaea), the remainder rates measured represent the contribution of the non-targeted groups (e.g., fungi and other microbial eukaryotes). Peng and Valentine (2021) combined seawater incubations with  $^{15}\text{N}$ -labeled nitrate with chloramphenicol treatments and found that fungi in the eastern tropical North Pacific Ocean can contribute up to 10% of the production of  $\text{N}_2\text{O}$ , a potent ozone-depleting greenhouse gas. However, one of the major limitations of this approach using antimicrobial compounds is that the inhibition of targeted groups may not be complete (Salkin & Hurd, 1972), sometimes due to antibiotic-resistance (Larsson & Flach, 2022), and the antimicrobial compound may affect non-targeted organisms (Castaldi & Smith, 1998; Rousk et al., 2009). Therefore, the results from incubations with antimicrobial compounds must be interpreted with caution, and ideally the optimal concentration of antimicrobial compounds that maximizes their effectiveness and specificity for each sample type (e.g., open ocean vs. coastal seawater) should be determined with pilot experiments.

## 5. Outlook

This review of the study of planktonic marine fungi shows the large potential for new discoveries of fungal diversity and functions in marine environments, especially in the open ocean. To reconcile some of the inconsistent reports of fungal diversity based on metabarcoding surveys using different primers, we recommend using primer pairs targeting the large subunit of the rRNA gene because they are less biased against early diverging fungi compared to ITS primers and they typically allow taxonomic classification at lower levels than the small subunit of the rRNA gene. Once long-read sequencing becomes more affordable and tractable, it will be another promising alternative as nearly the entire rRNA region can be sequenced. Metagenomics and metatranscriptomics are becoming increasingly useful tools for assessing the diversity, function, and activity of planktonic marine fungi.

In addition to assessing planktonic fungal diversity with metabarcoding surveys, it is essential to determine the biomass of marine fungi to elucidate the flow of energy and nutrients in marine food webs. Future cultivation efforts could learn from the development of marine bacteria studies, in which it took decades to isolate the most abundant and prevalent heterotrophic bacteria in the ocean. Specifically, enrichment and isolation media would ideally be designed to mimic in situ nutrient and oxygen concentrations, temperature, light level, and pressure. Analysis of marine fungal genomes and transcriptomic profiling will generate valuable insights into how planktonic fungi would adapt to a changing ocean and interact with other members of the marine microbiome.

## Data Availability Statement

Previously published data from the MetaPR2 project can be accessed at <https://app.metapr2.org/metapr2/> (Vaultot, 2022).

## References

- Agoué, H., Casamayor, E. O., Joux, F., Obermosterer, I., Dupuy, C., Lantoine, F., et al. (2004). Comparison of samplers for the biological characterization of the sea surface microlayer. *Limnology and Oceanography: Methods*, 2(7), 213–225. <https://doi.org/10.4319/lom.2004.2.213>
- Álvarez-Barragán, J., Cravo-Laureau, C., Xiong, B., Wick, L. Y., & Duran, R. (2023). Marine fungi select and transport aerobic and anaerobic bacterial populations from polycyclic aromatic hydrocarbon-contaminated sediments. *mBio*, 14(2), e02761-22. <https://doi.org/10.1128/mbio.02761-22>
- Ameen, F., AlNadhari, S., & Al-Homaidan, A. A. (2021). Marine microorganisms as an untapped source of bioactive compounds. *Saudi Journal of Biological Sciences*, 28(1), 224–231. <https://doi.org/10.1016/j.sjbs.2020.09.052>
- Amend, A., Burgaud, G., Cunliffe, M., Edgcomb, V. P., Ettinger, C. L., Gutiérrez, M. H., et al. (2019). Fungi in the marine environment: Open questions and unsolved problems. *mBio*, 10(2), e01189-18. <https://doi.org/10.1128/mBio.01189-18>
- Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyer-Reil, L.-A., & Thingstad, F. (1983). The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series*, 10(3), 257–263. <https://doi.org/10.3354/meps010257>
- Baakza, A., Vala, A. K., Dave, B. P., & Dube, H. C. (2004). A comparative study of siderophore production by fungi from marine and terrestrial habitats. *Journal of Experimental Marine Biology and Ecology*, 311(1), 1–9. <https://doi.org/10.1016/j.jembe.2003.12.028>
- Bacic, M. K., & Yoch, D. C. (1998). In vivo characterization of dimethylsulfoniopropionate lyase in the fungus *Fusarium lateritium*. *Applied and Environmental Microbiology*, 64(1), 106–111. <https://doi.org/10.1128/AEM.64.1.106-111.1998>
- Baltar, F., Zhao, Z., & Herndl, G. J. (2021). Potential and expression of carbohydrate utilization by marine fungi in the global ocean. *Microbiome*, 9(1), 106. <https://doi.org/10.1186/s40168-021-01063-4>
- Banos, S., Gysi, D. M., Richter-Heitmann, T., Glöckner, F. O., Boersma, M., Wiltshire, K. H., et al. (2020). Seasonal dynamics of pelagic mycoplanktonic communities: Interplay of taxon abundance, temporal occurrence, and biotic interactions. *Frontiers in Microbiology*, 11, 1305. <https://doi.org/10.3389/fmicb.2020.01305>
- Banos, S., Lentendu, G., Kopf, A., Wubet, T., Glöckner, F. O., & Reich, M. (2018). A comprehensive fungi-specific 18S rRNA gene sequence primer toolkit suited for diverse research issues and sequencing platforms. *BMC Microbiology*, 18(1), 190. <https://doi.org/10.1186/s12866-018-1331-4>
- Baschien, C., Manz, W., Neu, T. R., Marvanová, L., & Szewzyk, U. (2008). In situ detection of freshwater fungi in an alpine stream by new taxon-specific fluorescence in situ hybridization probes. *Applied and Environmental Microbiology*, 74(20), 6427–6436. <https://doi.org/10.1128/AEM.00815-08>
- Bass, D., Howe, A., Brown, N., Barton, H., Demidova, M., Michelle, H., et al. (2007). Yeast forms dominate fungal diversity in the deep oceans. *Proceedings of the Royal Society B: Biological Sciences*, 274(1629), 3069–3077. <https://doi.org/10.1098/rspb.2007.1067>
- Basu, S., & Mackey, K. R. M. (2018). Phytoplankton as key mediators of the biological carbon pump: Their responses to a changing climate. *Sustainability*, 10(3), 869. <https://doi.org/10.3390/su10030869>
- Baumas, C. M. J., Le Moigne, F. A. C., Garel, M., Bhairy, N., Guasco, S., Riou, V., et al. (2021). Mesopelagic microbial carbon production correlates with diversity across different marine particle fractions. *The ISME Journal*, 15(6), 1695–1708. <https://doi.org/10.1038/s41396-020-00880-z>
- Baztan, J., Chouinard, O., Jorgensen, B., Tett, P., Vanderlinden, J. P., & Vasseur, L. (2015). Introduction. In *Coastal zones: Solutions for the 21st century* (pp. xxi–xxiii). <https://doi.org/10.1016/B978-0-12-802748-6.02001-5>
- Beeck, M. O. D., Lievens, B., Busschaert, P., Declerck, S., Vangronsveld, J., & Colpaert, J. V. (2014). Comparison and validation of some ITS primer pairs useful for fungal metabarcoding studies. *PLoS One*, 9(6), e97629. <https://doi.org/10.1371/journal.pone.0097629>
- Beever, R. E., & Bollard, E. G. (1970). The nature of the stimulation of fungal growth by potato extract. *Microbiology*, 60(2), 273–279. <https://doi.org/10.1099/00221287-60-2-273>

## Acknowledgments

ASA and SW were supported by award GBMF9343 from the Gordon and Betty Moore Foundation. FB and EB were supported by the Austrian Science Fund (FWF) projects OCEANIDES (P34304-B), ENIGMA (TAI534), EXEBIO (P35248), and OCEANBIOPLAST (P35619-B). LBB was supported by the Simons Foundation International through the BIOSCOPE program. GB thanks the French National Research Agency ANR-19-CE04-0001-01 Mycoplast project. PM and VE were supported by the National Science Foundation Grants OCE-2046799 and OCE-1829903. EU-Biodiversa project FUNACTION by the German Science foundation (DFG) (GR1540/47-1) funded AR and HPG. DFG project Pycnocline (GR1540/37-1) funded HM and HPG. AKW thanks the Natural Sciences and Engineering Research Council of Canada for Discovery Grant support (no. NSERC—2017-04325). KLP was funded by the National Science and Technology Council, Taiwan (NSTC 111-2621-M-019-002-, NSTC 111-2621-B-019-001-MY3, NSTC 112-2621-M-019-003-). XP was supported by the National Science Foundation Award DEB-2303089. XP thanks Savannah Judge and Joshua Stone for their assistance in acquiring FlowCam images. CR was supported by an ARIES DTP PhD studentship funded from the UK Natural Environmental Research Council (NERC) (NE/S007334/1). CR and MC were supported by the European Research Council (ERC) (MYCO-CARB project Grant 772584).



- Biancalana, F., Kopprio, G. A., Lara, R. J., & Alonso, C. (2017). A protocol for the simultaneous identification of chitin-containing particles and their associated bacteria. *Systematic and Applied Microbiology*, 40(5), 314–320. <https://doi.org/10.1016/j.syapm.2017.05.004>
- Bik, H. M., Halanach, K. M., Sharma, J., & Thomas, W. K. (2012). Dramatic shifts in benthic microbial eukaryote communities following the deepwater horizon oil spill. *PLoS One*, 7(6), e38550. <https://doi.org/10.1371/journal.pone.0038550>
- Bochdandy, A. B., Clouse, M. A., & Herndl, G. J. (2017). Eukaryotic microbes, principally fungi and labyrinthulomycetes, dominate biomass on bathypelagic marine snow. *The ISME Journal*, 11(2), 362–373. <https://doi.org/10.1038/ismej.2016.113>
- Bond, C. (2007). Cryopreservation of yeast cultures. In J. G. Day & G. N. Stacey (Eds.), *Cryopreservation and freeze-drying protocols* (pp. 109–117). Humana Press. [https://doi.org/10.1007/978-1-59745-362-2\\_7](https://doi.org/10.1007/978-1-59745-362-2_7)
- Bonugli-Santos, R. C., Durrant, L. R., da Silva, M., & Sette, L. D. (2010). Production of laccase, manganese peroxidase and lignin peroxidase by Brazilian marine-derived fungi. *Enzyme and Microbial Technology*, 46(1), 32–37. <https://doi.org/10.1016/j.enzmictec.2009.07.014>
- Bradford-Grieve, J. M., Nodder, S. D., Jillett, J. B., Currie, K., & Lassey, K. R. (2001). Potential contribution that the copepod *Neocalanus tonsus* makes to downward carbon flux in the Southern Ocean. *Journal of Plankton Research*, 23(9), 963–975. <https://doi.org/10.1093/plankt/23.9.963>
- Breyer, E., & Baltar, F. (2023). The largely neglected ecological role of oceanic pelagic fungi. *Trends in Ecology & Evolution*, 38(9), 870–888. <https://doi.org/10.1016/j.tree.2023.05.002>
- Breyer, E., Böhm, M., Reitbauer, M., Amano, C., Heitger, M., & Baltar, F. (2021). Autofluorescence is a common trait in different oceanic fungi. *Journal of Fungi*, 7(9), 709. <https://doi.org/10.3390/jof7090709>
- Breyer, E., Espada-Hinojosa, S., Reitbauer, M., Karunarathna, S. C., & Baltar, F. (2023). Physiological properties of three pelagic fungi isolated from the Atlantic Ocean. *Journal of Fungi*, 9(4), 439. <https://doi.org/10.3390/jof9040439>
- Breyer, E., Zhao, Z., Herndl, G. J., & Baltar, F. (2022). Global contribution of pelagic fungi to protein degradation in the ocean. *Microbiome*, 10(1), 143. <https://doi.org/10.1186/s40168-022-01329-5>
- Bridson, E. Y., & Brecker, A. (1970). Chapter III Design and formulation of microbial culture media. In J. R. Norris & D. W. Ribbons (Eds.), *Methods in microbiology* (Vol. 3, pp. 229–295). Academic Press. [https://doi.org/10.1016/S0580-9517\(08\)70541-5](https://doi.org/10.1016/S0580-9517(08)70541-5)
- Bronk, D. A., & Glibert, P. M. (1993). Application of a <sup>15</sup>N tracer method to the study of dissolved organic nitrogen uptake during spring and summer in Chesapeake Bay. *Marine Biology*, 115(3), 501–508. <https://doi.org/10.1007/BF00349849>
- Burgaud, G., Edgcomb, V., Hassett, B. T., Kumar, A., Li, W., Mara, P., et al. (2022). Marine fungi. In L. J. Stal & M. S. Cretoiu (Eds.), *The marine microbiome* (pp. 243–295). Springer International Publishing. [https://doi.org/10.1007/978-3-030-90383-1\\_5](https://doi.org/10.1007/978-3-030-90383-1_5)
- Byrne, P. J., & Gareth Jones, E. B. (1975). Effect of salinity on the reproduction of terrestrial and marine fungi. *Transactions of the British Mycological Society*, 65(2), 185–200. [https://doi.org/10.1016/S0007-1536\(75\)80002-7](https://doi.org/10.1016/S0007-1536(75)80002-7)
- Calabon, M. S., Jones, E. B. G., Pang, K.-L., Abdel-Wahab, M. A., Jin, J., Devadatha, B., et al. (2023). Updates on the classification and numbers of marine fungi. *Botanica Marina*, 66(4), 213–238. <https://doi.org/10.1515/bot-2023-0032>
- Castaldi, S., & Smith, K. A. (1998). Effect of cycloheximide on N<sub>2</sub>O and NO<sub>3</sub><sup>-</sup> production in a forest and an agricultural soil. *Biology and Fertility of Soils*, 27(1), 27–34. <https://doi.org/10.1007/s003740050395>
- Chen, H., Mothapo, N. V., & Shi, W. (2014). The significant contribution of fungi to soil N<sub>2</sub>O production across diverse ecosystems. *Applied Soil Ecology*, 73, 70–77. <https://doi.org/10.1016/j.apsoil.2013.08.011>
- Chen, S. F., & Chien, C.-Y. (1998). Some chytrids of Taiwan (II). *Botanical Bulletin of Academia Sinica*, 39, 47–56.
- Chen, W., Lee, M.-K., Jefcoate, C., Kim, S.-C., Chen, F., & Yu, J.-H. (2014). Fungal cytochrome P450 monooxygenases: Their distribution, structure, functions, family expansion, and evolutionary origin. *Genome Biology and Evolution*, 6(7), 1620–1634. <https://doi.org/10.1093/gbe/evu132>
- Cheng, J., Jacquin, J., Conan, P., Pujo-Pay, M., Barbe, V., George, M., et al. (2021). Relative influence of plastic debris size and shape, chemical composition and phytoplankton-bacteria interactions in driving seawater plastisphere abundance, diversity and activity. *Frontiers in Microbiology*, 11, 610231. <https://doi.org/10.3389/fmicb.2020.610231>
- Chou, H.-Y., Chiang, M. W.-L., Lin, W.-R., Hsieh, S.-Y., Jones, E. B. G., Guo, S.-Y., & Pang, K.-L. (2022). Metabolic activity on bioglyph MicroPlate suggests organic substrate decomposition by *Aspergillus terreus* NT0U4989 isolated from Kueishan Island Hydrothermal Vent Field, Taiwan. *Fungal Ecology*, 60, 101157. <https://doi.org/10.1016/j.funeco.2022.101157>
- Christmas, N., Allen, R., Allen, M. J., Bird, K., & Cunliffe, M. (2023). A 17-year time-series of fungal environmental DNA from a coastal marine ecosystem reveals long-term seasonal-scale and inter-annual diversity patterns. *Proceedings of the Royal Society B: Biological Sciences*, 290(1992), 20222129. <https://doi.org/10.1098/rspb.2022.2129>
- Christmas, N., & Cunliffe, M. (2020). Depth-dependent mycoplankton glycoside hydrolase gene activity in the open ocean—Evidence from the Tara Oceans eukaryote metatranscriptomes. *The ISME Journal*, 14(9), 2361–2365. <https://doi.org/10.1038/s41396-020-0687-2>
- Clipson, N., Landy, E., & Otte, M. (2005). Biogeochemical roles of fungi in marine and estuarine habitats. In *Micro-organisms and earth systems - Advances in geomicrobiology* (pp. 321–344). Society for General Microbiology. <https://doi.org/10.1017/CBO9780511754852.016>
- Collado, J., Platas, G., Paulus, B., & Bills, G. F. (2007). High-throughput culturing of fungi from plant litter by a dilution-to-extinction technique. *FEMS Microbiology Ecology*, 60(3), 521–533. <https://doi.org/10.1111/j.1574-6941.2007.00294.x>
- Conrad, R., & Seiler, W. (1988). Influence of the surface microlayer on the flux of nonconservative trace gases (CO, H<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O) across the ocean-atmosphere interface. *Journal of Atmospheric Chemistry*, 6(1), 83–94. <https://doi.org/10.1007/BF00048333>
- Cunliffe, M. (2023). Who are the marine fungi? *Environmental Microbiology*, 25(1), 131–134. <https://doi.org/10.1111/1462-2920.16240>
- Cunliffe, M., Hollingsworth, A., Bain, C., Sharma, V., & Taylor, J. D. (2017). Algal polysaccharide utilisation by saprotrophic planktonic marine fungi. *Fungal Ecology*, 30, 135–138. <https://doi.org/10.1016/j.funeco.2017.08.009>
- Cunliffe, M., Schäfer, H., Harrison, E., Cleave, S., Upstill-Goddard, R., & Murrell, J. C. (2008). Phylogenetic and functional gene analysis of the bacterial and archaeal communities associated with the surface microlayer of an estuary. *The ISME Journal*, 2(7), 776–789. <https://doi.org/10.1038/ismej.2008.28>
- Cunliffe, M., Upstill-Goddard, R. C., & Murrell, J. C. (2011). Microbiology of aquatic surface microlayers. *FEMS Microbiology Reviews*, 35(2), 233–246. <https://doi.org/10.1111/j.1574-6976.2010.00246.x>
- Damare, S. R., Nagarajan, M., & Raghukumar, C. (2008). Spore germination of fungi belonging to *Aspergillus* species under deep-sea conditions. *Deep Sea Research Part I: Oceanographic Research Papers*, 55(5), 670–678. <https://doi.org/10.1016/j.dsr.2008.02.004>
- Damare, S. R., & Raghukumar, C. (2008). Fungi and macroaggregation in deep-sea sediments. *Microbial Ecology*, 56(1), 168–177. <https://doi.org/10.1007/s00248-007-9334-y>
- Danger, M., & Chauvet, E. (2013). Elemental composition and degree of homeostasis of fungi: Are aquatic hyphomycetes more like metazoans, bacteria or plants? *Fungal Ecology*, 6(5), 453–457. <https://doi.org/10.1016/j.funeco.2013.05.007>
- Danger, M., Gessner, M. O., & Bärlocher, F. (2016). Ecological stoichiometry of aquatic fungi: Current knowledge and perspectives. *Fungal Ecology*, 19, 100–111. <https://doi.org/10.1016/j.funeco.2015.09.004>

- Debeljak, P., & Baltar, F. (2023). Fungal diversity and community composition across ecosystems. *Journal of Fungi*, 9(5), 510. <https://doi.org/10.3390/jof9050510>
- De Tender, C., Devriese, L. I., Haegeman, A., Maes, S., Vangeyte, J., Cattrijsse, A., et al. (2017). Temporal dynamics of bacterial and fungal colonization on plastic debris in the North Sea. *Environmental Science & Technology*, 51(13), 7350–7360. <https://doi.org/10.1021/acs.est.7b00697>
- Duan, Y., Xie, N., Song, Z., Ward, C. S., Yung, C.-M., Hunt, D. E., et al. (2018). A high-resolution time series reveals distinct seasonal patterns of planktonic fungi at a temperate coastal ocean site (Beaufort, North Carolina, USA). *Applied and Environmental Microbiology*, 84(21), e00967-18. <https://doi.org/10.1128/AEM.00967-18>
- Duan, Y., Xie, N., Wang, Z., Johnson, Z. I., Hunt, D. E., & Wang, G. (2021). Patchy distributions and distinct niche partitioning of mycoplankton populations across a nearshore to open ocean gradient. *Microbiology Spectrum*, 9(3), e01470-21. <https://doi.org/10.1128/spectrum.01470-21>
- Duret, M. T., Lampitt, R. S., & Lam, P. (2019). Prokaryotic niche partitioning between suspended and sinking marine particles. *Environmental Microbiology Reports*, 11(3), 386–400. <https://doi.org/10.1111/1758-2229.12692>
- Duret, M. T., Lampitt, R. S., & Lam, P. (2020). Eukaryotic influence on the oceanic biological carbon pump in the Scotia Sea as revealed by 18S rRNA gene sequencing of suspended and sinking particles. *Limnology and Oceanography*, 65(S1), S49–S70. <https://doi.org/10.1002/lno.11319>
- Durkin, C. A., Mock, T., & Armbrust, E. V. (2009). Chitin in diatoms and its association with the cell wall. *Eukaryotic Cell*, 8(7), 1038–1050. <https://doi.org/10.1128/ec.00079-09>
- Engel, A., Bange, H. W., Cunliffe, M., Burrows, S. M., Friedrichs, G., Galgani, L., et al. (2017). The ocean's vital skin: Toward an integrated understanding of the sea surface microlayer. *Frontiers in Marine Science*, 4, 165. <https://doi.org/10.3389/fmars.2017.00165>
- Evans, J. B., & Niven, C., Jr. (1951). Nutrition of the heterofermentative lactobacilli that cause greening of cured meat products. *Journal of Bacteriology*, 62(5), 599–603. <https://doi.org/10.1128/jb.62.5.599-603.1951>
- Falkowski, P. G., Barber, R. T., & Smetacek, V. (1998). Biogeochemical controls and feedbacks on ocean primary production. *Science*, 281(5374), 200–206. <https://doi.org/10.1126/science.281.5374.200>
- Fell, J. W. (1967). Distribution of yeasts in the Indian Ocean. *Bulletin of Marine Science*, 17(2), 454–470.
- Fernández-Valero, A. D., Reñé, A., Timoneda, N., Sampedro, N., & Garcés, E. (2022). Dinoflagellate hosts determine the community structure of marine Chytridiomycota: Demonstration of their prominent interactions. *Environmental Microbiology*, 24(12), 5951–5965. <https://doi.org/10.1111/1462-2920.16182>
- Florio Furno, M., Poli, A., Ferrero, D., Tardelli, F., Manzini, C., Oliva, M., et al. (2022). The culturable mycobiota of sediments and associated microplastics: From a harbor to a marine protected area, a comparative study. *Journal of Fungi*, 8(9), 927. <https://doi.org/10.3390/jof8090927>
- Fontanez, K. M., Eppley, J. M., Samo, T. J., Karl, D. M., & DeLong, E. F. (2015). Microbial community structure and function on sinking particles in the North Pacific Subtropical Gyre. *Frontiers in Microbiology*, 6, 469. <https://doi.org/10.3389/fmicb.2015.00469>
- Franklin, M. P., McDonald, I. R., Bourne, D. G., Owens, N. J. P., Upstill-Goddard, R. C., & Murrell, J. C. (2005). Bacterial diversity in the bacterioneuston (sea surface microlayer): The bacterioneuston through the looking glass. *Environmental Microbiology*, 7(5), 723–736. <https://doi.org/10.1111/j.1462-2920.2004.00736.x>
- Frenken, T., Velthuis, M., de Senerpont Domis, L. N., Stephan, S., Aben, R., Kosten, S., et al. (2016). Warming accelerates termination of a phytoplankton spring bloom by fungal parasites. *Global Change Biology*, 22(1), 299–309. <https://doi.org/10.1111/gcb.13095>
- Fröhlich-Nowoisky, J., Burrows, S. M., Xie, Z., Engling, G., Solomon, P. A., Fraser, M. P., et al. (2012). Biogeography in the air: Fungal diversity over land and oceans. *Biogeosciences*, 9(3), 1125–1136. <https://doi.org/10.5194/bg-9-1125-2012>
- Frost, T. (1999). Environmental controls of air-water gas exchange.
- Gao, R., Liu, R., & Sun, C. (2022). A marine fungus *Alternaria alternata* FB1 efficiently degrades polyethylene. *Journal of Hazardous Materials*, 431, 128617. <https://doi.org/10.1016/j.jhazmat.2022.128617>
- Gao, Z., Johnson, Z. I., & Wang, G. (2010). Molecular characterization of the spatial diversity and novel lineages of mycoplankton in Hawaiian coastal waters. *The ISME Journal*, 4(1), 111–120. <https://doi.org/10.1038/ismej.2009.87>
- Geller-McGrath, D., Mara, P., Taylor, G. T., Suter, E., Edgcomb, V., & Pachiadaki, M. (2023). Diverse secondary metabolites are expressed in particle-associated and free-living microorganisms of the permanently anoxic Cariaco Basin. *Nature Communications*, 14(1), 656. <https://doi.org/10.1038/s41467-023-36026-w>
- Gessner, M. O. (2020). Ergosterol as a measure of fungal biomass. In F. Bärlocher, M. O. Gessner, & M. A. S. Graça (Eds.), *Methods to study litter decomposition: A practical guide* (pp. 247–255). Springer International Publishing. [https://doi.org/10.1007/978-3-030-30515-4\\_27](https://doi.org/10.1007/978-3-030-30515-4_27)
- Gladfelter, A. S., James, T. Y., & Amend, A. S. (2019). Marine fungi. *Current Biology*, 29(6), R191–R195. <https://doi.org/10.1016/j.cub.2019.02.009>
- Gleason, F. H., Küpper, F. C., Amon, J. P., Picard, K., Gachon, C. M. M., Marano, A. V., et al. (2011). Zoospore true fungi in marine ecosystems: A review. *Marine and Freshwater Research*, 62(4), 383–393. <https://doi.org/10.1071/MF10294>
- Gleason, F. H., Mozley-Standridge, S. E., Porter, D., Boyle, D. G., & Hyatt, A. D. (2007). Preservation of Chytridiomycota in culture collections. *Mycological Research*, 111(2), 129–136. <https://doi.org/10.1016/j.mycres.2006.10.009>
- Glibert, P. M., Middelburg, J. J., McClelland, J. W., & Jake Vander Zanden, M. (2019). Stable isotope tracers: Enriching our perspectives and questions on sources, fates, rates, and pathways of major elements in aquatic systems. *Limnology and Oceanography*, 64(3), 950–981. <https://doi.org/10.1002/lno.11087>
- Gonçalves, M. F. M., Esteves, A. C., & Alves, A. (2022). Marine fungi: Opportunities and challenges. *Encyclopedia*, 2(1), 559–577. <https://doi.org/10.3390/encyclopedia2010037>
- Gonçalves, M. F. M., Hilário, S., Van de Peer, Y., Esteves, A. C., & Alves, A. (2022). Genomic and metabolomic analyses of the marine fungus *Emericellopsis cladophorae*: Insights into saltwater adaptability mechanisms and its biosynthetic potential. *Journal of Fungi*, 8(1), 31. <https://doi.org/10.3390/jof8010031>
- Grossart, H.-P., Van den Wyngaert, S., Kagami, M., Wurzbacher, C., Cunliffe, M., & Rojas-Jimenez, K. (2019). Fungi in aquatic ecosystems. *Nature Reviews Microbiology*, 17(6), 339–354. <https://doi.org/10.1038/s41579-019-0175-8>
- Grossart, H.-P., Wurzbacher, C., James, T. Y., & Kagami, M. (2016). Discovery of dark matter fungi in aquatic ecosystems demands a reappraisal of the phylogeny and ecology of zoospore fungi. *Fungal Ecology*, 19, 28–38. <https://doi.org/10.1016/j.funeco.2015.06.004>
- Gruber, N. (2008). The marine nitrogen cycle: Overview and challenges. *Nitrogen in the Marine Environment*, 2, 1–50.
- Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., et al. (2019). The oceanic sink for anthropogenic CO<sub>2</sub> from 1994 to 2007. *Science*, 363(6432), 1193–1199. <https://doi.org/10.1126/science.aau5153>
- Guo, S.-Y., Jones, E. B. G., Chiang, M. W. L., & Pang, K.-L. (2023). Salinity and temperature affect growth rate of *Alphamyces chaetifer* and *Gorgonomycetes haynaldii* (Chytridiomycota) isolated from coastal habitats of Taiwan. *Botanica Marina*, 66(4), 345–352. <https://doi.org/10.1515/bot-2023-0011>

- Gutiérrez, M. H., Jara, A. M., & Pantoja, S. (2016). Fungal parasites infect marine diatoms in the upwelling ecosystem of the Humboldt current system off central Chile. *Environmental Microbiology*, *18*(5), 1646–1653. <https://doi.org/10.1111/1462-2920.13257>
- Gutiérrez, M. H., Pantoja, S., Quiñones, R. A., & González, R. R. (2010). First record of filamentous fungi in the coastal upwelling ecosystem off central Chile. *Gayana (Concepción)*, *74*(1), 66–73. <https://doi.org/10.4067/s0717-65382010000100010>
- Gutiérrez, M. H., Pantoja, S., Tejos, E., & Quiñones, R. A. (2011). The role of fungi in processing marine organic matter in the upwelling ecosystem off Chile. *Marine Biology*, *158*(1), 205–219. <https://doi.org/10.1007/s00227-010-1552-z>
- Gutiérrez, M. H., Vera, J., Strain, B., Quiñones, R. A., Wörmer, L., Hinrichs, K.-U., & Pantoja-Gutiérrez, S. (2020). Biochemical fingerprints of marine fungi: Implications for trophic and biogeochemical studies. *Aquatic Microbial Ecology*, *84*, 75–90. <https://doi.org/10.3354/ame01927>
- Hagestad, O. C., Hou, L., Andersen, J. H., Hansen, E. H., Altermark, B., Li, C., et al. (2021). Genomic characterization of three marine fungi, including *Emericellopsis atlantica* sp. nov. with signatures of a generalist lifestyle and marine biomass degradation. *IMA Fungus*, *12*(1), 21. <https://doi.org/10.1186/s43008-021-00072-0>
- Hansel, C. M., Zeiner, C. A., Santelli, C. M., & Webb, S. M. (2012). Mn(II) oxidation by an ascomycete fungus is linked to superoxide production during asexual reproduction. *Proceedings of the National Academy of Sciences*, *109*(31), 12621–12625. <https://doi.org/10.1073/pnas.1203885109>
- Hardy, J. T. (1982). The sea surface microlayer: Biology, chemistry and anthropogenic enrichment. *Progress in Oceanography*, *11*(4), 307–328. [https://doi.org/10.1016/0079-6611\(82\)90001-5](https://doi.org/10.1016/0079-6611(82)90001-5)
- Harirchi, S., Rosta, N., Varjani, S., & Taherzadeh, M. J. (2023). 5 - Sampling, preservation, and growth monitoring of filamentous fungi. In M. J. Taherzadeh, J. A. Ferreira, & A. Pandey (Eds.), *Current developments in biotechnology and bioengineering* (pp. 149–180). Elsevier. <https://doi.org/10.1016/B978-0-323-91872-5.00014-4>
- Harrington, B. J., & Hageage, G. J., Jr. (2003). Calcofluor white: A review of its uses and applications in clinical mycology and parasitology. *Laboratory Medicine*, *34*(5), 361–367. <https://doi.org/10.1309/eph2tdt8335gh03>
- Hassett, B. T., Borrego, E. J., Vonnahme, T. R., Rämä, T., Kolomiets, M. V., & Gradinger, R. (2019). Arctic marine fungi: Biomass, functional genes, and putative ecological roles. *The ISME Journal*, *1*(6), 1484–1496. <https://doi.org/10.1038/s41396-019-0368-1>
- Hassett, B. T., Ducluzeau, A.-L. L., Collins, R. E., & Gradinger, R. (2017). Spatial distribution of aquatic marine fungi across the western Arctic and sub-Arctic. *Environmental Microbiology*, *19*(2), 475–484. <https://doi.org/10.1111/1462-2920.13371>
- Hassett, B. T., & Gradinger, R. (2016). Chytrids dominate Arctic marine fungal communities. *Environmental Microbiology*, *18*(6), 2001–2009. <https://doi.org/10.1111/1462-2920.13216>
- Hassett, B. T., Vonnahme, T. R., Peng, X., Jones, E. B. G., & Heuzé, C. (2020). Global diversity and geography of planktonic marine fungi. *Botanica Marina*, *63*(2), 121–139. <https://doi.org/10.1515/bot-2018-0113>
- Hazen, E. L., & Reed, F. C. (1960). Laboratory identification of pathogenic fungi simplified (2nd ed.).
- Heeger, F., Bourne, E. C., Baschien, C., Yurkov, A., Bunk, B., Spröer, C., et al. (2018). Long-read DNA metabarcoding of ribosomal RNA in the analysis of fungi from aquatic environments. *Molecular Ecology Resources*, *18*(6), 1500–1514. <https://doi.org/10.1111/1755-0998.12937>
- Heitger, M., & Baltar, F. (2023). Respiration, production, and growth efficiency of marine pelagic fungal isolates. *Journal of Fungi*, *9*(4), 417. <https://doi.org/10.3390/jof9040417>
- Higgins, S. A., Schadt, C. W., Matheny, P. B., & Löffler, F. E. (2018). Phylogenomics reveal the dynamic evolution of fungal nitric oxide reductases and their relationship to secondary metabolism. *Genome Biology and Evolution*, *10*(9), 2474–2489. <https://doi.org/10.1093/gbe/evy187>
- Hodge, A., & Fitter, A. H. (2010). Substantial nitrogen acquisition by arbuscular mycorrhizal fungi from organic material has implications for N cycling. *Proceedings of the National Academy of Sciences*, *107*(31), 13754–13759. <https://doi.org/10.1073/pnas.1005874107>
- Holinsworth, B., & Martin, J. D. (2009). Siderophore production by marine-derived fungi. *BioMetals*, *22*(4), 625–632. <https://doi.org/10.1007/s10534-009-9239-y>
- Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J. (2010). Prodigal: Prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*, *11*(1), 119. <https://doi.org/10.1186/1471-2105-11-119>
- Ibelings, B. W., De Bruin, A., Kagami, M., Rijkeboer, M., Brehm, M., & Donk, E. V. (2004). Host parasite interactions between freshwater phytoplankton and chytrid fungi (Chytridiomycota) I. *Journal of Phycology*, *40*(3), 437–453. <https://doi.org/10.1111/j.1529-8817.2004.03117.x>
- Ihrmark, K., Bödeker, I. T. M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., et al. (2012). New primers to amplify the fungal ITS2 region – Evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology*, *82*(3), 666–677. <https://doi.org/10.1111/j.1574-6941.2012.01437.x>
- Ilicic, D., & Grossart, H.-P. (2022). Basal parasitic fungi in marine food webs—A mystery yet to unravel. *Journal of Fungi*, *8*(2), 114. <https://doi.org/10.3390/jof8020114>
- Janssen, F., Schrum, C., & Backhaus, J. O. (1999). A climatological data set of temperature and salinity for the Baltic Sea and the North Sea. *Deutsche Hydrographische Zeitschrift*, *51*(9), 5–245. <https://doi.org/10.1007/bf02933676>
- Jeffries, T. C., Curlevski, N. J., Brown, M. V., Harrison, D. P., Doblin, M. A., Petrou, K., et al. (2016). Partitioning of fungal assemblages across different marine habitats. *Environmental Microbiology Reports*, *8*(2), 235–238. <https://doi.org/10.1111/1758-2229.12373>
- Jirout, J. (2015). Nitrous oxide productivity of soil fungi along a gradient of cattle impact. *Fungal Ecology*, *17*, 155–163. <https://doi.org/10.1016/j.funeco.2015.07.003>
- Jobard, M., Rasconi, S., & Sime-Ngando, T. (2010). Fluorescence in situ hybridization of uncultured zoospore fungi: Testing with clone-FISH and application to freshwater samples using CARD-FISH. *Journal of Microbiological Methods*, *83*(2), 236–243. <https://doi.org/10.1016/j.mimet.2010.09.006>
- Johnson, T. W., & Sparrow, F. K. (1961). Fungi in oceans and estuaries.
- Jones, E. B. G. (2011). Are there more marine fungi to be described? *Botanica Marina*, *54*(4), 343–354. <https://doi.org/10.1515/BOT.2011.043>
- Jones, E. B. G., & Pang, K.-L. (2012). *Marine fungi: And fungal-like organisms*. Walter de Gruyter.
- Jones, E. B. G., Pang, K.-L., Abdel-Wahab, M. A., Scholz, B., Hyde, K. D., Boekhout, T., et al. (2019). An online resource for marine fungi. *Fungal Diversity*, *96*(1), 347–433. <https://doi.org/10.1007/s13225-019-00426-5>
- Julianti, E., Abrian, I. A., Wibowo, M. S., Azhari, M., Tsurayya, N., Izzati, F., et al. (2022). Secondary metabolites from marine-derived fungi and actinobacteria as potential sources of novel colorectal cancer drugs. *Marine Drugs*, *20*(1), 67. <https://doi.org/10.3390/md20010067>
- Kagami, M., de Bruin, A., Ibelings, B. W., & Van Donk, E. (2007). Parasitic chytrids: Their effects on phytoplankton communities and food-web dynamics. *Hydrobiologia*, *578*(1), 113–129. <https://doi.org/10.1007/s10750-006-0438-z>
- Kagami, M., Gurung, T. B., Yoshida, T., & Urabe, J. (2006). To sink or to be lysed? Contrasting fate of two large phytoplankton species in Lake Biwa. *Limnology and Oceanography*, *51*(6), 2775–2786. <https://doi.org/10.4319/lo.2006.51.6.2775>

- Kagami, M., Miki, T., & Takimoto, G. (2014). Mycoloop: Chytrids in aquatic food webs. *Frontiers in Microbiology*, *5*, 166. <https://doi.org/10.3389/fmicb.2014.00166>
- Kagami, M., Seto, K., Nozaki, D., Nakamura, T., Wakana, H., & Wurzbacher, C. (2021). Single dominant diatom can host diverse parasitic fungi with different degree of host specificity. *Limnology and Oceanography*, *66*(3), 667–677. <https://doi.org/10.1002/lno.11631>
- Kagami, M., von Elert, E., Ibelings, B. W., de Bruin, A., & Van Donk, E. (2007). The parasitic chytrid, *Zygorhizidium*, facilitates the growth of the cladoceran zooplankton, *Daphnia*, in cultures of the inedible alga, *Asterionella*. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1617), 1561–1566. <https://doi.org/10.1098/rspb.2007.0425>
- Kamei, I., Daikoku, C., Tsutsumi, Y., & Kondo, R. (2008). Saline-dependent regulation of manganese peroxidase genes in the hypersaline-tolerant white rot fungus *Phlebia* sp. Strain MG-60. *Applied and Environmental Microbiology*, *74*(9), 2709–2716. <https://doi.org/10.1128/AEM.02257-07>
- Karl, D. M., & Björkman, K. M. (2015). Chapter 5 - Dynamics of dissolved organic phosphorus. In D. A. Hansell & C. A. Carlson (Eds.), *Biogeochemistry of marine dissolved organic matter* (2nd ed., pp. 233–334). Academic Press. <https://doi.org/10.1016/B978-0-12-405940-5.00005-4>
- Keeling, P. J., Burki, F., Wilcox, H. M., Allam, B., Allen, E. E., Amaral-Zettler, L. A., et al. (2014). The Marine Microbial Eukaryote Transcriptome Sequencing Project (MMETSP): Illuminating the functional diversity of eukaryotic life in the oceans through transcriptome sequencing. *PLoS Biology*, *12*(6), e1001889. <https://doi.org/10.1371/journal.pbio.1001889>
- Kell, D. B. (2004). Metabolomics and systems biology: Making sense of the soup. *Current Opinion in Microbiology*, *7*(3), 296–307. <https://doi.org/10.1016/j.mib.2004.04.012>
- Kempken, F. (2023). Marine fungi: A treasure trove of novel natural products and for biological discovery. *PLoS Pathogens*, *19*(9), e1011624. <https://doi.org/10.1371/journal.ppat.1011624>
- Ketchum, B. H. (1972). *The water's edge: Critical problems of the coastal zone*. MIT Press.
- Kettner, M. T., Oberbeckmann, S., Labrenz, M., & Grossart, H.-P. (2019). The eukaryotic life on microplastics in brackish ecosystems. *Frontiers in Microbiology*, *10*, 538. <https://doi.org/10.3389/fmicb.2019.00538>
- Kettner, M. T., Rojas-Jimenez, K., Oberbeckmann, S., Labrenz, M., & Grossart, H.-P. (2017). Microplastics alter composition of fungal communities in aquatic ecosystems. *Environmental Microbiology*, *19*(11), 4447–4459. <https://doi.org/10.1111/1462-2920.13891>
- Kiliams, E. S., Junges, L., Supraha, L., Leonard, G., Metfies, K., & Richards, T. A. (2020). Chytrid fungi distribution and co-occurrence with diatoms correlate with sea ice melt in the Arctic Ocean. *Communications Biology*, *3*(1), 1–13. <https://doi.org/10.1038/s42003-020-0891-7>
- Kim, S.-K. (2016). *Marine OMICS: Principles and applications*. CRC Press.
- Kimura, H., Fukuba, T., & Naganuma, T. (1999). Biomass of thraustochytrid protists in coastal water. *Marine Ecology Progress Series*, *189*, 27–33. <https://doi.org/10.3354/meps189027>
- Kirstein, I. V., Wichels, A., Krohne, G., & Gerdtz, G. (2018). Mature biofilm communities on synthetic polymers in seawater - Specific or general? *Marine Environmental Research*, *142*, 147–154. <https://doi.org/10.1016/j.marenvres.2018.09.028>
- Klawonn, I., Dunker, S., Kagami, M., Grossart, H.-P., & Van den Wyngaert, S. (2023). Intercomparison of two fluorescent dyes to visualize parasitic fungi (Chytridiomycota) on phytoplankton. *Microbial Ecology*, *85*(1), 9–23. <https://doi.org/10.1007/s00248-021-01893-7>
- Klawonn, I., Van den Wyngaert, S., Iversen, M. H., Walles, T. J. W., Flintrop, C. M., Cisternas-Novoa, C., et al. (2023). Fungal parasitism on diatoms alters formation and bio-physical properties of sinking aggregates. *Communications Biology*, *6*(1), 1–14. <https://doi.org/10.1038/s42003-023-04453-6>
- Klawonn, I., Van den Wyngaert, S., Parada, A. E., Arandia-Gorostidi, N., Whitehouse, M. J., Grossart, H.-P., & Dekas, A. E. (2021). Characterizing the “fungal shunt”: Parasitic fungi on diatoms affect carbon flow and bacterial communities in aquatic microbial food webs. *Proceedings of the National Academy of Sciences*, *118*(23), e2102225118. <https://doi.org/10.1073/pnas.2102225118>
- Kobari, T., Steinberg, D. K., Ueda, A., Tsuda, A., Silver, M. W., & Kitamura, M. (2008). Impacts of ontogenetically migrating copepods on downward carbon flux in the western subarctic Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography*, *55*(14), 1648–1660. <https://doi.org/10.1016/j.dsr2.2008.04.016>
- Kohlmeyer, J., & Kohlmeyer, E. (1979). *Marine mycology: The higher fungi*. Academic Press.
- Kolody, B. C., McCrow, J. P., Allen, L. Z., Aylward, F. O., Fontanez, K. M., Moustafa, A., et al. (2019). Diel transcriptional response of a California Current plankton microbiome to light, low iron, and enduring viral infection. *The ISME Journal*, *13*(11), 2817–2833. <https://doi.org/10.1038/s41396-019-0472-2>
- Korban, S. A., Bobrov, K. S., Maynskova, M. A., Naryzhny, S. N., Vlasova, O. L., Eneyskaya, E. V., & Kulminskaya, A. A. (2017). Heterologous expression in *Pichia pastoris* and biochemical characterization of the unmodified sulfatase from *Fusarium proliferatum* LE1. *Protein Engineering, Design and Selection*, *30*(7), 477–488. <https://doi.org/10.1093/protein/gzx033>
- Korth, F., Deutsch, B., Liskow, I., & Voss, M. (2012). Uptake of dissolved organic nitrogen by size-fractionated plankton along a salinity gradient from the North Sea to the Baltic Sea. *Biogeochemistry*, *111*(1), 347–360. <https://doi.org/10.1007/s10533-011-9656-1>
- Kramer, A., Beck, H. C., Kumar, A., Kristensen, L. P., Imhoff, J. F., & Labes, A. (2015). Proteomic analysis of anti-cancerous scopularide production by a marine *Microascus brevicaulis* strain and its UV mutant. *PLoS One*, *10*(10), e0140047. <https://doi.org/10.1371/journal.pone.0140047>
- Kumar, A., Sørensen, J. L., Hansen, F. T., Arvas, M., Syed, M. F., Hassan, L., et al. (2018). Genome sequencing and analyses of two marine fungi from the North Sea unraveled a plethora of novel biosynthetic gene clusters. *Scientific Reports*, *8*(1), 10187. <https://doi.org/10.1038/s41598-018-28473-z>
- Kumar, V., Sarma, V. V., Thambugala, K. M., Huang, J.-J., Li, X.-Y., & Hao, G.-F. (2021). Ecology and evolution of marine fungi with their adaptation to climate change. *Frontiers in Microbiology*, *12*, 719000. <https://doi.org/10.3389/fmicb.2021.719000>
- Kurata, N., Vella, K., Hamilton, B., Shivji, M., Soloviev, A., Matt, S., et al. (2016). Surfactant-associated bacteria in the near-surface layer of the ocean. *Scientific Reports*, *6*(1), 19123. <https://doi.org/10.1038/srep19123>
- Kutty, S. N., & Philip, R. (2008). Marine yeasts—A review. *Yeast*, *25*(7), 465–483. <https://doi.org/10.1002/yea.1599>
- Kydd, J., Rajakaruna, H., Briski, E., & Bailey, S. (2018). Examination of a high resolution laser optical plankton counter and FlowCAM for measuring plankton concentration and size. *Journal of Sea Research*, *133*, 2–10. <https://doi.org/10.1016/j.seares.2017.01.003>
- Lacerda, A. L. D. F., Proietti, M. C., Secchi, E. R., & Taylor, J. D. (2020). Diverse groups of fungi are associated with plastics in the surface waters of the Western South Atlantic and the Antarctic Peninsula. *Molecular Ecology*, *29*(10), 1903–1918. <https://doi.org/10.1111/mec.15444>
- Lacerda, A. L. D. F., Taylor, J. D., Rodrigues, L. D. S., Kessler, F., Secchi, E., & Proietti, M. C. (2022). Floating plastics and their associated biota in the Western South Atlantic. *Science of the Total Environment*, *805*, 150186. <https://doi.org/10.1016/j.scitotenv.2021.150186>
- Lan, Y., Sun, J., Chen, C., Wang, H., Xiao, Y., Perez, M., et al. (2022). Endosymbiont population genomics sheds light on transmission mode, partner specificity, and stability of the scaly-foot snail holobiont. *The ISME Journal*, *16*(9), 2132–2143. <https://doi.org/10.1038/s41396-022-01261-4>

- Lane, D. M., Valentine, D. L., & Peng, X. (2023). Genomic analysis of the marine yeast *Rhodotorula sphaerocarpa* ETNP2018 reveals adaptation to the open ocean. *BMC Genomics*, 24(1), 695. <https://doi.org/10.1186/s12864-023-09791-7>
- Langvad, F. (1999). A rapid and efficient method for growth measurement of filamentous fungi. *Journal of Microbiological Methods*, 37(1), 97–100. [https://doi.org/10.1016/S0167-7012\(99\)00053-6](https://doi.org/10.1016/S0167-7012(99)00053-6)
- Larsson, D. G. J., & Flach, C.-F. (2022). Antibiotic resistance in the environment. *Nature Reviews Microbiology*, 20(5), 257–269. <https://doi.org/10.1038/s41579-021-00649-x>
- Latva, M., Dedman, C. J., Wright, R. J., Polin, M., & Christie-Oleza, J. A. (2022). Microbial pioneers of plastic colonisation in coastal seawaters. *Marine Pollution Bulletin*, 179, 113701. <https://doi.org/10.1016/j.marpolbul.2022.113701>
- Laughlin, R. J., & Stevens, R. J. (2002). Evidence for fungal dominance of denitrification and codenitrification in a grassland soil. *Soil Science Society of America Journal*, 66(5), 1540–1548. <https://doi.org/10.2136/sssaj2002.1540>
- Lazo-Murphy, B. M., Larson, S., Staines, S., Bruck, H., McHenry, J., Bourbonnais, A., & Peng, X. (2022). Nitrous oxide production and isotopomer composition by fungi isolated from salt marsh sediments. *Frontiers in Marine Science*, 9, 1098508. <https://doi.org/10.3389/fmars.2022.1098508>
- Lepelletier, F., Karpov, S. A., Alacid, E., Le Panse, S., Bigeard, E., Garcés, E., et al. (2014). *Dinomyces arenysensis* gen. et sp. nov. (Rhizophydiales, Dinomycetaceae fam. nov.), a Chytrid Infecting Marine Dinoflagellates. *Protist*, 165(2), 230–244. <https://doi.org/10.1016/j.protis.2014.02.004>
- Leu, A. O., Eppley, J. M., Burger, A., & DeLong, E. F. (2022). Diverse genomic traits differentiate sinking-particle-associated versus free-living microbes throughout the oligotrophic open ocean water column. *mBio*, 13(4), e01569-22. <https://doi.org/10.1128/mbio.01569-22>
- Li, G., Jian, T., Liu, X., Lv, Q., Zhang, G., & Ling, J. (2022). Application of metabolomics in fungal research. *Molecules*, 27(21), 7365. <https://doi.org/10.3390/molecules27217365>
- Li, Q., Liu, D., Jia, Z., Csetenyi, L., & Gadd, G. M. (2016). Fungal biomineralization of manganese as a novel source of electrochemical materials. *Current Biology*, 26(7), 950–955. <https://doi.org/10.1016/j.cub.2016.01.068>
- Li, W., Wang, M., Burgaud, G., Yu, H., & Cai, L. (2019). Fungal community composition and potential depth-related driving factors impacting distribution pattern and trophic modes from epi- to abyssopelagic zones of the western Pacific Ocean. *Microbial Ecology*, 78(4), 820–831. <https://doi.org/10.1007/s00248-019-01374-y>
- Linder, T. (2018). Assimilation of alternative sulfur sources in fungi. *World Journal of Microbiology and Biotechnology*, 34(4), 51. <https://doi.org/10.1007/s11274-018-2435-6>
- Liu, Z., Li, M., Wang, S., Huang, H., & Zhang, W. (2022). Sulfur-containing metabolites from marine and terrestrial fungal sources: Origin, structures, and bioactivities. *Marine Drugs*, 20(12), 765. <https://doi.org/10.3390/md20120765>
- Maeda, K., Spor, A., Edel-Hermann, V., Heraud, C., Breuil, M.-C., Bizouard, F., et al. (2015). N<sub>2</sub>O production, a widespread trait in fungi. *Scientific Reports*, 5(1), srep09697. <https://doi.org/10.1038/srep09697>
- Mahajan, A. S., Fadnavis, S., Thomas, M. A., Pozzoli, L., Gupta, S., Royer, S.-J., et al. (2015). Quantifying the impacts of an updated global dimethyl sulfide climatology on cloud microphysics and aerosol radiative forcing. *Journal of Geophysical Research: Atmospheres*, 120(6), 2524–2536. <https://doi.org/10.1002/2014JD022687>
- Marcelino, V. R., Irinyi, L., Eden, J.-S., Meyer, W., Holmes, E. C., & Sorrell, T. C. (2019). Metatranscriptomics as a tool to identify fungal species and subspecies in mixed communities – A proof of concept under laboratory conditions. *IMA Fungus*, 10(1), 12. <https://doi.org/10.1186/s43008-019-0012-8>
- Martens-Habbena, W., Berube, P. M., Urakawa, H., de la Torre, J. R., & Stahl, D. A. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature*, 461(7266), 976–979. <https://doi.org/10.1038/nature08465>
- Masigol, H., Woodhouse, J. N., van West, P., Mostowfizadeh-Ghalamfarsa, R., Rojas-Jimenez, K., Goldhammer, T., et al. (2021). Phylogenetic and functional diversity of Saprolegniales and fungi isolated from temperate lakes in northeast Germany. *Journal of Fungi*, 7(11), 968. <https://doi.org/10.3390/jof7110968>
- Mayali, X. (2020). NanoSIMS: Microscale quantification of biogeochemical activity with large-scale impacts. *Annual Review of Marine Science*, 12(1), 449–467. <https://doi.org/10.1146/annurev-marine-010419-010714>
- Mescioglu, E., Paytan, A., Mitchell, B. W., & Griffin, D. W. (2021). Efficiency of bioaerosol samplers: A comparison study. *Aerobiologia*, 37(3), 447–459. <https://doi.org/10.1007/s10453-020-09686-0>
- Meyberg, M. (1988). Selective staining of fungal hyphae in parasitic and symbiotic plant-fungus associations. *Histochemistry*, 88(2), 197–199. <https://doi.org/10.1007/BF00493305>
- Montgomery, M. T., Welschmeyer, N. A., & Kirchman, D. L. (1990). A simple assay for chitin: Application to sediment trap samples from the subarctic Pacific. *Marine Ecology Progress Series*, 64(3), 301–308. <https://doi.org/10.3354/meps064301>
- Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., et al. (2013). Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*, 6(9), 701–710. <https://doi.org/10.1038/ngeo1765>
- Morales, S. E., Biswas, A., Herndl, G. J., & Baltar, F. (2019). Global structuring of phylogenetic and functional diversity of pelagic fungi by depth and temperature. *Frontiers in Marine Science*, 6, 131. <https://doi.org/10.3389/fmars.2019.00131>
- Mulholland, M. R., & Lomas, M. W. (2008). Nitrogen uptake and assimilation. In *Nitrogen in the marine environment* (pp. 303–384).
- Nakagiri, A. (2012). Culture collections and maintenance of marine fungi. In *Marine fungi and fungal-like organisms* (pp. 501–508).
- Nakagiri, A., & Jones, E. (2000). Long term maintenance of cultures. *Fungal Diversity Research Series*, 1, 62–68.
- Naranjo-Ortiz, M. A., & Gabaldón, T. (2019). Fungal evolution: Diversity, taxonomy and phylogeny of the fungi. *Biological Reviews*, 94(6), 2101–2137. <https://doi.org/10.1111/brv.12550>
- Neethu, C. S., Saravanakumar, C., Purvaja, R., Robin, R. S., & Ramesh, R. (2019). Oil-spill triggered shift in indigenous microbial structure and functional dynamics in different marine environmental matrices. *Scientific Reports*, 9(1), 1354. <https://doi.org/10.1038/s41598-018-37903-x>
- Neufeld, J. D., Vohra, J., Dumont, M. G., Lueders, T., Manefield, M., Friedrich, M. W., & Murrell, J. C. (2007). DNA stable-isotope probing. *Nature Protocols*, 2(4), 860–866. <https://doi.org/10.1038/nprot.2007.109>
- Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., et al. (2016). FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology*, 20, 241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Nguyen Van Long, N., Vasseur, V., Coroller, L., Dantigny, P., Le Panse, S., Weill, A., et al. (2017). Temperature, water activity and pH during conidia production affect the physiological state and germination time of *Penicillium* species. *International Journal of Food Microbiology*, 241, 151–160. <https://doi.org/10.1016/j.ijfoodmicro.2016.10.022>
- Nilsson, R. H., Wurzbacher, C., Bahram, M., Coimbra, V. R., Larsson, E., Tedersoo, L., et al. (2016). Top 50 most wanted fungi. *Mycosphere*, 12(12), 29–40. <https://doi.org/10.3897/mycokeys.12.7553>
- Oberbeckmann, S., Osborn, A. M., & Duhaime, M. B. (2016). Microbes on a bottle: Substrate, season and geography influence community composition of microbes colonizing marine plastic debris. *PLoS One*, 11(8), e0159289. <https://doi.org/10.1371/journal.pone.0159289>

- Odds, F. C. (1991). Sabouraud's agar. *Journal of Medical and Veterinary Mycology*, 29(6), 355–359. <https://doi.org/10.1080/02681219180000581>
- Odobel, C., Dussud, C., Philip, L., Derippe, G., Lauters, M., Eyheraguibel, B., et al. (2021). Bacterial abundance, diversity and activity during long-term colonization of non-biodegradable and biodegradable plastics in seawater. *Frontiers in Microbiology*, 12, 734782. <https://doi.org/10.3389/fmicb.2021.734782>
- Ollison, G. A., Hu, S. K., Mesrop, L. Y., DeLong, E. F., & Caron, D. A. (2021). Come rain or shine: Depth not season shapes the active protistan community at station ALOHA in the North Pacific Subtropical Gyre. *Deep Sea Research Part I: Oceanographic Research Papers*, 170, 103494. <https://doi.org/10.1016/j.dsr.2021.103494>
- Orsi, W. D., Edgcomb, V. P., Christman, G. D., & Biddle, J. F. (2013). Gene expression in the deep biosphere. *Nature*, 499(7457), 205–208. <https://doi.org/10.1038/nature12230>
- Orsi, W. D., Vuillemin, A., Coskun, Ö. K., Rodriguez, P., Oertel, Y., Niggemann, J., et al. (2022). Carbon assimilating fungi from surface ocean to seafloor revealed by coupled phylogenetic and stable isotope analysis. *The ISME Journal*, 16(5), 1245–1261. <https://doi.org/10.1038/s41396-021-01169-5>
- Ottow, J. C. G. (1972). Rose Bengal as a selective aid in the isolation of fungi and actinomycetes from natural sources. *Mycologia*, 64(2), 304–315. <https://doi.org/10.1080/00275514.1972.12019265>
- Overy, D. P., Rämä, T., Oosterhuis, R., Walker, A. K., & Pang, K.-L. (2019). The neglected marine fungi, *Sensu stricto*, and their isolation for natural products' discovery. *Marine Drugs*, 17(1), 42. <https://doi.org/10.3390/md17010042>
- Paço, A., Duarte, K., da Costa, J. P., Santos, P. S. M., Pereira, R., Pereira, M. E., et al. (2017). Biodegradation of polyethylene microplastics by the marine fungus *Zalerion maritimum*. *Science of the Total Environment*, 586, 10–15. <https://doi.org/10.1016/j.scitotenv.2017.02.017>
- Pang, K.-L., Chiang, M. W.-L., Guo, S.-Y., Shih, C.-Y., Dahms, H. U., Hwang, J.-S., & Cha, H.-J. (2020). Growth study under combined effects of temperature, pH and salinity and transcriptome analysis revealed adaptations of *Aspergillus terreus* NT04989 to the extreme conditions at Kueishan Island Hydrothermal Vent Field, Taiwan. *PLoS One*, 15(5), e0233621. <https://doi.org/10.1371/journal.pone.0233621>
- Pang, K.-L., Jones, E. B. G., Abdel-Wahab, M. A., Adams, S. J., Alves, A., Azevedo, E., et al. (2023). Recent progress in marine mycological research in different countries, and prospects for future developments worldwide. *Botanica Marina*, 66(4), 239–269. <https://doi.org/10.1515/bot-2023-0015>
- Pang, K.-L., Overy, D. P., Jones, E. B. G., Calado, M. D. L., Burgaud, G., Walker, A. K., et al. (2016). 'Marine fungi' and 'marine-derived fungi' in natural product chemistry research: Toward a new consensual definition. *Fungal Biology Reviews*, 30(4), 163–175. <https://doi.org/10.1016/j.fbr.2016.08.001>
- Panno, L., Bruno, M., Voyron, S., Anastasi, A., Gnani, G., Miserere, L., & Varese, G. C. (2013). Diversity, ecological role and potential biotechnological applications of marine fungi associated to the seagrass *Posidonia oceanica*. *New Biotechnology*, 30(6), 685–694. <https://doi.org/10.1016/j.nbt.2013.01.010>
- Peng, X., Fuchsman, C. A., Jayakumar, A., Oleynik, S., Martens-Habbena, W., Devol, A. H., & Ward, B. B. (2015). Ammonia and nitrite oxidation in the Eastern Tropical North Pacific. *Global Biogeochemical Cycles*, 29(12), 2034–2049. <https://doi.org/10.1002/2015GB005278>
- Peng, X., Gat, D., Paytan, A., & Rudich, Y. (2021). The response of airborne mycobiome to dust storms in the Eastern Mediterranean. *Journal of Fungi*, 7(10), 802. <https://doi.org/10.3390/jof7100802>
- Peng, X., & Valentine, D. L. (2021). Diversity and N<sub>2</sub>O production potential of fungi in an oceanic oxygen minimum zone. *Journal of Fungi*, 7(3), 218. <https://doi.org/10.3390/jof7030218>
- Pham, T. T., Dinh, K. V., & Nguyen, V. D. (2021). Biodiversity and enzyme activity of marine fungi with 28 new records from the tropical coastal ecosystems in Vietnam. *Mycobiology*, 49(6), 559–581. <https://doi.org/10.1080/12298093.2021.2008103>
- Philippe, A., Noël, C., Eyheraguibel, B., Briand, J.-F., Paul-Pont, I., Ghiglione, J.-F., et al. (2023). Fungal diversity and dynamics during long-term immersion of conventional and biodegradable plastics in the marine environment. *Diversity*, 15(4), 579. <https://doi.org/10.3390/d15040579>
- Phillips, R., Grelet, G., McMillan, A., Song, B., Weir, B., Palmada, T., & Tobias, C. (2016). Fungal denitrification: *Bipolaris sorokiniana* exclusively denitrifies inorganic nitrogen in the presence and absence of oxygen. *FEMS Microbiology Letters*, 363(4), fnw007. <https://doi.org/10.1093/femsle/fnw007>
- Pilgaard, B., Wilkens, C., Herbst, F.-A., Vuillemin, M., Rhein-Knudsen, N., Meyer, A. S., & Lange, L. (2019). Proteomic enzyme analysis of the marine fungus *Paradendryphiella salina* reveals alginate lyase as a minimal adaptation strategy for brown algae degradation. *Scientific Reports*, 9(1), 12338. <https://doi.org/10.1038/s41598-019-48823-9>
- Piontek, K., Strittmatter, E., Ullrich, R., Gröbe, G., Pecyna, M. J., Kluge, M., et al. (2013). Structural basis of substrate conversion in a new aromatic peroxigenase: Cytochrome P450 functionality with benefits\*. *Journal of Biological Chemistry*, 288(48), 34767–34776. <https://doi.org/10.1074/jbc.M113.514521>
- Pointing, S. B., & Hyde, K. D. (2000). Lignocellulose-degrading marine fungi. *Biofouling*, 15(1–3), 221–229. <https://doi.org/10.1080/08927010009386312>
- Pointing, S. B., Vrijmoed, L. L. P., & Jones, E. B. G. (1998). A qualitative assessment of lignocellulose degrading enzyme activity in marine fungi. *Botanica Marina*, 41(1–6), 293–298. <https://doi.org/10.1515/botm.1998.41.1-6.293>
- Prévost-Bouré, N. C., Christen, R., Dequiedt, S., Mougél, C., Lelièvre, M., Jolivet, C., et al. (2011). Validation and application of a PCR primer set to quantify fungal communities in the soil environment by real-time quantitative PCR. *PLoS One*, 6(9), e24166. <https://doi.org/10.1371/journal.pone.0024166>
- Priest, T., Fuchs, B., Amann, R., & Reich, M. (2021). Diversity and biomass dynamics of unicellular marine fungi during a spring phytoplankton bloom. *Environmental Microbiology*, 23(1), 448–463. <https://doi.org/10.1111/1462-2920.15331>
- Quemener, M., Dayras, M., Frotté, N., Debaets, S., Le Meur, C., Barbier, G., et al. (2021). Highlighting the biotechnological potential of deep oceanic crust fungi through the prism of their antimicrobial activity. *Marine Drugs*, 19(8), 411. <https://doi.org/10.3390/md19080411>
- Quemener, M., Mara, P., Schubotz, F., Beaudoin, D., Li, W., Pachiadaki, M., et al. (2020). Meta-omics highlights the diversity, activity and adaptations of fungi in deep oceanic crust. *Environmental Microbiology*, 22(9), 3950–3967. <https://doi.org/10.1111/1462-2920.15181>
- Raghukumar, S. (2017a). *Fungi in coastal and oceanic marine ecosystems*. Springer International Publishing. <https://doi.org/10.1007/978-3-319-54304-8>
- Raghukumar, S. (2017b). Origin and evolution of marine fungi. In S. Raghukumar (Ed.), *Fungi in coastal and oceanic marine ecosystems: Marine fungi* (pp. 307–321). Springer International Publishing. [https://doi.org/10.1007/978-3-319-54304-8\\_14](https://doi.org/10.1007/978-3-319-54304-8_14)
- Rao, S. (1977). Soil microorganisms and plant growth.
- Rasconi, S., Jobard, M., Jouve, L., & Sime-Ngando, T. (2009). Use of calcofluor white for detection, identification, and quantification of phytoplanktonic fungal parasites. *Applied and Environmental Microbiology*, 75(8), 2545–2553. <https://doi.org/10.1128/AEM.02211-08>
- Rasconi, S., Niquil, N., & Sime-Ngando, T. (2012). Phytoplankton chytridiomycosis: Community structure and infectivity of fungal parasites in aquatic ecosystems. *Environmental Microbiology*, 14(8), 2151–2170. <https://doi.org/10.1111/j.1462-2920.2011.02690.x>

- Rasconi, S., Ptacnik, R., Danner, S., Van den Wyngaert, S., Rohrlack, T., Pilecky, M., & Kainz, M. J. (2020). Parasitic chytrids upgrade and convey primary produced carbon during inedible algae proliferation. *Protist*, *171*(5), 125768. <https://doi.org/10.1016/j.protis.2020.125768>
- Ravishankara, A. R., Daniel, J. S., & Portmann, R. W. (2009). Nitrous oxide (N<sub>2</sub>O): The dominant ozone-depleting substance emitted in the 21st century. *Science*, *326*(5949), 123–125. <https://doi.org/10.1126/science.1176985>
- Rédou, V., Navarri, M., Meslet-Cladière, L., Barbier, G., & Burgaud, G. (2015). Species richness and adaptation of marine fungi from deep-subseafloor sediments. *Applied and Environmental Microbiology*, *81*(10), 3571–3583. <https://doi.org/10.1128/AEM.04064-14>
- Rédou, V., Vallet, M., Meslet-Cladière, L., Kumar, A., Pang, K.-L., Pouchus, Y.-F., et al. (2016). Marine fungi. In L. J. Stal & M. S. Cretoiu (Eds.), *The marine microbiome* (pp. 99–153). Springer International Publishing. [https://doi.org/10.1007/978-3-319-33000-6\\_4](https://doi.org/10.1007/978-3-319-33000-6_4)
- Reinthal, T., Sintes, E., & Herndl, G. J. (2008). Dissolved organic matter and bacterial production and respiration in the sea-surface microlayer of the open Atlantic and the western Mediterranean Sea. *Limnology and Oceanography*, *53*(1), 122–136. <https://doi.org/10.4319/lo.2008.53.1.0122>
- Reñé, A., Timoneda, N., Sarno, D., Piredda, R., Zampicini, G., Zingone, A., et al. (2023). Vertical and temporal distribution of chytrids infecting diatoms in the Gulf of Naples (Italy, Mediterranean Sea). *Marine Ecology*, *44*(3), e12726. <https://doi.org/10.1111/maec.12726>
- Richards, T. A., Jones, M. D. M., Leonard, G., & Bass, D. (2012). Marine fungi: Their ecology and molecular diversity. *Annual Review of Marine Science*, *4*(1), 495–522. <https://doi.org/10.1146/annurev-marine-120710-100802>
- Richards, T. A., Leonard, G., Mahé, F., Del Campo, J., Romac, S., Jones, M. D. M., et al. (2015). Molecular diversity and distribution of marine fungi across 130 European environmental samples. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1819), 20152243. <https://doi.org/10.1098/rspb.2015.2243>
- Riley, J. S., Sanders, R., Marsay, C., Le Moigne, F. A. C., Achterberg, E. P., & Poulton, A. J. (2012). The relative contribution of fast and slow sinking particles to ocean carbon export. *Global Biogeochemical Cycles*, *26*(1), GB1026. <https://doi.org/10.1029/2011GB004085>
- Robbertse, B. (2023). Fungal 28S ribosomal RNA (LSU) RefSeq targeted loci project. <https://doi.org/10.15468/jzfdew>
- Rosenfeld, C. E., Sabuda, M. C., Hinkle, M. A. G., James, B. R., & Santelli, C. M. (2020). A fungal-mediated cryptic selenium cycle linked to manganese biogeochemistry. *Environmental Science & Technology*, *54*(6), 3570–3580. <https://doi.org/10.1021/acs.est.9b06022>
- Rousk, J., Demoling, L. A., & Bååth, E. (2009). Contrasting short-term antibiotic effects on respiration and bacterial growth compromises the validity of the selective respiratory inhibition technique to distinguish fungi and bacteria. *Microbial Ecology*, *58*(1), 75–85. <https://doi.org/10.1007/s00248-008-9444-1>
- Runnel, K., Abarenkov, K., Copo, O., Mikryukov, V., Kõljalg, U., Saar, I., & Tedersoo, L. (2022). DNA barcoding of fungal specimens using PacBio long-read high-throughput sequencing. *Molecular Ecology Resources*, *22*(8), 2871–2879. <https://doi.org/10.1111/1755-0998.13663>
- Saito, M. A., Bertrand, E. M., Duffy, M. E., Gaylord, D. A., Held, N. A., Hervey, W. J. I., et al. (2019). Progress and challenges in ocean metaproteomics and proposed best practices for data sharing. *Journal of Proteome Research*, *18*(4), 1461–1476. <https://doi.org/10.1021/acs.jproteome.8b00761>
- Salazar Alekseyeva, K., Herndl, G. J., & Baltar, F. (2022). Extracellular enzymatic activities of oceanic pelagic fungal strains and the influence of temperature. *Journal of Fungi*, *8*(6), 571. <https://doi.org/10.3390/jof8060571>
- Salkin, I. F., & Hurd, N. (1972). Quantitative evaluation of the antifungal properties of cycloheximide. *Antimicrobial Agents and Chemotherapy*, *1*(3), 177–184. <https://doi.org/10.1128/aac.1.3.177>
- Sánchez Barranco, V., Van der Meer, M. T. J., Kagami, M., Van den Wyngaert, S., Van de Waal, D. B., Van Donk, E., & Gsell, A. S. (2020). Trophic position, elemental ratios and nitrogen transfer in a planktonic host–parasite–consumer food chain including a fungal parasite. *Oecologia*, *194*(4), 541–554. <https://doi.org/10.1007/s00442-020-04721-w>
- Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., et al. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences*, *109*(16), 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Scholz, B., Guillou, L., Marano, A. V., Neuhauser, S., Sullivan, B. K., Karsten, U., et al. (2016). Zoospore parasites infecting marine diatoms – A black box that needs to be opened. *Fungal Ecology*, *19*, 59–76. <https://doi.org/10.1016/j.funeco.2015.09.002>
- Scholz, B., Vyverman, W., Küpper, F. C., Ólafsson, H. G., & Karsten, U. (2017). Effects of environmental parameters on chytrid infection prevalence of four marine diatoms: A laboratory case study. *Botanica Marina*, *60*(4), 419–431. <https://doi.org/10.1515/bot-2016-0105>
- Seifert, K. A., & Rossman, A. Y. (2010). How to describe a new fungal species. *IMA Fungus*, *1*(2), 109–111. <https://doi.org/10.5598/ima fungus.2010.01.02.02>
- Sen, K., Bai, M., Sen, B., & Wang, G. (2021). Disentangling the structure and function of mycoplankton communities in the context of marine environmental heterogeneity. *Science of the Total Environment*, *766*, 142635. <https://doi.org/10.1016/j.scitotenv.2020.142635>
- Sen, K., Sen, B., & Wang, G. (2022). Diversity, abundance, and ecological roles of planktonic fungi in marine environments. *Journal of Fungi*, *8*(5), 491. <https://doi.org/10.3390/jof8050491>
- Senga, Y., Yabe, S., Nakamura, T., & Kagami, M. (2018). Influence of parasitic chytrids on the quantity and quality of algal dissolved organic matter (AOM). *Water Research*, *145*, 346–353. <https://doi.org/10.1016/j.watres.2018.08.037>
- Sérvulo, T., Taylor, J. D., Proietti, M. C., Rodrigues, L. D. S., Puertas, I. P., Barutot, R. A., & Lacerda, A. L. D. F. (2023). Plastisphere composition in a subtropical estuary: Influence of season, incubation time and polymer type on plastic biofouling. *Environmental Pollution*, *332*, 121873. <https://doi.org/10.1016/j.envpol.2023.121873>
- Shoun, H., & Fushinobu, S. (2016). Chapter 14: Denitrification in fungi. In *Metalloenzymes in denitrification* (pp. 331–348). <https://doi.org/10.1039/9781782623762-00331>
- Shoun, H., & Tanimoto, T. (1991). Denitrification by the fungus *Fusarium oxysporum* and involvement of cytochrome P-450 in the respiratory nitrite reduction. *Journal of Biological Chemistry*, *266*(17), 11078–11082. [https://doi.org/10.1016/s0021-9258\(18\)99130-1](https://doi.org/10.1016/s0021-9258(18)99130-1)
- Sime-Ngando, T. (2012). Phytoplankton chytridiomycosis: Fungal parasites of phytoplankton and their imprints on the food web dynamics. *Frontiers in Microbiology*, *3*, 361. <https://doi.org/10.3389/fmicb.2012.00361>
- Smith, C. J., & Osborn, A. M. (2009). Advantages and limitations of quantitative PCR (Q-PCR)-based approaches in microbial ecology. *FEMS Microbiology Ecology*, *67*(1), 6–20. <https://doi.org/10.1111/j.1574-6941.2008.00629.x>
- Sommer, U., Adrian, R., De Senerpont Domis, L., Elser, J. J., Gaedke, U., Ibelings, B., et al. (2012). Beyond the Plankton Ecology Group (PEG) model: Mechanisms driving plankton succession. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 429–448. <https://doi.org/10.1146/annurev-ecolsys-110411-160251>
- Sparrow, F. K. (1969). Zoospore marine fungi from the Pacific Northwest (U.S.A.). *Archiv für Mikrobiologie*, *66*(2), 129–146. <https://doi.org/10.1007/BF00410220>
- Spatafora, J. W., Aime, M. C., Grigoriev, I. V., Martin, F., Stajich, J. E., & Blackwell, M. (2017). The fungal tree of life: From molecular systematics to genome-scale phylogenies. In *The fungal kingdom* (pp. 1–34). John Wiley & Sons, Ltd. <https://doi.org/10.1128/9781555819583.ch1>

- Sridhar, K. R. (2020). Dimensions, diversity and ecology of aquatic mycobiome. *Kavaka*, 54, 10. <https://doi.org/10.36460/kavaka/54/2020/10-23>
- Stanke, M., & Waack, S. (2003). Gene prediction with a hidden Markov model and a new intron submodel. *Bioinformatics*, 19(suppl\_2), ii215–ii225. <https://doi.org/10.1093/bioinformatics/btg1080>
- Stoeck, T., Hayward, B., Taylor, G. T., Varela, R., & Epstein, S. S. (2006). A multiple PCR-primer approach to access the microeukaryotic diversity in environmental samples. *Protist*, 157(1), 31–43. <https://doi.org/10.1016/j.protis.2005.10.004>
- Street, J. H., & Paytan, A. (2005). Iron, phytoplankton growth, and the carbon cycle. *Metal Ions in Biological Systems*, 43, 153–193.
- Stuart, K. A., Welsh, K., Walker, M. C., & Edrada-Ebel, R. (2020). Metabolomic tools used in marine natural product drug discovery. *Expert Opinion on Drug Discovery*, 15(4), 499–522. <https://doi.org/10.1080/17460441.2020.1722636>
- Su, X., Wen, T., Wang, Y., Xu, J., Cui, L., Zhang, J., et al. (2021). Stimulation of N<sub>2</sub>O emission via bacterial denitrification driven by acidification in estuarine sediments. *Global Change Biology*, 27(21), 5564–5579. <https://doi.org/10.1111/gcb.15863>
- Sunda, W. G., & Huntsman, S. A. (1994). Photoreduction of manganese oxides in seawater. *Marine Chemistry*, 46(1), 133–152. [https://doi.org/10.1016/0304-4203\(94\)90051-5](https://doi.org/10.1016/0304-4203(94)90051-5)
- Sutherland, K. M., Wankel, S. D., & Hansel, C. M. (2018). Oxygen isotope analysis of bacterial and fungal manganese oxidation. *Geobiology*, 16(4), 399–411. <https://doi.org/10.1111/gbi.12288>
- Tan, C. S., van Ingen, C. W., & Stalpers, J. A. (2007). Freeze-drying fungi using a shelf-freeze drier. In J. G. Day & G. N. Stacey (Eds.), *Cryopreservation and freeze-drying protocols* (pp. 119–125). Humana Press. [https://doi.org/10.1007/978-1-59745-362-2\\_8](https://doi.org/10.1007/978-1-59745-362-2_8)
- Tang, Y., Zeiner, C. A., Santelli, C. M., & Hansel, C. M. (2013). Fungal oxidative dissolution of the Mn(II)-bearing mineral rhodochrosite and the role of metabolites in manganese oxide formation. *Environmental Microbiology*, 15(4), 1063–1077. <https://doi.org/10.1111/1462-2920.12029>
- Taylor, J. D., & Cunliffe, M. (2014). High-throughput sequencing reveals neustonic and planktonic microbial eukaryote diversity in coastal waters. *Journal of Phycology*, 50(5), 960–965. <https://doi.org/10.1111/jpy.12228>
- Taylor, J. D., & Cunliffe, M. (2016). Multi-year assessment of coastal planktonic fungi reveals environmental drivers of diversity and abundance. *The ISME Journal*, 10(9), 2118–2128. <https://doi.org/10.1038/ismej.2016.24>
- Tebo, B. M., Johnson, H. A., McCarthy, J. K., & Templeton, A. S. (2005). Geomicrobiology of manganese(II) oxidation. *Trends in Microbiology*, 13(9), 421–428. <https://doi.org/10.1016/j.tim.2005.07.009>
- Tedersoo, L., Anslan, S., Bahram, M., Pölme, S., Riit, T., Liiv, I., et al. (2015). Shotgun metagenomes and multiple primer pair-barcode combinations of amplicons reveal biases in metabarcoding analyses of fungi. *MycKeys*, 10, 1–43. <https://doi.org/10.3897/mycokeys.10.4852>
- Tedersoo, L., & Lindahl, B. (2016). Fungal identification biases in microbiome projects. *Environmental Microbiology Reports*, 8(5), 774–779. <https://doi.org/10.1111/1758-2229.12438>
- Tedersoo, L., Tooming-Klunderud, A., & Anslan, S. (2018). PacBio metabarcoding of fungi and other eukaryotes: Errors, biases and perspectives. *New Phytologist*, 217(3), 1370–1385. <https://doi.org/10.1111/nph.14776>
- Teng, Z.-J., Qin, Q.-L., Zhang, W., Li, J., Fu, H.-H., Wang, P., et al. (2021). Biogeographic traits of dimethyl sulfide and dimethylsulfoniopropionate cycling in polar oceans. *Microbiome*, 9(1), 207. <https://doi.org/10.1186/s40168-021-01153-3>
- Tisthammer, K. H., Cobian, G. M., & Amend, A. S. (2016). Global geography of marine fungi is shaped by the environment. *Fungal Ecology*, 19, 39–46. <https://doi.org/10.1016/j.funeco.2015.09.003>
- Todd, J. D., Curson, A. R. J., Dupont, C. L., Nicholson, P., & Johnston, A. W. B. (2009). The *dddP* gene, encoding a novel enzyme that converts dimethylsulfoniopropionate into dimethyl sulfide, is widespread in ocean metagenomes and marine bacteria and also occurs in some Ascomycete fungi. *Environmental Microbiology*, 11(6), 1376–1385. <https://doi.org/10.1111/j.1462-2920.2009.01864.x>
- Trefault, N., De la Iglesia, R., Moreno-Pino, M., Lopes dos Santos, A., Gériques Ribeiro, C., Parada-Pozo, G., et al. (2021). Annual phytoplankton dynamics in coastal waters from Fildes Bay, Western Antarctic Peninsula. *Scientific Reports*, 11(1), 1368. <https://doi.org/10.1038/s41598-020-80568-8>
- Upstill-Goddard, R. C., Frost, T., Henry, G. R., Franklin, M., Murrell, J. C., & Owens, N. J. P. (2003). Bacterioneuston control of air-water methane exchange determined with a laboratory gas exchange tank. *Global Biogeochemical Cycles*, 17(4), 19–19–15. <https://doi.org/10.1029/2003GB002043>
- Vaksmas, A., Knittel, K., Abdala Asbun, A., Goudriaan, M., Ellrott, A., Witte, H. J., et al. (2021). Microbial communities on plastic polymers in the Mediterranean Sea. *Frontiers in Microbiology*, 12, 673553. <https://doi.org/10.3389/fmicb.2021.673553>
- Vaksmas, A., Polerecky, L., Dombrowski, N., Kienhuis, M. V. M., Posthuma, I., Gerritse, J., et al. (2023). Polyethylene degradation and assimilation by the marine yeast *Rhodotorula mucilaginosa*. *ISME Communications*, 3(1), 1–8. <https://doi.org/10.1038/s43705-023-00267-z>
- Vala, A. K., Dave, B. P., & Dube, H. C. (2006). Chemical characterization and quantification of siderophores produced by marine and terrestrial aspergilli. *Canadian Journal of Microbiology*, 52(6), 603–607. <https://doi.org/10.1139/w06-012>
- Van den Wyngaert, S., Ganzert, L., Seto, K., Rojas-Jimenez, K., Agha, R., Berger, S. A., et al. (2022). Seasonality of parasitic and saprotrophic zoospore fungi: Linking sequence data to ecological traits. *The ISME Journal*, 16(9), 2242–2254. <https://doi.org/10.1038/s41396-022-01267-y>
- van Hulst, M., Middag, R., Dutay, J.-C., de Baar, H., Roy-Barman, M., Gehlen, M., et al. (2017). Manganese in the west Atlantic Ocean in the context of the first global ocean circulation model of manganese. *Biogeosciences*, 14(5), 1123–1152. <https://doi.org/10.5194/bg-14-1123-2017>
- Vaulot, D. (2022). MeaPR2. (Version v2.1.1) [Dataset]. <https://doi.org/10.1111/1755-0998.13674>
- Vaulot, D., Sim, C. W. H., Ong, D., Teo, B., Biver, C., Jamy, M., & Lopes dos Santos, A. (2022). metaPR2: A database of eukaryotic 18S rRNA metabarcodes with an emphasis on protists. *Molecular Ecology Resources*, 22(8), 3188–3201. <https://doi.org/10.1111/1755-0998.13674>
- Velez, P., Alejandri-Ramírez, N. D., González, M. C., Estrada, K. J., Sanchez-Flores, A., & Dinkova, T. D. (2015). Comparative transcriptome analysis of the cosmopolitan marine fungus *Corollospora maritima* under two physiological conditions. *G3 Genes/Genomes/Genetics*, 5(9), 1805–1814. <https://doi.org/10.1534/g3.115.019620>
- Vila, T., Frases, S., & Gomes, F. M. (2022). Lessons from protozoans: Phosphate sensing and polyphosphate storage in fungi. *PLoS Pathogens*, 18(3), e1010298. <https://doi.org/10.1371/journal.ppat.1010298>
- Vrijmoed, L. (2000). Isolation and culture of higher filamentous fungi. *Fungal Diversity Research Series*, 1, 1–20.
- Waksman, S. A. (1927). *Principles of soil microbiology*. Williams & Wilkins.
- Walker, G. M., & White, N. A. (2017). Introduction to fungal physiology. In *Fungi* (pp. 1–35). John Wiley & Sons, Ltd. <https://doi.org/10.1002/9781119374312.ch1>
- Wang, G., Wang, X., Liu, X., & Li, Q. (2012). Diversity and biogeochemical function of planktonic fungi in the ocean. In *Biology of marine fungi* (pp. 71–88). Springer. [https://doi.org/10.1007/978-3-642-23342-5\\_4](https://doi.org/10.1007/978-3-642-23342-5_4)
- Wang, M., Mara, P., Burgaud, G., Edgcomb, V., Long, X., Yang, H., et al. (2023). Metatranscriptomics and metabarcoding reveal spatiotemporal shifts in fungal communities and their activities in Chinese coastal waters. *Molecular Ecology*, 32(11), 2750–2765. <https://doi.org/10.1111/mec.16905>
- Wang, X., Singh, P., Gao, Z., Zhang, X., Johnson, Z. I., & Wang, G. (2014). Distribution and diversity of planktonic fungi in the west Pacific warm pool. *PLoS One*, 9(7), e101523. <https://doi.org/10.1371/journal.pone.0101523>



- Wang, Y., Barth, D., Tamminen, A., & Wiebe, M. G. (2016). Growth of marine fungi on polymeric substrates. *BMC Biotechnology*, *16*(1), 3. <https://doi.org/10.1186/s12896-016-0233-5>
- Wang, Y., Sen, B., He, Y., Xie, N., & Wang, G. (2018). Spatiotemporal distribution and assemblages of planktonic fungi in the coastal waters of the Bohai Sea. *Frontiers in Microbiology*, *9*, 584. <https://doi.org/10.3389/fmicb.2018.00584>
- Wang, Y., Sen, K., He, Y., Xie, Y., & Wang, G. (2019). Impact of environmental gradients on the abundance and diversity of planktonic fungi across coastal habitats of contrasting trophic status. *Science of the Total Environment*, *683*, 822–833. <https://doi.org/10.1016/j.scitotenv.2019.05.204>
- Wankel, S. D., Ziebis, W., Buchwald, C., Charoenpong, C., de Beer, D., Dentinger, J., et al. (2017). Evidence for fungal and chemodenitrification based N<sub>2</sub>O flux from nitrogen impacted coastal sediments. *Nature Communications*, *8*(1), 15595. <https://doi.org/10.1038/ncomms15595>
- Ward, N. D., Bianchi, T. S., Medeiros, P. M., Seidel, M., Richey, J. E., Keil, R. G., & Sawakuchi, H. O. (2017). Where carbon goes when water flows: Carbon cycling across the aquatic continuum. *Frontiers in Marine Science*, *4*(January), 1–28. <https://doi.org/10.3389/fmars.2017.00007>
- Watson, A. J., Schuster, U., Shuttler, J. D., Holding, T., Ashton, I. G. C., Landschützer, P., et al. (2020). Revised estimates of ocean-atmosphere CO<sub>2</sub> flux are consistent with ocean carbon inventory. *Nature Communications*, *11*(1), 4422. <https://doi.org/10.1038/s41467-020-18203-3>
- West, P. T., Probst, A. J., Grigoriev, I. V., Thomas, B. C., & Banfield, J. F. (2018). Genome-reconstruction for eukaryotes from complex natural microbial communities. *Genome Research*, *28*(4), 569–580. <https://doi.org/10.1101/gr.228429.117>
- Worden, A. Z., Follows, M. J., Giovannoni, S. J., Wilken, S., Zimmerman, A. E., & Keeling, P. J. (2015). Rethinking the marine carbon cycle: Factoring in the multifarious lifestyles of microbes. *Science*, *347*(6223), 1257594. <https://doi.org/10.1126/science.1257594>
- Wright, R. J., Erni-Cassola, G., Zadjelovic, V., Latva, M., & Christie-Oleza, J. A. (2020). Marine plastic debris: A new surface for microbial colonization. *Environmental Science & Technology*, *54*(19), 11657–11672. <https://doi.org/10.1021/acs.est.0c02305>
- Wright, R. J., Langille, M. G. I., & Walker, T. R. (2021). Food or just a free ride? A meta-analysis reveals the global diversity of the Plastisphere. *The ISME Journal*, *15*(3), 789–806. <https://doi.org/10.1038/s41396-020-00814-9>
- Wurl, O., Bird, K., Cunliffe, M., Landing, W. M., Miller, U., Mustaffa, N. I. H., et al. (2018). Warming and inhibition of salinization at the ocean's surface by cyanobacteria. *Geophysical Research Letters*, *45*(9), 4230–4237. <https://doi.org/10.1029/2018GL077946>
- Wurl, O., & Holmes, M. (2008). The gelatinous nature of the sea-surface microlayer. *Marine Chemistry*, *110*(1), 89–97. <https://doi.org/10.1016/j.marchem.2008.02.009>
- Wurzbacher, C., & Grossart, H.-P. (2012). Improved detection and identification of aquatic fungi and chitin in aquatic environments. *Mycologia*, *104*(6), 1267–1271. <https://doi.org/10.3852/11-225>
- Wurzbacher, C., Larsson, E., Bengtsson-Palme, J., den Wyngaert, S. V., Svantesson, S., Kristiansson, E., et al. (2019). Introducing ribosomal tandem repeat barcoding for fungi. *Molecular Ecology Resources*, *19*(1), 118–127. <https://doi.org/10.1111/1755-0998.12944>
- Xu, W., Luo, Z.-H., Guo, S., & Pang, K.-L. (2016). Fungal community analysis in the deep-sea sediments of the Pacific Ocean assessed by comparison of ITS, 18S and 28S ribosomal DNA regions. *Deep Sea Research Part I: Oceanographic Research Papers*, *109*, 51–60. <https://doi.org/10.1016/j.dsr.2016.01.001>
- Xue, X.-X., Chen, L., & Tang, M.-C. (2022). Genome mining discovery of a new benzazepine alkaloid pseudofisinin A from the marine fungus *Neosartorya pseudofischeri* F27-1. *Antibiotics*, *11*(10), 1444. <https://doi.org/10.3390/antibiotics11101444>
- Yandell, M., & Ence, D. (2012). A beginner's guide to eukaryotic genome annotation. *Nature Reviews Genetics*, *13*(5), 329–342. <https://doi.org/10.1038/nrg3174>
- Yang, S., Xu, W., Zhang, K., Hu, J., Gao, Y., Cui, G., et al. (2022). Fungal communities differ with microplastic types in deep sea sediment enrichments of the Eastern Pacific. *International Biodeterioration & Biodegradation*, *173*, 105461. <https://doi.org/10.1016/j.ibiod.2022.105461>
- Yang, Y., Liu, W., Zhang, Z., Grossart, H.-P., & Gadd, G. M. (2020). Microplastics provide new microbial niches in aquatic environments. *Applied Microbiology and Biotechnology*, *104*(15), 6501–6511. <https://doi.org/10.1007/s00253-020-10704-x>
- Yoch, D. C. (2002). Dimethylsulfoniopropionate: Its sources, role in the marine food web, and biological degradation to dimethylsulfide. *Applied and Environmental Microbiology*, *68*(12), 5804–5815. <https://doi.org/10.1128/AEM.68.12.5804-5815.2002>
- Yu, J., Hu, Q., Xie, Z., Kang, H., Li, M., Li, Z., & Ye, P. (2013). Concentration and size distribution of fungi aerosol over oceans along a cruise path during the fourth Chinese Arctic research expedition. *Atmosphere*, *4*(4), 337–348. <https://doi.org/10.3390/atmos4040337>
- Yunis, A. A. (1988). Chloramphenicol: Relation of structure to activity and toxicity. *Annual Review of Pharmacology and Toxicology*, *28*(1), 83–100. <https://doi.org/10.1146/annurev.pa.28.040188.000503>
- Zäncker, B., Cunliffe, M., & Engel, A. (2021). Eukaryotic community composition in the sea surface microlayer across an east–west transect in the Mediterranean Sea. *Biogeosciences*, *18*(6), 2107–2118. <https://doi.org/10.5194/bg-18-2107-2021>
- Zeghal, E., Vaksmaa, A., Vielfaure, H., Boekhout, T., & Niemann, H. (2021). The potential role of marine fungi in plastic degradation – A review. *Frontiers in Marine Science*, *8*, 738877. <https://doi.org/10.3389/fmars.2021.738877>
- Zeiner, C. A., Purvine, S. O., Zink, E., Wu, S., Paša-Tolić, L., Chaput, D. L., et al. (2021). Mechanisms of Manganese(II) oxidation by filamentous ascomycete fungi vary with species and time as a function of secretome composition. *Frontiers in Microbiology*, *12*, 610497. <https://doi.org/10.3389/fmicb.2021.610497>
- Zettler, E. R., Mincer, T. J., & Amaral-Zettler, L. A. (2013). Life in the “Plastisphere”: Microbial communities on plastic marine debris. *Environmental Science & Technology*, *47*(13), 7137–7146. <https://doi.org/10.1021/es401288x>
- Zhang, F., Wen, Z., Wang, S., Tang, W., Luo, Y.-W., Kranz, S. A., et al. (2022). Phosphate limitation intensifies negative effects of ocean acidification on globally important nitrogen fixing cyanobacterium. *Nature Communications*, *13*(1), 6730. <https://doi.org/10.1038/s41467-022-34586-x>
- Zhang, K., Hu, J., Yang, S., Xu, W., Wang, Z., Zhuang, P., et al. (2022). Biodegradation of polyester polyurethane by the marine fungus *Cladosporium halotolerans* 6UPA1. *Journal of Hazardous Materials*, *437*, 129406. <https://doi.org/10.1016/j.jhazmat.2022.129406>
- Zhao, H., Brearley, F. Q., Huang, L., Tang, J., Xu, Q., Li, X., et al. (2023). Abundant and rare taxa of planktonic fungal community exhibit distinct assembly patterns along coastal eutrophication gradient. *Microbial Ecology*, *85*(2), 495–507. <https://doi.org/10.1007/s00248-022-01976-z>
- Zhao, Z., Baltar, F., & Herndl, G. J. (2020). Linking extracellular enzymes to phylogeny indicates a predominantly particle-associated lifestyle of deep-sea prokaryotes. *Science Advances*, *6*(16), eaaz4354. <https://doi.org/10.1126/sciadv.aaz4354>
- Zuo, X., Xu, W., Wei, S., Jiang, S., Luo, Y., Ling, M., et al. (2023). Aerobic denitrifying bacterial-fungal consortium mediating nitrate removal: Dynamics, network patterns and interactions. *iScience*, *26*(6), 106824. <https://doi.org/10.1016/j.isci.2023.106824>