04 University of Plymouth Research Theses

01 Research Theses Main Collection

2024

# Predictors of Weight Gain in the Rehabilitation of Eastern North Atlantic Grey Seals (Halichoerus grypus)

# McGivern, Samantha

https://pearl.plymouth.ac.uk/handle/10026.1/22590

http://dx.doi.org/10.24382/5223 University of Plymouth

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Samantha McGivern; Predictors of Weight Gain in the Rehabilitation of Eastern North Atlantic Grey Seals (*Halichoerus grypus*).

# **COPYRIGHT STATEMENT**

Copyright and Moral rights arising from original work in this thesis and (where relevant), any accompanying data, rests with the Author unless stated otherwise<sup>1</sup>.

Re-use of the work is allowed under fair dealing exceptions outlined in the Copyright, Designs and Patents Act 1988 (amended)<sup>2</sup>, and the terms of the copyright licence assigned to the thesis by the Author.

In practice, and unless the copyright licence assigned by the author allows for more permissive use, this means,

- that any content or accompanying data cannot be extensively quoted, reproduced or changed without the written permission of the author / rights holder; and
- that the work in whole or part may not be sold commercially in any format or medium without the written permission of the author/rights holder.

Any third-party copyright material in this thesis remains the property of the original owner. Such third party copyright work included in the thesis will be clearly marked and attributed, and the original licence under which it was released will be specified. This material is not covered by the licence or terms assigned to the wider thesis and must be used in accordance with the original licence; or separate permission must be sought from the copyright holder.

The author assigns certain rights to the University of Plymouth including the right to make the thesis accessible and discoverable via the British Library's Electronic Thesis Online Service (ETHOS) and the University research repository, and to undertake activities to migrate, preserve and maintain the medium, format and integrity of the deposited file for future discovery and use.

 $<sup>^{1}</sup>$  *E.g.* in the example of third party copyright materials reused in the thesis.

<sup>&</sup>lt;sup>2</sup> In accordance with best practice principles such as, *Marking/Creators/Marking third party content* (2013). Available from:

https://wiki.creativecommons.org/wiki/Marking/Creators/Marking\_third\_party\_content [accessed 28<sup>th</sup> February 2022]



# PREDICTORS OF WEIGHT GAIN IN THE REHABILITATION OF EASTERN NORTH ATLANTIC

# GREY SEALS (HALICHOERUS GRYPUS).

by

SAMANTHA MCGIVERN

A thesis submitted to the University of Plymouth in partial fulfilment for the degree of

**RESEARCH MASTERS** 

School of Biological and Marine Sciences.

April 2024

Samantha McGivern; Predictors of Weight Gain in the Rehabilitation of Eastern North Atlantic Grey Seals (*Halichoerus grypus*).

#### Acknowledgements:

I would like to thank my supervisory team for all of their support and advice during this process. I want to specially thank Dr K. Herborn and Dr C. Boschetti for all the extra support and advice they have provided in the last 18 months.

I would like to thank Seal Rescue Ireland (S.R.I.) and Duncan Kenny for their incredible support, involvement, and resourcefulness within this project. To Seal Rescue Ireland, it has been an honour to work alongside you. A special thank you to Hayleigh at the S.R.I., you have been incredibly patient and amazing to export that data from your software and get it over to me. To Duncan Kenny, without your tenacity, knowledge of cameras, and incredible knowledge of seals, I wouldn't have collected an amazing behavioural dataset nor have learnt so much about identifying grey seals. I still cannot believe you can identify them from head shape alone!

I would additionally like to thank my partner, Alex, for all of your support, patience, and wit. You have kept me going through all the hardship and challenges this project has subjected me too. I would also like to thank two fellow research master's students: Kathryn Wiltshire and Victoria Wakeham. Kathryn, you have been incredible, and I have learnt so much about seal genetics from you as our projects interwove so much. You have been incredibly supportive and thank you for helping to collect the behavioural data. To Vikki, we have spent many a time being each other's therapist. We finally got there in the end!

### Author's Declaration:

At no time during the registration for the degree of Research Masters in Biological Sciences has the author been registered for any other University award without prior agreement of the Doctoral College Quality Sub-Committee.

Work submitted for this research degree at the University of Plymouth has not formed part of any other degree either at the University of Plymouth or at another establishment.

A programme of advanced study was undertaken, which included taught modules taken (BIOL5131: Postgraduate Research Skills & Methods. ANIM5010: Animal Behaviour and Welfare Research).

The following external institutions were visited for research and consultation purposes:

Seal Rescue Ireland, Courtown, Ireland.

The Cove Project, Brixham, Devon, U.K. (specifically, Duncan Kenny).

Presentations at conferences:

University of Plymouth STEM Showcase (26<sup>th</sup> January 2024)

Word count of main body of thesis: 10045

Signed: SMcGivern

Date: 11/04/2024

Samantha McGivern; Predictors of Weight Gain in the Rehabilitation of Eastern North Atlantic Grey Seals (*Halichoerus grypus*).

#### <u>Abstract:</u>

As large adult body size is subjected to strong selective pressures, other processes can be compromised before growth when there is resource limitation or stress during the growth phase. Stress has been linked with both weight loss and weight gain in a variety of taxa, and some stressors, such as physical disturbance, are known to affect the development of important physiological processes, e.g., diving physiology in marine mammals. With 40% of the world's grey seal (Halichoerus grypus) population on UK shores, grey seal pups are frequently admitted to rehabilitation centres within their first four weeks – a crucial growth phase – across the UK. Common rehabilitation practices (e.g., change in environment, social grouping, and disturbance) are known to cause stress to individuals in other contexts. Within grey seal rehabilitation, there are known factors outside the control of the centre that affect weight gain and survival, e.g., being underweight upon admittance. Yet, there are management practices, such as movement of individuals between pools (requires handling, movement, and regrouping) that may cause stress. The effects of these management practices on weight gain and mortality is unknown. By studying the behaviour of individuals, it can provide insights into how or why there are any changes in weight gain and mortality. This study investigates impacts of factors known to affect rehabilitation outcomes (e.g., admittance reason and intake weight) on weight gain and mortality in one rehabilitation centre over six pupping seasons. In a case study year, where data were available, this study also investigates impacts of unexplored factors: pool design and movement between pools, on individuallevel weight gain (<70 seals) and behaviour (<39 seals). No evidence was found that management practices (pool characteristics, pool movements) affected weight gain, which was explained by duration of rehabilitation, and admittance reasons: underweight and injured. Pools varied in movement, rest, and head elevated behaviour (may indicate disturbance), suggesting some characteristics such as the visual blockade, might modify stress exposure or energetic costs during growth. No clear pattern across pools with respect to weight emerged, suggesting that although there are indicators of disturbance it might not be affecting weight gain.

Samantha McGivern; Predictors of Weight Gain in the Rehabilitation of Eastern North Atlantic Grey Seals (*Halichoerus grypus*).

# Contents:

Tables and Illustrations			Page 5						
1. Introduction			Page 6						
	1.1: W	'eight ar	nd Stress						Page 7
	1.2: Gi	rey Seal	Phenology						Page 10
	1.3: Re	ehabilita	ition Centre	S					Page 14
	1.4: Tł	nis Study	/						Page 21
2. Met	hodolo	gy							Page 23
	2.1: Re	ehabilita	ition Centre						Page 23
	2.2: W	'eight ar	nd Mortality						Page 25
	2.3: Be	ehaviour	r						Page 27
	2.4: Da	ata Anal	ysis						Page 30
		2.4.1: \	Weight Char	nge					Page 30
		2.4.2: 1	Mortality						Page 31
		2.4.3: [	Behaviour						Page 31
	2.5: Et	hics							Page 32
3. Resu	ults								Page 33
	3.1: W	eight/							Page 33
		3.1.1:	Variables	Influencing	Weight	Change	for	Six	
		Consec	cutive Puppi	ng Seasons.					Page 33

3.1.2: Variables Influencing Weight Change for Case Study	
Pupping Season	Page 34
3.2: Mortality	Page 34
3.2.1: Variables Influencing Mortality for Six Consecutive	
Pupping Season	Page 34
3.3: Behaviour	Page 35
4. Discussion	Page 37
4.1: Impact of Management Practices	Page 37
4.2: Impact of Management Practices Outside the Rehabilitation	
Centre's Control	Page 42
4.3: Individuals within Rehabilitation	Page 45
4.4: Recommendations and Further Research	Page 47
References	Page 49
Appendix	Page 59
A.1: Behavioural Models	Page 59
A.2: Justification for Removing Admittance Reasons from the Model	Page 61
A.3: Six Consecutive Pupping Seasons Main Weight Results Figures	Page 62
A.4: Case Study Pupping Season Main Weight Results Figure	Page 63
Supplementary Information	Page 64
S.1: Definitions of Reasons for Admittance	Page 64
S.2: Camera Locations	Page 65

S.3: AIC Numl	bers for Comparison of Behavioural Models	Page 66
S3.1: (	Combined Behavioural Dataset	Page 66
S3.2: I	Day Behavioural Dataset	Page 66
S3.3: I	Night Behavioural Dataset	Page 67
S.4: Behaviou	Iral Model Results (Including Repeatability)	Page 68
S4.1: [	Day Data	Page 68
	S4.1.1: Movement Behaviour	Page 68
	S4.1.2: Rest Behaviour	Page 69
	S4.1.3: Head Elevated Behaviour	Page 70
S4.2: I	Night Data	Page 71
	S4.2.1: Movement Behaviour	Page 71
	S4.2.2: Rest Behaviour	Page 71
	S4.2.3: Head Elevated Behaviour	Page 72
S4.3: (	Combined Night and Day	Page 73
	S4.3.1: Movement Behaviour	Page 73
	S4.3.2: Rest Behaviour	Page 73
	S4.3.3: Head Elevated Behaviour	Page 74

# Tables and Illustrations:

# Tables:

Table 1: The natural progression of grey seals (Halichoerus grypus) from birth	
to adulthood.	Page 11
Table 2: Attributes for four pools at Seal Rescue Ireland, Courtown, Wexford,	
Ireland.	Page 24
Table 3: Behavioural ethogram created for data collection on time-lapse	
images of seals.	Page 28
Table 4: Contextual variables considered to affect the behaviour of	
individuals across pools.	Page 29
Table 5: Variables influencing weight change over six consecutive pupping	
season.	Page 33
Table 6: Variables influencing weight change for case study pupping season.	Page 34
Table 7: Variables influencing mortality over six consecutive pupping season	Page 35
Figures:	
Figure 1: Adapted from Silva et al., 2020, and Spotte and Stake, 1982.	Page 13
Figure 2: Seal rehabilitation process example; Seal Rescue Ireland (S.R.I.	),
Courtown, Wexford, Ireland.	Page 20
Figure 3: Aerial view of the Seal Rescue Ireland site.	Page 24
Figure 4: The likelihood of different factors influencing three grey seal	
(Halichoerus grpyus) behaviours.	Page 36

#### 1. Introduction:

Rehabilitation centres have become a key part of conserving wild populations (Aitken, 2004), mitigating the effects on populations from human disturbance and severe weather events, by rehabilitating and releasing individuals back into the wild (Cope et al., 2022). Grey seal (Halichoerus grypus) pups often enter rehabilitation centres within their first four weeks (e.g., 21% of individuals admitted to Seal Rescue Ireland, 2018 -2024) - a crucial growth phase for surviving the post-weaning period (Noren et al., 2008). Therefore, it is imperative that we understand the implications of management practices on growth. A recent review showed that there are many known factors that could affect weight gain and pup survival (e.g., intake weight, weight gain, admittance reasons, age, sex, and pupping seasons), providing management recommendations on protocols during the triage and treatment phase (Zatrak et al., 2022). However, there could be unknown management factors (e.g., routine movement between pools (part of standard management in captivity), other pool disturbances (e.g., staff within the pools and water drainage as part of cleaning (part of standard management in captivity), and other wildlife present), and duration of rehabilitation that are not considered which could be impacting weight gain and survival of grey seal pups. Individuals are routinely moved between pools as part of standard management in captivity and it largely remains unknown what the impact of these moves have on individuals versus the variation in disturbance levels across pools within centres. This thesis will explore both these known and unknown factors to understand if there are further practices that should be considered when rehabilitating wild grey seal pups.

The introduction will cover relationships between weight and stress, hence how stress during management might affect the mortality and growth of seals. A further

explanation on the relevant aspects of seal phenology and physiology, and an introduction into rehabilitation centres (management practices, characteristics, and grey seal rehabilitation).

1.1: Weight and Stress:

Growth rate, according to life history theory, can be subjected to strong directional selection i.e., the reproductive and survival advantages associated with large body size including fecundity, mate selection, predation avoidance, and offspring survival (Dmitriew, 2011). However, optimal growth is seldom maximal, shaped by selection pressures to reduce the negative physiological impacts of rapid growth, such as reduced immune function due to resource allocation to growth, and reduced resistance to physiological stressors, for example drought causes decreased resistance to elevated sodium concentrations (Mangel and Stamps, 2001)

As growth is subjected to strong selective pressures that favour fast growth; other processes are often observed to be compromised before growth when there is resource limitation. The costs associated with rapid growth are not evident until specifically after reaching the reproductive phase (Hector and Nakagawa, 2012). For example, there is cumulative damage and development flaws evident in faster cellular senescence (Ben-Porath and Weinberg, 2004). In European shags (*Phalacrocorax aristotelis*), high levels of stress during early post-natal life has potentially been mechanistically linked to increased shorten telomeres in nestlings, reducing longevity (Herborn et al., 2014). Herborn et al., (2014) also found that it provided, at minimum, evidence of oxidative damage resulting from oxidative stress. Higher rates of instantaneous mortality is another potential cost of rapid growth. In three-spined stickleback fish, rapid growth has

been shown to also reduce the lifespan through rapid skeletal growth (Inness and Metcalfe, 2008).

There are environmental constraints during growth that can also impact the pattern of growth with immediate and long-term impacts for animals. Food deprivation is a prime example of this (Moe et al., 2004) but stress is a further example. When an individual experiences a stressor, glucocorticoid hormones, such as cortisol, and corticosterone (dependent on species), are released to facilitate the response to the stressor and then to return the body to baseline function. Yet, if an individual receives prolonged or recurrent stress exposure this can disrupt homeostasis, leading to stress-induced diseases (Romero and Beattie, 2022). Stress has been linked with both weight loss and weight gain in a variety of taxa; mammals (Kannan et al., 2000; Reeder and Kramer, 2005; Schuamann et al., 2014; Kershaw and Hall., 2016, Malik and Spencer, 2019), birds (Moe et al., 2004; Dickens et al., 2009), amphibians (Crespi and Denver, 2004), and reptiles (Dunlap, 1995). It has been noted that early life stress, including maternal gestational and juvenile stress, can cause weight gain as it can alter feeding and metabolism (Malik and Spencer, 2019). In female sheep, when corticosteroids are given at the end stage of pregnancy, it leads to increased weight and fat masses in the adult offspring (Berry et al., 2013). Repeated exposure to stressors in late pregnancy can also lead to metabolic vulnerability to a high fat or a high fat-high sugar diet in adult offspring (Malik and Spencer, 2019). Malik and Spencer (2019) further suggested that parental or combined parental influence is important in the programming of metabolic and hypothalamus-pituitary-adrenal (HPA) axis outcomes. Early life stress in juveniles can cause alterations to the limbic system. In rats, early life stress can cause juveniles to perform poorly in spatial learning tasks (Avital et al., 2006). In rodent pups, when the dam alters her behaviour due to a stressor (e.g., the loss of nesting material (Ivy et al.,

2008)), the stressor leads to detrimental long-term metabolic outcomes of the pup even with access to an adequate milk supply from the dam. The pups exhibit persistent reductions in white adipose tissue mass but experience long-term weight regulation difficulties with increased weight gain and body fat when exposed to unhealthy diets later in life (Yam et al., 2017). Within marine mammals, there is further evidence of parental stress exposure impacting offspring weight gain through glucocorticoids passing through lactation (Stead et al., 2021; Armstrong et al., 2023).

Stress can have a direct effect on food intake, i.e., weight, but it can also have indirect effects through energetic costs, i.e., diversion of resources. Stress-exposed individuals can respond by increased food intake hence weight gain or through appetite suppression hence weight loss (Reeder and Kramer, 2005; Malik and Spencer, 2019), seen in many species such as fish (Bernier, 2006), toads (Crespi and Denver, 2004), and mammals (Harris et al., 1998; Maniam and Morris, 2012). Another way stress can affect weight is through time budget and energetic costs of lower rest and heightened vigilance. Rest is involved in the regulation of body weight, and the amount of rest and synchronisation of the biological clock are necessary for energy balance and the secretion of hormones that contribute to weight regulation (Leger et al., 2015). Sleep deprivation has been shown to cause weight change in both directions. In humans, it has been shown to cause weight gain through increased food intake (Chaput and Tremblay, 2012), but in mice, sleep deprivation is associated with weight loss as the amount of food eaten during sleep deprivation was insufficient to cover the energy expenditure further leading to a metabolic deficit (Đukanović et al., 2022). Complete rest is important, but fragmented sleep, in humans, has been associated with impaired alertness, memory, mood regulation, changes to brain activity and metabolism (Short and Banks, 2013).

Environmental stressors, such as physical disturbance, can also play a part in regulating behaviour. Social and physical disturbance has been shown to alter the behaviour of Australian field crickets (*Teleogryllus oceanicus*). Rudin et al., (2018) found that individuals switched from a silent environment to an acoustic environment became less bold, active, and explorative. However, when moved from an undisturbed to disturbed environment, the crickets only became less explorative, indicating that changes within the social environment had a greater effect. In meerkats (*Suricata suricatta*), there is some evidence that an increase in vigilance may occur due to previously experienced stressors through an increase in glucocorticoids, potentially causing weight loss (Voellmy et al., 2014). Evidence of these behaviours and how they might be impacting weight, are important welfare indicators that may be particularly relevant in the rehabilitation of grey seals.

#### 1.2: Grey Seal Phenology:

There are three stages in the annual cycle of the eastern north Atlantic grey seal: breeding season (August to December), moulting (December to April), and a period of time feeding and hauled out either on land, ice or at sea (Lidgard, 2003; In the UK, these are public available records from North Wales Wildlife Trust, 2021a, 2021b, 2021c). Female grey seals become sexually mature at five years old, reproducing annually. At most breeding sites, there are overlapping generations as grey seals are heroparous capital breeders with a high degree of site fidelity (Bull et al., 2017). Females give birth to one pup a year weighing 10 – 15kgs on average (Based on publicly available records from Cumbria Wildlife Trust, 2023). Grey seal pups progress through four different phases before becoming a juvenile (Table 1). Grey seal pups are not water resistant at birth and are unable to swim due to the lanugo (white) coat. Grey seals have a threeweek nursing period, where for the first 16 – 20 days, they rely solely upon the mother's milk to build up strength and fat reserves (feeding up to six times a day) (Based on public available records from Cumbria Wildlife Trust, 2023). The milk transferred from the mother to the pup contains fat (50%), protein (12%), and water (36.2%) (Baker, 1990). During the lactation period, female grey seals can lose 84% of stored energy reserves, with the pup gaining 46% of the mass lost by the mother (Reidman, 1990a). At three weeks, the pup is abandoned by the mother, weighing between 30 and 40kgs. During the larger the fat reserves, the better the chance the seal pup has to survive (Hall et al., 2008). The diet post-weaning mainly consists of fish such as salmon, cod, herring, mackerel, cephalopods (such as squid), and crustaceans (Reidman, 1990b).

Table 1: The natural progression of grey seals (*Halichoerus grypus*) from birth to adulthood. Age classes are a functional way of aging pups from within each pupping season. In southwest England, pups appear to complete the moult before weaning (Barnett et al., 2000). Table based upon data gathered from Barnett et al., (2000) and Zatrak et al., (2022). These age classes are also applied within rehabilitation centres.

Age Class	Phase	Age	Description
	Phase 1	0 – 2 days	Umbilicus is pink, moist, and bloody. Neonatal lanugo coat is present.
Neonate		3 – 4 days	Umbilicus is pink and dry at the top. Neonatal lanugo coat is present.
		5 – 6 days	Umbilicus is dry, shrivelled, and grey/black in colour. Neonatal lanugo coat present.
White-coat	Phase 2	7 – 14 days	Umbilicus lost, neonatal lanugo coat present.
Mid-moults	Phase 3	11 - 16 days	Lanugo coat partially lost.
Moulted	Phase 4	11 days – 10	Lanugo coat lost.
		months old	16 – 21 days weaning occurs.
			At 21 days, mother abandons the pup and learning phase commences.

Juveniles	1 – 2 years	During second moult, fur loses structure and becomes orangey brown. Remain solitary or with other juveniles.
Adolescent	3 – 5 years	Behaviour is learned though interactions with adults and other adolescents.
Adult	>5 years	Reached full size and weight. Females start breeding between five and six.

Wild grey seals also do not follow a linear weight trajectory throughout their lifespan (Figure 1) due to important life history processes and seasonal environmental dynamics such as reproduction and moulting (Figure 1C) (Silva et al., 2020). Pup growth is faster during the weaning period before decreasing post-weaning, before slowly increasing again (Figure 1B). Weight gain decreases post-weaning due to the required foraging skill acquisition. In captive grey seals, growth is much slower, gradually catching up to their wild counterparts (Figure 2).



Figure 1: Adapted from Silva et al., 2020, and Spotte and Stake, 1982. Simulated growth data from a lifecycle dynamic energy budget (DEB) model vs. empirical observations of female grey seal body growth. (A) Body weight during foetal growth. (B) Body weight of weaning and juveniles. Points represent empirical data from five studies, solid lines represent the mean simulated body growth and dashed lined represent the range (min – max) of the simulated data (Silva et al., 2020). Weaning phase is between 0 and 21 days, post weaning occurs after 21 days. (C) Weight gain of two captive-reared grey seal (*Halichoerus grypus*) pups, compared with wild pups on Sable Island, Nova Scotia (Spotte and Stake, 1982).

Phocid pups in general have a high first year mortality rate (Hall et al., 2002; Harding et

al., 2005; Hall et al., 2008; Jenssen et al., 2010), with 43% mortality of juvenile grey seals

reported in the Netherlands over a 30-year period (1988 and 2002 excluded due to high

mortality caused by phocine distemper virus (Phocine morbillivirus) epidemic) (Osinga

et al., 2012). There is evidence of a positive correlation between weight at weaning and

survival post-weaning (Hall et al., 2008; Jenssen et al., 2010), potentially stronger in male

pups (Hall et al., 2008). At two breeding sites in the U.K., the primary cause of pup mortality was starvation (19.6%), but other causes included septicaemia (14%), peritonitis (12%), and drowning (12%) (Anderson et al., 1979). There is further evidence of a wider range of anthropogenic effects on grey seals, including entanglement within fishing lines and bycatch (Sayer et al., 2021). During a 10-year period (2010 – 2020), 81.4% of grey seals were reported as entangled, with young grey seals the most susceptible (Salazar-Casals et al., 2022). Salazar-Casals et al., (2022) reported that entanglements have gradually been increasing from 2017, with a spike after 2018. Other evidence has shown that maternal condition is a significant factor in the likelihood of pup survival. Mellish et al., (1999) found clear evidence that heavier females not only lactated for longer and had higher milk outputs, but also produced larger pups at weaning. Further evidence suggests that due to the increased frequency of severe winter storms within the last five years, 70% of seal pups die within their first year, with large numbers of pups being separated prematurely from their mothers (Jarvis and Marsh, 2017). Jarvis and Marsh (2017) further reported that wild grey seal pups were often underweight due to the inability to feed effectively in the post-weaning period.

1.3: Rehabilitation Centres:

Rehabilitation centres, or rescue centres, are centres that rescue and take in injured or ailing marine mammals such as seals, rehabilitate them and then release them back into the wild. Rehabilitation centres employ a variety of management practices to increase the likelihood of post-release survival and to reduce the impact on their welfare, including the replication of wild environment (e.g., enrichment, haul-out areas, access to water of varying depths), to meets the animals physiological and psychological needs, and the type of human interaction (British Veterinary Zoological Society, 2016). The

majority of rehabilitation centres are designed to meet a minimum set of criteria in weight and health before release (Burroughes et al, 2021), but there are rehabilitation centres such as Seal Rescue Ireland that design practices to increase the likelihood of post-release survival. Most rehabilitation centres follow a set pathway; a hospital or intensive care unit (ICU) for initial care and treatment, a recovery area (sometimes referred to as Kennels), pools or enclosures for social and/or physical development, and an area with reduced or removed human interaction pre-release. Wild animals enter rehabilitation centres for a variety of reasons, including injury, illness, entanglement (often in human made objects such as netting), and human disturbance. These reasons as well as the condition (e.g., weight) on entering the centre, can all impact management decisions as well as the likelihood of survival both pre- and post-release.

All rehabilitation centres have a large database of information recorded on individuals upon admittance. This information can be used for a variety of purposes, including tracking of populations, disease outbreaks (e.g., phocine distemper (*Morbillivirus*) outbreak in 2002 (Härkönen et al., 2006)), changes in admittance patterns for particular species (e.g., if there are more entanglement cases in one particular year) (Zatrak et al., 2022), reviewing management practices (Burroughes et al., 2021), and further factors that could be influencing survival. European hedgehogs (*Erinaceus europaeus*) are a great example as this species is frequently seen in rehabilitation centres; the Royal Society for the Prevention of Cruelty to Animals (RSPCA) saw over 19 thousand hedgehogs admitted over a 13-year period (Burroughes et al., 2021). Due to the plethora of data available from this period, it allowed for management care practices to be greatly improved (e.g., through increased experience and expertise of rehabilitation staff), increasing the number of hedgehogs released (Burroughes et al., 2021). A survey by Guy et al., (2013) on current mammal rehabilitation and release practices, found that

three primate rehabilitation centres reported habituation to humans as a potential problem. A very recent study on orphaned lynx in rehabilitation centres are another example as habituation to humans developed during the rehabilitation process, especially as they associated humans with food (Molinari-Jobin et al., 2024).

Although there is an extensive database available, in phocids, there is less research into management practices that could be influencing survival. If this data was harnessed, management decisions and practices would be better informed. Evidence of this has been seen in a recent study by Zatrak et al., (2022). This study evidenced that weight upon admittance of juvenile seals had a significant effect on survival; juvenile seals surviving to release increased by 1.07 times for every kilogram of weight over the predicted weight for their age. This was consistent with another study by Hall et al. (2002) where upon a heavier juvenile had higher survival probabilities. Zatrak et al., (2022) recommended that weight during the triage and treatment phase should be paid special attention to, making sure that individuals do not lose weight and therefore, decreasing their chance of survival.

It is estimated that 40% of the world's population of grey seals are in UK and Ireland (Zatrak et al., 2022). Within the U.K. and Ireland, there are 13 specialist seal rescue centres and hospitals, and the British Divers Marine Life Rescue (BDMLR) reported that the total rehabilitation capacity within the U.K. often reached its limit during peak pupping season, due to severe storms, reduced ability to feed effectively post-weaning, and an increase in human disturbance (Jarvis and Marsh, 2017). In a recent review of grey seal and harbour seal admittances between 1988 and 2020 and 4126 individuals (2691 grey seals and 1435 harbour seals) over five rehabilitation centres (Zatrak et al., 2022), the most common nonexclusive reasons have included: malnourishment (37%),

injuries (37%), maternal abandonment pre-weaning (15%), lethargy (12%), and parasite infections (8%). Other evidence suggests that the most common presenting condition is trauma: in a review of 205 seals admitted between 2005-2011, Silpa et al. (2015) found that 81.95% of pups also exhibited at least one form of trauma: puncture trauma (68.78%), abrasions (36.59%), orthopaedic disorders (11.7%), and netting injuries (3.4%). These trauma injuries can further lead to other health conditions developing including malnourishment and hypothermia.

Weight-gain is crucial for an individual's health, but it can also be indicative of success upon release (survival of an individual once released and not returned to rehabilitation). In the wild, grey seal pups gain weight rapidly (Figure 1B), but individuals within rehabilitation centres and captivity grow and gain weight at a lower rate (Figure 1C)). O'Hara (2019) found that grey seals pups in rehabilitation gained weight 7.7% of the mean daily weight that individuals in the wild were gaining. However, weight-gain is not always easily attainable within rehabilitation centres, as, for example, replicating the milk for pre-weaning pups is difficult (MacRae et al., 2011). A common feeding method used for seals is gavage feeding (an immediate feeding method, involving restraint as a tube is forced into the oesophagus). However, this feeding method can trigger the stress response which halts the digestive process, further slowing the weight gain of individuals (MacRae et al., 2011). This has been evidenced in harbour seals; MacRae et al. (2011) found that individuals fed on milk-replacers had higher weight gain and a lower mortality rate compared with a fish formula (a common feeding practice within rehabilitation centres due to cost of milk replacement). Rehabilitation centres have tried to replicate the milk with some success (MacRae et al., 2011). Typical hand-rearing diets include artificial milk-replacers and diets based on macerated fish that is fed via gavage, but often weight gain is slow. It is also important that individuals increase in weight

rapidly to reduce the time spent in the centres and further risk of habituation to humans (Cope et al., 2022). There is evidence that habituation to humans happens in a wide variety of taxa in rehabilitation centres (Molinari-Jobin et al., 2024). Pups at postweaning are then fed a different diet comprising of deceased whole herring. This is to both replicate the natural progression of diet, and some aspects of behaviour such as food handling, of seal pups in the wild, but also herring is an easily accessible and costeffective product.

Stressors associated with weight change in experimental research are often similar to management practices within rehabilitation centres (e.g., handling (Harris et al., 1998), transportation (Kannen et al., 2000; Schuamann et al., 2014), changes in social grouping (McLaren et al., 2023), and disturbance (Jayakody et al., 2008)). Recent evidence has shown, for example, that handling of wood mice (Apodemus sylvaticus) in a captive breeding context causes weight loss, with intensive handling having greater weight loss (McLaren et al., 2023). Evidence has also shown that changes in social grouping can cause stress (indicated through physiological and behavioural indices) within social species such as bottlenose dolphins (*Tursiops truncatus*) and can cause weight loss in captive individuals (Waples and Gales, 2002). Environmental stressors are also known to obstruct weight gain through stress-induced susceptibility to disease and can additionally impact their behaviour (Stead et al., 2021). Stress can alter feeding behaviours which can further cause weight loss (Maniam and Morris, 2012). In captivity, when an observer is present, as a potential source of disturbance, behaviours are known to change. A UK study saw resting and clustering behaviours in captive breeding Edwards' pheasants significantly increase when an observer was present but feeding and locomotion behaviours significantly decreased (Hoy and Brereton, 2022). Human disturbance in captive management of wild species is also associated with increased

wariness (Merrick and Koprowski, 2017), for example increased vigilance in wild red deer (*Cervus elaphus*) (Jayakody et al., 2008).

All rehabilitation centres have a variety of pools that the pup passes through once they meet a set criteria (which can vary slightly from centre to centre, this is illustrated below for Seal Rescue Ireland (Figure 2)) that can include a certain weight, completion of treatment or loss of disease signs, recovered from injury, and are able to feed on their own (based on publicly available records from Cornish Seal Sanctuary, 2024). There are optimal paths based on design features, but in practice, management decisions are affected by the capacity of the centre or by disease quarantine. The design of pools are intended to provide opportunities for development of vital skills (e.g., swimming, diving, social skills) and physiological attributes (e.g., muscle mass through exercise), that may support survival in the wild. Rehabilitation centres have a goal release weight which replicates their wild counterparts (typical weight of weaned wild pups is 35 – 40kgs in grey seals) (Based upon publicly available data from Cornish Seal Sanctuary, 2020). Once they have reached this weight, pups are then released back into the wild.



Figure 2: Seal rehabilitation process example; Seal Rescue Ireland (S.R.I.), Courtown, Wexford, Ireland. This a sample of the pupping season 2022 - 2023 (a total of 38 individuals were released, recorded between January and March 2023). When a seal pup is admitted to the centre, it follows a pathway. First, the seal goes to the Intensive Care Unit (ICU) or Hospital, to receive immediate treatment and observation. From here, they go into a recovery area, also known as kennels at the S.R.I. The individual is then moved into one of the three pools (pathways blue, purple, and orange) dependent on management factors such as social structure, weight, and health. Within each of the different stages the number in brackets represents the number of individuals moving between each section in the 2022 - 2023 pupping season at the S.R.I. On the odd occasion, individuals can sometimes return to the kennels due to, e.g., the development of an injury (N=1 in 2022 - 2023 pupping season sample (green pathway)). They progress through the system of pool before reaching the final two: Physio and Pre-Release Pool. Once they have reached the target weight of 35 - 40kgs, individuals are then released (usually with a few individuals from the final social group).

1.4: This Study:

Common rehabilitation practices (e.g., change in environment (Stead et al., 2021), changes in social grouping (Waples and Gales, 2002), and disturbance (Jayakody et al., 2008; Rudin et al., 2018; Hoy and Brereton, 2022)), are known to cause stress to individuals in other contexts. Yet, within a rehabilitation context, the impacts of these stressors on grey seal pup weight gain and survival are less known. These practices are expected to have certain behavioural effects that should also be measurable. Therefore, it is important to measure factors such as these and how they impact growth, behaviour, and success of individuals, to better inform management practices.

There are factors outside of the rehabilitation centres control such as the admittance reason, sex, intake weight, and pupping year, that have already been evidenced to impact survivorship (as seen by Zatrak et al., 2022). This study aims to see if there are management practices inside the rehabilitation centres' control, such as movement between pools (e.g., number of pool changes and the pool characteristics) and natural disturbances (e.g., presence of wildlife (cause for sudden disturbance (Gaspari, 1994)), that could further be impacting weight gain, mortality, and the behaviour of grey seal pups alongside those key factors previously identified within a rehabilitation setting over a six-year period. The objectives of this study are as follows:

- To investigate the effect of known management practices on weight gain (N = 292 grey seals) on multiple pupping seasons and one case study year using a linear regression analysis and mortality on multiple pupping seasons (N = 289 grey seals) using a generalised logistic regression over a six-year period.
- To Investigate the effect of unknown management practices (i.e., pool movement) between pools on weight and mortality of individuals on one case

study pupping season through a repeated measures regression analysis of <70 individuals capturing effects of the pool changes. I hypothesise that given associations between changes in context and stress, I predicted that the higher the number of pool movements, the great the effect on weight gain and mortality.

 To investigate the impact of specific management practices on behaviour (specifically rest, "head elevated", and movement) between each pool through a regression analysis. Repeatability, as described by Dingemanse et al. (2010), was also used to account for repeated measures on individuals to look at pool effects. I hypothesise that grey seal behaviour could be impacted by management practices currently in place, such as water level and pool movement. I further hypothesise that these behaviours are expected to indicate either stress or disturbance hence an effect on growth and mortality.

#### 2: Methodology:

Data was collected from a rehabilitation centre in Ireland; Seal Rescue Ireland (S.R.I.), Courtown, Wexford, Ireland. The S.R.I. have a rehabilitation system involving an ICU unit, kennels, and four separate pools. These pool systems are designed to allow for the development of physical attributes and for natural competition between individuals, preparing the seal pups for release. The S.R.I provided weight and mortality data in a raw format (.csv files), exported straight from their Salesforce Lite database. Individual data collection from the raw format to understand management effects on weight (Seal pupping years 2018 ± 2024, N weight measurements = 3433, N individuals = 292) and mortality (Seal pupping years 2018 ± 2024, N mortality measurements = 289 (26 mortalities)). Due to a malfunction on the S.R.I.'s Salesforce Lite software, data had been lost for pool movements pre-2022. To understand management effects, i.e., pool, on weight, a subset of individual data was collected from the raw format files for seal pupping year 2022-2023 only (Seal pupping year 2022 – 2023, N weight measurements = 368, N individuals = 67). Individual behaviour data was collected on a subset of individuals (pupping year 2022 – 2023, N behavioural samples = 413, N individuals = 31), through Reolink Go PT Plus cameras (Reolink, China) placed by each of the four pools, to understand pool effects on behaviour.

2.1: Rehabilitation Centre:

The S.R.I. site is made up of a visitor centre, an intensive care unit (ICU), kennels, and four separate pools: Nursery, Rock, Physio, and Pre-Release (Figure 4 and Table 2). The water is from a local freshwater source and is UV filtered as pools are refilled. Water drains from each of the pools allowing for cleaning of the pools base. The Pre-Release pool has a small section within the pool area where there is still accessible water during

drainage. The number of grey seals within the centre changes continually between August and March with new intakes and releases.



Figure 3: Aerial view of the Seal Rescue Ireland site, Courtown, Wexford, Ireland (Duncan Kenny Drone photo). The site is located near a leisure centre, a publicly accessible forest trail, and a housing estate. Locations of Reolink Go PT Plus (Reolink, China) cameras are indicated. See Table 2 for specific pool details.

Table 2: Attributes for four pools at Seal Rescue Ireland, Courtown, Wexford, Ireland.

Pool	Pool Attributes	Pool Measurements
Nursery Pool	Smaller and shallower than other pools. Primary use is for seals coming out of ICU/ Kennels.	Length: 10.5m. Width largest point: 7m. Width smallest point: 4m. Pool depth (deepest point): 1m. Pool volume estimate: 38 – 40000L.
Rock Pool	Larger and deeper than Nursery Pool (suited for pups further along the process).	Length: 17.5m. Width largest point: 8.5m. Width smallest point: 6m. Pool depth (deepest point): 1.5m. Pool volume estimate: 115000L.
Physio Pool	Larger and deeper than Nursery Pool (suited for pups further along the process). Section within pool that requires seals to climb over it; intended to	Length: 15m. Width largest point: 8m. Width smallest point: 6m. Pool depth (deepest point): 1.5m. Pool volume estimate: 95000L.

	develop skills and essential physical attributes.	
Pre-Release Pool	Largest pool and final stage of rehabilitation. Pool is enclosed with tall fencing designed to minimise human interaction for the final few weeks before release. Small section within pool area where water is still accessible during drainage.	Length: 20m. Width largest point: 11.5m. Width smallest point: 6m. Pool depth (deepest point): 1.5m. Pool volume estimate: 155000L.

2.2: Weight and Mortality:

The following information for 466 seals from 2012 to 2024 was collated from Salesforce Lite, via manual collation from database and notes provided by the S.R.I.: field identification number (Field ID), sex (male/female), age class (1-3 days, 4-7 days, 7-14 days, 2-4 weeks, 4-6 weeks, 4-8 weeks, 2-4 months, 4-6 months, 4-8 months, 6-12 months, unknown), intake weight (kgs), actual weights (kgs), date of measure, admission reason (emaciated, underweight, illness, injured, dehydrated, orphaned, premature, entanglement, bad location, harassed, and unknown (see Supplementary 1)), location (ICU, Kennels, Nursery Pool, Rock Pool, Physio Pool, and Pre-Release Pool), date moved, mortality (released/deceased), decease date, cause of death if known, pupping year, and additional comments.

Individuals were excluded from the analysis using the following exclusion criteria: minimum of two individuals per year (N = 3), minimum of four recorded weights (N = 139), not currently in rehabilitation (i.e. outcome unknown) (N = 26), not have an unknown age classes (N = 4) and are not in the 6–12-month age class (i.e., unknown whether in the linear growth phase (approx. 200 – 400 days (Figure 1)) (N = 2). Two individuals were re-admitted to SRI after release, for which only data from the first admittance was included. Given the inconsistency in reporting and increasing uncertainty with increasing age class, age class was conservatively modified into dependent (still reliant on the mother (0 - 4 weeks), N individuals = 55, range: 9.2 – 33.4kgs, average weight: 15.71kgs) and non-dependent (not reliant on the mother (4+ weeks), N individuals = 239, range: 9.4 – 37.45kgs, average weight: 16.86kgs).

The duration of time spent in the rehabilitation centre from initial intake was calculated for each weight (Date of measurement – Intake date). Weight change between days was calculated per individual by (mean: 0.25kgs, std. dev: 0.38):

# $\frac{Wd}{ND}$

Where *Wd* = Weight difference between each weight measurement (mean: 1.76kgs, std. dev: 2.18) and *ND* = Number of days between each weight measurements (mean: 6.27 days, std. dev: 3.07). Weight difference across days will be referred to as weight gain herein.

Due to a malfunction on the S.R.I.'s Salesforce Lite software, data had been lost for pool movements pre-2022. To understand management effects, i.e., pool, on weight, a subset of individual data was collected from the raw format files for seal pupping year 2022-2023 only. Therefore, final sample sizes for weight were N individuals = 292, N weight measurements = 3433 for the multiple pupping seasons (2018 – 2024), and N individuals = 67, N weight measurements = 378 for the case study pupping season (2022 – 2023).

Mortality (0 = alive, 1 = deceased) during rehabilitation was censored to <40 days in rehabilitation (threshold set at 40 days as average death occurred at 40 days). Weight change for analysis of predictors of mortality (mean: 8.92kgs, std. dev: 4.8) was calculated by actual weight (kgs) at end of 40-day period (or closest final weight) – intake weight (kgs). Total duration of rehabilitation was noted for all individuals within this analysis (numbers of days between intake date and release/decease date). Individuals

with an accidental or catastrophic death were removed from the analysis (N seals = 3). Due to the aforementioned issue on the S.R.I's software, data had been lost for pool movements pre-2022. To understand management effects, i.e., pool, on mortality, a subset of individual data was collected from the raw format files for seal pupping year 2022-2023 only. However, due to only seven mortalities within this case study year, and only one individual being moved into the pool systems, this data was not analysed. Final sample sizes for mortality were therefore N measurements = 289 for years 2018 – 2024.

#### 2.3 Behaviour:

Behavioural data was collected on 39 seals on eight days between 14<sup>th</sup> February 2023 and 16<sup>th</sup> March 2023. All seal pups were moved only according to S.R.I. practice including for medical treatment, weighing, changing pools, or release when they reached the desired weight of at least 35kgs. The behavioural data was collected using a Reolink Go PT Plus camera (Reolink, China) stationed at each pools (Figure 4, see Supplementary materials 2 for further details). The cameras recorded onto 32GB SD cards. Cameras recorded on a 30 second interval timelapse between 00:00:00 and 23:59:59 on a threeday rotation, recording through infrared at night.

Due to the extensive data (n = 32873 snapshots) and camera malfunctions/faults, behavioural data was only collected between 12:00 and 16:00 at 5-minute intervals during the day (collected by myself and a second observer, n = 768; henceforth 'day data') and in three-time categories during the night (collected by myself): 00:00 - 01:00 (n = 2400), 02:00 - 04:00 (n = 720), 05:00 - 07:00 (n = 1440; henceforth 'night data'). Night and day data were recorded separately to investigate some management practices (such as pool cleans, staff feeds, weighing, medical treatment) that only occur during the day. A combined dataset was compiled using both night and day data (day: n

= 2102, night: n = 910; N = 31; henceforth 'combined data'). Individual ID was collected for 95% of individuals within images using an ID guide that I created from photos collated by the S.R.I. upon instruction. Identification was based on pelage markings, head shape, visible injures, and colouration caused by treatments (Beaumont and Goold, 2007; In the UK, these are public available records from Dorset Wildlife Trust, 2023). All unidentifiable individuals were removed from the analysis.

Four non-mutually exclusive behaviours were collected for both day and night data: "head elevation", "movement", "resting", and "aggression" (Table 3). These four behaviours were selected through a pilot study (behaviours were added as they were observed in 240h timelapse in one pool to construct an expanded ethogram), to identify common behaviours that could be objectively classified (unlike e.g., distance of proximity to another seal)), and was informed by some studies on wild grey seals (Chilvers et al., 1999; Twiss and Franklin, 2010). These four behaviours are biologically important to link weight gain and stress. Head elevated was included with the intention of capturing alertness or vigilance or may equally capture neutral or positive interactions, e.g., social or feeding opportunities.

Table 3: Behavioural ethogram created for data collection on time-lapse images of seals.

Behaviour	Definition
Head elevated	Head is up from head down position. Individual's head can be slightly raised from head down position or raise significantly with neck extended.
Movement	Any movement from original position that is active and require energy expenditure. This can include one of the following: swimming (both at surface and underwater), looking underwater, galumphing (moving in a humped motion forward), climbing, scratching.
	Can coincide with head elevated behaviour.
Rest	Head down, eyes can be open or closed. Can be in the water or on land. Includes bottling and logging behaviour seen within the footage (Seals are resting but are able to surface and go back

Aggression	underwater without waking, eyes remained closed, head can be in a particular direction. Logging is horizontal, bottling is vertical).
	Can coincide with head elevated behaviour (8.5% of observations) but can never be moving.
	Individuals within a close distance, teeth can be visible, neck forward, lips raised. Two individuals face on, nose to nose (wit muzzles touching). Can be seal directed, human (staff) directed or wildlife (e.g., heron) directed.
	Can coincide with head elevated and movement behaviour.

Contextual variables were recorded from the snapshots to capture disturbances, social

factors and environmental factors that may affect behaviour on the specific individual.

During pilot data collection, in comparison to other wildlife such as seabirds, only herons

were observed to alter seal behaviour as seal pups were observed chasing herons. This

could be causing energy expenditure and therefore impacting weight gain, or

alternatively, could be a source of enrichment for the seals, so was included within the

data collection.

Table 4: Contextual variables considered to affect the behaviour of individuals across pools.

Environmental Factor	Description:
In Proximity	0 = not in proximity. 1 = within a seal body length of another individual.
Water Level	0 = little water to no water, 1 = Shoulder depth, 2 = swimmable/full
Intensity of people	0: No people in the pool.
	1 (Low Intensity): People in enclosure, seals not reactive. Staff very briefly in pool (less than five minutes recorded via timestamp in snapshots).
	<ul> <li>2 (Medium intensity): People in enclosure, seals are reactive, but staff are not interacting with seals. Staff are in pool for short period of time or an extended period of time but no disturbance to seals, e.g. pool clean (over 10 minutes recorded via timestamp in snapshots).</li> <li>3 (High intensity): People in enclosure. Staff are interacting with seals (e.g., weighing, medication administration).</li> </ul>

Staff Feed	Staff are throwing fish into pool. This occurs between two		
	and three times at varied times throughout the day		
	between 07:00 and 19:00 during the observation periods.		
Weather	Rain versus dry, visible in image.		
Heron	0: Absent from the pool. 1: Present in the pool.		

2.4: Data Analyses:

Data was analysed using R (version 4.3.1, (R Core Team, 2023)) via the software R Studio, version 2023.12.0+369 (R Core Team, 2023). All visual outputs have been produced through "sjPlot" (Lüdecke, 2023) and "tidyverse" (Wickham et al., 2019).

2.4.1: Weight Change:

Weight change was analysed using linear mixed effects models (LMER's) for six consecutive over-winter pupping seasons between 2018-2024 (N individuals = 292, n weight measurements = 3433), using the package "Ime4" (Bates et al., 2015). Weight change between consecutive measurements (kgs) was the dependent variable instead of actual weight per measurement to increase sensitivity for capturing effects of current context. Overall weight and age might both affect capacity to change weight, were controlled for by including actual weight as an independent variable. Further fixed independent variables were duration of rehabilitation to date of weighing (days), sex (male/female), age (Dependent or Non-Dependent), and admittance reasons: underweight (yes/no), illness (yes/no), injured (yes/no), and seal pupping season (2018 – 2019 (n = 192, N individuals = 10), 2019 – 2020 (n = 877, N individuals = 73), 2020 – 2021 (n = 924, N individuals = 59), 2021 – 2022 (n = 619, N individuals = 48), 2022 – 2023 (n = 571, N individuals = 72), 2023 – 2024 (n = 70, N individuals = 10)). Individual ID (Field ID) was controlled for as a random effect in the analysis.

As pool data was only available for 2022 – 2023 seal pupping season, a separate analysis was done for weight. The same fixed independent variables and random effects were

considered with the addition of pool ID at measurements, and if they changed pool between measurements (yes/no). ICU and Kennel were not included in the pool movement as the focus was specifically on the pools. Changes included movement from kennel to pool to include the first weight.

#### 2.4.2: Mortality

Mortality (0 = "alive", 1 = "deceased") was analysed using a generalised logistic regression model (GLM, family = binomial) for the same six consecutive over-winter pupping seasons (2018-2024) (N = 292), through package "lme4" (Bates et al., 2015). There were 23 mortalities (including individuals euthanised (n = 5 seals)) and 268 released. Fixed independent variables were age (Dependent or Non-Dependent), sex (male/female), intake weight (kgs), weight change at 40 days, standard deviation of weight change (0 – 40 days) (as a measure of consistency in expected weight gain), duration of rehabilitation (total time), admission reasons: underweight (yes/no), injured (yes/no), and illness (yes/no). Seal pupping season was removed due to uneven mortalities and limited data within some years to explore mortality (2018 – 2019, 10 data points, 0 mortalities).

#### 2.4.3: Behaviour

Behaviour was analysed using generalised linear mixed effects models (GLMM's) on the combined data (n observations = 415, N individuals = 31), using package "Ime4" (Bates et al., 2015), for three behaviours (head elevated, movement, and rest). Aggression was not analysed due to rarity of behaviour (combined night and day: n observations = 28). Fixed independents variables were pool ID, time of day (day v. night), heron (absent v present), and in proximity of another seal. The reference levels were Pre-Release Pool, day, heron present, and not in proximity of another seal. Individual ID was included as a
random effect to account for repeated measures of individuals, through comparing AIC numbers (see Supplementary 4 for further details). Confounding effects were tested for using package "car" (Fox and Weisberg, 2019). Observations of people and rain were confounded with day/night, so are not included in these models, but analyses for day and night data are presented separately in Supplementary materials 4.1 - 4.2.

The focus of the analyses was to explore pool effects on behaviour. To determine the extent to which behavioural variation observed was driven by individuals', consistency repeatability estimates were used following Nakagawa and Schielzeth, 2010, using package "rptR" (Stoffel et al., 2017). Estimated adjusted repeatability (an estimate adjusting for confounding effects) was used as there were confounding effects within the GLMM's (a variable distorting the observed relationship between the independent and dependent variables). To be classed as repeatable, the estimates needed to be higher than 0.2 (Harper 1994).

### 2.5: Ethics

This study was approved by Seal Rescue Ireland and the University of Plymouth's School of Biological and Marine Sciences Animal Ethical Review Committee (2023). The study did not require any changes to normal management practices of the Seal Rescue Centre staff. As the cameras were facing away from the public, human ethical review was limited to notification of staff regarding timelapse recording, appropriate storage of footage, and protocols to ensure personally identifiable information was omitted from files shared by S.R.I. for analysis.

### 3.1: Weight

3.1.1: Variables Influencing Weight Change for Six Consecutive Pupping Seasons.

The actual weight (kgs) of an individual is positively correlated with weight gain between days (Table 5, Appendix 3, Figure A3A)). The duration of rehabilitation is negatively correlated with weight gain between days (i.e., the loner the duration of rehabilitation, the lower the weight gain between days) (Appendix 3, Figure A3B). Compared to individuals not underweight upon admittance, underweight individuals had greater weight gain between days (Appendix 3, Figure A3C)). Compared to individuals without an illness, individuals admitted with an illness had lower weight gain between days (Appendix 3, Figure A3C)). Compared to individuals without an illness, individuals admitted with an illness had lower weight gain between days (Appendix 3, Figure A3C)). Compared to the 2018 – 2019 pupping season, 2019 – 2020, 2020 – 2021, and 2022 – 2023 pupping seasons all had greater weight gain between days.

Table 5: Variables influencing weight gain over six consecutive pupping seasons (N = 292 seals, n = 3433 weight measurements). Reference level: dependent age (0 – 4 weeks); female, not underweight, no injury, no illness, 2018 – 2019 Seal Pupping Season. Random effects was individual ID (Field ID). Random effects: residual variance ( $\sigma^2$ ) = 0.13, Between subject variance ( $\tau_{00}$ ) = 0.00, Marginal R<sup>2</sup> = 0.097 Conditional R<sup>2</sup> = NA.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	-0.27	-0.37 – -0.17	<0.001
Actual Weight (Kgs)	0.02	0.02 - 0.02	<0.001
Age: Non-dependent	-0.02	-0.05 - 0.02	0.287
Male	-0.01	-0.04 - 0.01	0.251
Duration of Rehabilitation	-0.00	-0.000.00	<0.001
Underweight	0.10	0.07 - 0.14	<0.001
Injured	0.01	-0.02 - 0.04	0.413
Illness	-0.04	-0.070.01	0.016
2019 – 2020 Seal Pupping Season	0.07	0.01 - 0.13	0.014
2020 – 2021 Seal Pupping Season	0.09	0.03 - 0.15	0.002
2021 – 2022 Seal Pupping Season	0.06	-0.00 - 0.12	0.059
2022 – 2023 Seal Pupping Season	0.12	0.06 - 0.18	<0.001
2023 – 2024 Seal Pupping Season	0.10	-0.00 - 0.20	0.053

3.1.2: Variables Influencing Weight Change for Case Study Pupping Season.

The actual weight (kgs) of an individual is positively correlated with weight gain between days (Table 6, Appendix 4, Figure A4A). The duration of rehabilitation is negatively correlated with weight gain between days (i.e., the longer the duration of rehabilitation, the lower the weight gain between days) (Appendix 4, Figure A4B). Compared to individuals without an injury, individuals admitted with an injury had higher weight gain between days (Appendix 4, Figure A4C). Although marginally significant, compared to Pre-Release Pool, individuals within Rock Pool gained more weight between days (Appendix 4. Figure A4D). There was no effect of individuals on weight gain between days.

Table 6: Variables influencing weight change for case study pupping season (N = 67 seals, n = 378 weight measurements). Reference level: dependent age (0 – 4 weeks), female, Pre-Release Pool, not changed pool, not underweight, no illness, and no injury. Random effects: individual ID (Field ID), residual variance ( $\sigma^2$ ) = 0.15, Between subject variance ( $\tau_{00}$ ) = 0.00, Marginal R<sup>2</sup> = 0.124, Conditional R<sup>2</sup> = NA.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	-0.36	-0.67 – -0.05	0.025
Actual Weight (Kgs)	0.03	0.02 - 0.04	<0.001
Age: Non-Dependent	-0.06	-0.17 – 0.05	0.321
Male	-0.02	-0.10 - 0.06	0.628
Duration of Rehabilitation	-0.00	-0.010.00	<0.001
Nursery Pool	0.11	-0.07 – 0.30	0.227
Rock Pool	0.12	-0.01 - 0.24	0.065
Physio Pool	0.01	-0.09 - 0.10	0.901
Changed Pool (Yes)	0.02	-0.07 - 0.12	0.634
Underweight	0.09	-0.08 – 0.26	0.308
Illness	-0.07	-0.24 - 0.11	0.445
Injured	0.09	0.01 - 0.17	0.035

3.2: Mortality:

3.2.1: Variables Influencing Mortality for Six Consecutive Pupping Seasons.

There were no significant variables influencing mortality within the six consecutive pupping seasons (Table 7). Although marginally significant, compared to individuals who

are not underweight upon admittance, underweight individuals had a higher likelihood

of mortality. Although marginally significant, intake weight is positively correlated with

mortality. There was no effect of individuals on mortality.

Table 7: Variables influencing mortality (0 = alive, 1 = death) for six consecutive pupping seasons (N = 289 measurements). Reference level: female, age dependent (0 – 4 weeks), not underweight, not injured, and no illness.  $R^2$  Tjur = 0.085.

Predictors	Estimates	<b>Confidence Interval</b>	p-value
(Intercept)	11.52	0.43 - 376.04	0.154
Male	1.81	0.68 – 5.39	0.251
Age: Non-Dependent	0.52	0.19 - 1.54	0.218
Intake Weight (Kgs)	0.88	0.77 – 0.99	0.052
Total Duration of Rehabilitation	0.99	0.97 – 1.00	0.138
Weight Change at 40 days	0.98	0.88 - 1.09	0.664
St.d. Deviation of Weight Change	0.92	0.77 – 1.07	0.326
Underweight	0.33	0.09 - 1.23	0.088
Injured	0.46	0.17 – 1.18	0.118
Illness	1.72	0.53 – 5.14	0.345

3.3: Behaviour

There were significant differences in behaviour across pools. Head elevated also had a higher likelihood of being observing in Rock Pool when compared to Pre-Release pool (Figure 5a). Compared to Pre-Release Pool, there was a higher likelihood of observing movement in Rock Pool and Physio Pool (Figure 5b). Compared to Pre-Release Pool, there was a lower likelihood of observing rest in Nursery Pool, Rock Pool, and Physio Pool, with Rock Pool being the least likely to observe resting behaviour (Figure 5c). When in close proximity to another seal, there was a lower likelihood of being observed in movement behaviour. When in close proximity to another seal, there was a lower likelihood of being observed when herons were present. Compared to during the day, head elevated, and movement behaviour were less likely to be observed during the night. However, compared to during the day, rest was more likely to be observed during the night. The combined day-night results are presented for greatest utility to management (to allow for a general

overview of changes within behaviour and pool effects) (Supplementary S4.3), but when day and night were analysed separately, time of day and intensity of staff (i.e., medium) were also significant (Supplementary S4.1 and S4.2). All three behaviours scored lower than the threshold for repeatability ( $R_M = 0.02$ ), but movement and rest were statistically significant (see Supplementary 4).



Figure 4: The likelihood of different management practices influencing three grey seal (*Halichoerus grpyus*) behaviours ((a) head elevated, (b) movement, (c) rest) within a rehabilitation centre based in Ireland. Individual ID was factored in as a random effect (N = 31, n = 415). Reference level: Pre-Release Pool, Day, heron absent, and not in proximity of another seal. Blue coloured factors have positive odds of the behaviour occurring and red coloured factors have negative odds of the behaviour occurring. The red line indicates zero and if the factor crosses this line, this factor for this behaviour is more down to chance. P-values are indicated as Asterix, three Asterix is <0.001.

#### 4. Discussion:

Investigating the management practices at S.R.I has identified that factors inside the centres control, particularly pool movement, largely did not influence weight gain between days. Most variables affecting weight gain and mortality were outside of the rehabilitation centres control (e.g., pupping season, intake weight, and admittance reason). There were marginally significant differences in weight gain between pools, however, this may capture some aspect of design or management. Pool type did also influence the behaviour of the grey seal pups. Rock Pool (a large pool with no fencing) was found to have increased likelihood of observing head elevated behaviour and decreased likelihood of observing resting behaviour, when compared with Pre-Release Pool (shielded pool), further suggesting that there is a difference in the seals' perception or experience of environmental factors in this pool.

4.1: Impact of Management Practices:

Pool movements are a regular part of management practices within rescue centres, and movements, including not only how often but also which pools individuals are moved across, and who moves where, are decided upon using criteria including behavioural traits as classified by rehabilitation centre, weight, and the number of individuals within the centre. Across a wide taxonomic range, previous research on known stressors such as transportation (Kannen et al., 2000; Schuamann et al., 2014), handling (Harris et al., 1998), changes in social grouping (Waples and Gales, 2002; McLaren et al., 2023), and disturbance (Jayakody et al., 2008) has been shown to affect weight gain. There was no evidence found within this study that movement of an individual between pools affected weight gain ("Changed Pool"). This is surprising as it is expected that the individuals would lose weight due to potential stress associated with movement. This

may suggest that young seals are robust to the level of disturbance required to manage them effectively.

Another management factor is pool design, including the inclusion of shields around the Pre-Release Pool. I predicted that this would reduce human and visual disturbance that may cause stress, hence increase weight gain. This was not observed, and unexpectedly, within the case study year, one of the unshielded pools, Rock Pool, had marginally higher weight gain between days (p = 0.065, est. 0.12kgs difference) compared with the shielded pool (Pre-Release Pool). This may not exclude stress: increased weight via eating or resting more potentially acts as a buffer to compensate potential weight loss under perceived environmental risks (Berry et al., 2013; Malik and Spencer, 2019). There is also evidence that once a restriction period ends, it can trigger an increase in food consumption and a larger energy store even when energy expenditure remains low (Jacuier et al., 2014).

Behavioural data also suggest that, of the three pools, Rock Pool may have higher disturbance: in this pool seals were less likely to be observed resting behaviour and more likely to be observed showing head elevated behaviour when compared to Pre-Release Pool (shielded pool). If so, the source of the disturbance is unknown, but disturbance generally can cause disruption to rest. Beyond total rest, fragmented sleep also, in humans, has been associated with impaired alertness, memory, mood regulation, changes to brain activity and metabolism (Short and Banks, 2013). Within harbour seals, there is evidence that disturbance from pedestrians, within 50m distance, impacts resting behaviour when hauled-out (the act of coming onto a solid surface (e.g., land or ice)) on land, as when disturbed, they haul-in (the act of returning into the water after hauling out; can be a very rapid response to fear) reducing the time spent resting and

increasing the energetic output (Osinga et al., 2012). However, importantly, an alternative explanation for behavioural differences is that the behaviour is neutral, and that there were other differences (stressful or beneficial) not measured in this pool that may have affected weight. This is certainly possible as the site sits next to a leisure park and a public woodland which could be causing extra stimulus. Furthermore, these behaviours could be positive, and these differences emerge only when there is a surplus of energy to maintain their weight whilst also maintaining their natural behaviour. Consequently, it is hard to use the behavioural data to interpret the weight gain between days within this pool, so further research is necessary.

Head elevated behaviour may (or may not) be evidence disturbance. Within this study, there is evidence that head elevated could be an indicator of disturbance, as the likelihood of observing head elevated behaviour increased when herons were present. While Heron presence was confounded with feeding time during the day that may otherwise explain this behaviour, this association was also observed at night, suggesting a direct association. Interestingly, within this study there was an unusually high presence of herons observed across the pools (20% of observations), with seals observed chasing herons within the pilot data collection. One explanation for the increased likelihood could be that the herons cause a sudden disturbance to the seals. Sudden disturbances have been shown to increase vigilance behaviour in wild grey seals (Gaspari, 1994). If this were the case, this could increase energy expenditure and sleep deprivation. However, the results within this study show that in the pool with the highest instance of head elevation, there was marginally higher weight gain. One other potentially interesting explanation could be that the herons are providing a form of enrichment to the seals. They could be providing naturalistic enrichment which could be more valuable to the individuals than other forms of enrichment (Ruppert, 2017; Eagan, 2018). Further

research is needed to determine the context of relationship between the herons and the seals and to interpret head elevation as a behavioural indicator.

Group size is another management factor. Although group size per se is confounded with pool, so cannot be explored, the effects of proximity can provide an insight into this. The case study pupping season observed increased likelihood of resting behaviour whilst in proximity of another seal. One explanation for this is that grey seals are known to be social at haul-out sites during the mating/pupping season and moulting season as they congregate in large groups on land. This finding was consistent with Twiss et al., (2022) who also found that grey seals show pro-social behaviours. The increased resting behaviour whilst in proximity of another seal is consistent with behaviour already observed in other seal species (Chilvers et al., 1999), although there is currently no previous evidence of this in grey seals. In harbour seals, evidence has shown that when in groups there is increased allocation of time spent resting in larger and tighter groups (Krieber and Barrette, 1984), which could potentially be reducing stress within the grey seals observed in this study. Additional evidence from Twiss et al., (2022) has shown that grey seal pups can recognise individuals they have previously encountered, and reduce the level of costly interactions, such as aggression, with these familiars. Therefore, it is necessary for the rehabilitation centre to consider the groupings and/or regroupings as best as possible to avoid certain interactions between individuals, but also the size of the group (this will vary between centres and the number of pups admitted in the year).

When exploring management factors, variation of the path of individual seals through the pools gave scope to separate the effects of e.g., age and weight, that increased through time, from pool characteristics. Random allocation of individuals across the

pools required to fully separate these variables would necessitate significant and potentially risky changes to practice that are informed by knowledge on suitability of e.g. behavioural and weight combinations, as well as space and resource management more generally. Therefore, it is important to note that some selection of more similar individuals to go together could be introducing biases to this study. However, importantly, recommendations from the study are specific hence relevant to current management practice. To address this further in the future, data collection on the management decision on where individuals go would help address potential biases.

Additionally, I could not summarise the data in a reasonable way per individual in order to compare behaviour and weight change at the individual-level. As the behavioural data was collected through point sampling over shorter time periods (snapshot every 30s), and the weight data collected over longer time periods (once a week), it was not possible to derive one summary measure from varied numbers of point samples across the different conditions. As such, it remains unknown if the pool with the highest indicator of disturbances or the number of moves did cause any weight changes.

A limitation to the behavioural data is that due to camera malfunctions and use of point sampling, it is estimated only 5% of the recorded day footage was collected. Data was not collected between 07:00 and 12:00, so there could be events (including pool cleans, weighing, individuals added into/removed from the group) occurring during this time that could be influencing the behaviours later on, e.g., new individual added into or removed from the group, changing the social dynamic and grouping. However, as data was collected across the day and night, it removes the limitation of events occurring during the night impacting those of the day and *vice versa*. There were expected variations between the night and day data due to more management practices occurring

during the day (e.g., pool cleans, weighing, medication), and these effects emerged (e.g., more active during the day, more at rest at night), further validating the approach within this study at capturing ecologically relevant effects. Aggression was not analysed for two reasons; 1) point sampling was not appropriate for this behaviour due to the rare, shortlived behaviours, and 2) the time-lapse camera took a snapshot every 30s which is inadequate for the unpredictability and speed of this behaviour. Aggression would require focal follows and, if to be recorded via time-lapse cameras, would require more frequent timing, e.g., every 10s.

4.2: Impact of Factors Outside the Rehabilitation Centre's Control:

There are many factors outside the control of the centre, including the reason for admittance (e.g., disease, injury, and being underweight), age, sex, and differing environmental factors per pupping season. The main management objective within rehabilitation centres is to maximise weight gain but also to minimise the time in captivity, reducing the risk of habituation, and to minimise the resources required per seal. This study found that disease status was associated with lower weight gain between days, and this was greater in seals admitted for being underweight (and in the case study year, those admitted for injury). Another important finding of this study was that duration of rehabilitation was associated with lower weight gain between days, i.e., the longer the seals stayed in rehabilitation, the less weight they gained between days. Within marine mammals, there has been a lack of research to confirm if the duration of rehabilitation has been related to survival (Wells et al., 2012). This study appears to be the first to find evidence that the total duration of rehabilitation had no effect on mortality. This suggests that although the longer rehabilitation time sees reduced weight gain, it does not cause mortality in the long-term. The slowing weight gain could be explained by a number of factors including plateauing growth, stress-induced weight gain early in admission or loss with prolonged admission, or delayed effects of the initial condition upon admittance.

This study confirms that being underweight (84%), having an illness (a broad category that included parasites (e.g., lungworm or roundworm) and suspected viruses (e.g., seal pox or phocine distemper)) (22%), and injury (including entanglement) (52%), are common admittance reasons for grey seals. Furthermore, 73% of mortalities within this study were admitted as underweight. This finding is consistent with Zatrak et al. (2020) where 35% of seals were underweight. This high admittance has been previously attributed to interruptions in suckling regime of mother and juveniles (Anderson et al., 1979) as well as post-weaning fasting caused by the learning period (Zatrak et al., 2020). However, Zatrak et al., (2020) sample size was 15.75 times larger than this study (N individuals = 292), and over a 32-year period (1996 – 2020). This larger sample allows for better evaluation of admittance trends. However, since 2020, within this study alone, there has been an increase of individuals admitted with underweight as the cause (180 individuals between 2020 and 2024). This is cause for further concern as previous evidence has shown that heavier individuals had higher survival probabilities (Hall et al., 2002, 2008; Zartak et al., 2020).

Interestingly, while weight gain was higher in individuals classed as underweight on admittance, this study also found marginal evidence that these individuals are more likely to die than those who were not. A potential explanation could be that rapid weight gain in individuals who were more severely underweight increases instantaneous mortality risk. This is observed in humans, following increased food intake after a period of starvation, known as "refeeding syndrome" (Brooks and Melnik, 1995). Refeeding

syndrome has also been seen in other mammalian species such as domestic cats (Chan, 2015), so it is possible that it could also be happening within grey seal pups. Higher weight gain may also indirectly indicate individuals with higher admittance-related stress, as glucocorticoids (GC) terminate fasting in pinnipeds (Bennett et al., 2013) However, regulation of weight is complex, and Bennett et al., (2013) evidenced that in an experimental setting, high GC levels can alter the allocation of resources and alter mass loss rate, specifically by increasing in the protein breakdown in fasting grey seal pups.

However, the use of the admittance term "underweight" could be influencing the results of this study. The centre within this study uses the term emaciated for those severely underweight, but as this term is subjective, and some individuals are recorded as both emaciated and underweight, I used the broader, more conservative description. The term 'underweight' also conflates varying degrees of severity in non-emaciated individuals. These differing severities could impact how much weight an individual gains and their mortality risk (see "refeeding syndrome" above). There is evidence, for example, that individuals above the age-predicted weight have increased survival odds (Zatrak et al., 2022). So, it is important to further categorise weight or combine with a skeletal measure both to increase sensitivity of the analysis and to help identify thresholds in risk.

Further evidence from this study revealed that individuals with the admittance reason illness were found to have lower weight gain. This potentially suggests that closer monitoring of these specific individuals is needed. Only within one pupping season (2022 – 2023), seals admitted with an injury had a higher weight gain between days than those without an injury. However, some of the variation within this year was explained by

other factors, such as pool changes and age. Additionally, this study found evidence that the actual weight (kgs) of an individual had a positive correlation with weight gain between days. This adds to the evidence presented by Zatrak et al., (2022); odds of juvenile seal survival increased by 1.05 times per kilogram when over the age predicted weight. There were also significant year effects on weight gain between days, with three years having greater weight gain (2019 - 2020, 2020 – 2021, 2022 – 2023). An explanation for this could be that for these years, the sample sizes were larger, potentially capturing more of the variation within these years.

This study found marginally increased likelihood of mortality for individuals who had relatively high intake weights. One potential explanation is that these are individuals that were thriving until admittance, due to more catastrophic reasons, including injury (Cases within this study: 2) and expression of genetic diseases such as Disseminated Intravascular Coagulation (DIC) (Cases within this study: 3). Another finding was that there was no effect of initial rate of weight gain (in the first 40-day of rehabilitation) on mortality. The average death occurred at 40 days in the focal individuals, but there was a much wider range (minimum: 13 days, maximum: 253 days), potentially meaning that the 40-day threshold excludes some important individuals and is not large enough to capture any changes in weight occurring just before mortality for individuals who stayed for longer.

4.3: Individuals within Rehabilitation:

Individuals are a vital part of conserving species, and it is important to make sure that management practices do not impact individuals (Aitken, 2004). Repeatability measures are a way to measure this, allowing for important management practices and decisions to correctly be put in place. In this study, ID explained a statistically significant

proportion of variation in for most behaviours (except head elevated), but the effect size was small with low repeatability ( $R_M = <0.2$ ). One possible interpretation of this result is that current management practices in place may not therefore have a different effect on different individuals. As all individuals within this study were pups and not adults, it is possible that behavioural traits often reported in adult seals have not yet developed. A 2015 review reported evidence that in juveniles, behavioural repeatability is often lower than adults (Brommer and Class, 2015). Brommer and Class (2015) theorised that although there is an absence of personality variation here, it need not exclude differences that develop between individuals later on. Therefore, it is important to investigate personality traits across the different life stages (pup, juveniles, adolescents, and adults) to understand if it will affect current management in rehabilitation centres.

However, there are some points to note regarding these interpretations. Firstly, as noted above, there are management factors that determine how individuals are allocated to pools (e.g., individual seal behavioural observations from the rehabilitation staff). This may systematically bias the relationship between behaviour and pool ID, making it difficult to separate individual from pool effects. Furthermore, behaviour was not measured under standardised conditions before seals were allocated to pools. This is required to quantify personality variation, as pool characteristics may amplify or suppress the expression of different behavioural traits. Therefore, to truly understand the effect of personality variation within rehabilitation centres, more research is needed. However, this pilot data showing no strong behavioural differences with respect to different management factors is encouraging that the rehabilitation environment might allow all personality types to thrive.

4.4: Recommendations and Further Research:

It would be advisable to monitor the weight of diseased individuals more closely due to the lower weight gain between days. A further recommendation would be to investigate the differing severities of admittance reason "underweight" and how this could be potentially impacting weight gain and mortality. Interestingly, a study by Zatrak et al., (2022), also recommended more consistency across sites in recording admittance reasons by seal rehabilitators to further enable research into admissions and survival probabilities of admitted seals. Additionally, a recommendation from this study to the S.R.I. is to monitor the Rock Pool to try to understand what could be causing the difference in behaviours/weight within this specific pool and if it is caused by a specific disturbance, or alternatively captures seals who are growing well hence have surplus energy to invest in more diverse behaviours. More generally, further research would be beneficial to understand the drivers of movement and head up behaviour to understand whether or not these indicate disturbance. Based on rest data however, it may be beneficial for pool designs to include a fence, creating a visual barrier between seals and humans, removing potential effects of disturbance from the presence of people. A further recommendation based on the literature motivating the study is to review the pool movements and its impact on mortality, as there was insufficient data to investigate this.

Additional research on the interactions between environmental cues and seals to feeding would further help with understanding the relationship between herons and head elevated behaviour, and if this is vigilance behaviour or if it is more specifically related to the presence food. The herons could be a source of natural enrichment, but

it could also be a stressor. Therefore, further research is also needed to truly understand

the relationship between them.

### **References:**

Aitken, G. 2004. A New Approach to Conservation: The Importance of the Individual Through Wildlife Rehabilitation, First edition, MPG Books Ltd, Bodmin, Cornwall.

Anderson, S.S., Baker, J.R., Prime, J.H., Bard, A. 1979. Mortality in Grey Seal Pups: Incidence and Causes. *Journal of Zoology*, 189(3), 407 – 417. https://doi.org/10.1111/j.1469-7998.1979.tb03972.x

Anderson, S.M., Teilmann, J., Dietz, R., Schmidt, N.M., Miller, L.A. 2011. Behavioural Responses of Harbour Seals to Human-Induced Disturbances. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 22(1), 113 – 121. <u>https://doi.org/10.1002/aqc.1244</u>

Armstrong, H.C., Russell., D.J.F., Moss, S.E.W., Pomeroy, P., Bennett, K.A. 2023. Fitness Correlates of Blubber Oxidative Stress and Cellular Defences in Grey Seals (*Halichoerus grypus*): Support for the Life-History-Oxidative Stress Theory from an Animal Model of Simultaneous Lactation and Fasting. *Cell Stress and Chaperones*, 28(5), 551 – 566. <u>https://doi.org/10.1007/s12192-023-01332-1</u>

Atkinson, S., Dieraug, L.A. 2018. Stress and Marine Mammals. In: Gulland, F.M.D., Dierauf, L.A., Whitman, K.L (eds) CRC Handbook of Marine Mammal Medicine, third edition, CRC Press. <u>https://doi.org/10.1201/9781315144931</u>

Avital, A., Ram, E., Maayan, R., Weizman, A., Richter-Levin, G. 2006. Effects of Early-Life Stress on Behaviour and Neurosteroid Levels in the Rat Hypothalamus and Entorhinal Cortext. *Brain Research Bulletin*, 68(6), 419 – 424. <u>https://doi.org/10.1016/j.brainresbull.2005.09.015</u>

Ben-Porath, I., Weinberg, R.A. 2004. When Cells Get Stressed: An Integrative View of Cellular Senescence. *The Journal of Clinical Investigation*, 113(1), 8 – 13. <u>https://doi.org/10.1172/JCI20663</u>

Barnett, J.E.F., Woodley, A.J., Hill, T.J., Turner, L. 2000. Conditions in Grey Seal Pups (*Halichoerus grypus*) Presented for Rehabilitation. *Veterinary Record*, 147(4), 98 – 104. https://doi.org/10.1136/vr.147.4.98

Barnett, J.E.F., Allen, R., Astley, K., Whitehouse, F., Wessels, M.E. 2021. Pathology of Grey Seals (*Halichoerus grypus*) in Southwest England Including Pups in Early Rehabilitation. *Disease of Aquatic Organisms*, 145, 35 – 50. https://doi.org/10.3354/dao03600

Bernier, N.J. 2006. The Corticotrophin-Releasing Factor System as a Mediator of the Appetite-Suppressing Effects of Stress in Fish. *General and Comparative Endocrinology*, 146(1), 45 – 55. <u>https://doi.org/10.1016/j.ygcen.2005.11.016</u>

Berry, M.J., Jaquiery, A.L., Oliver, M.J., Harding, J.E., Bloomfield, F.H. 2013. Antenatal Corticosteroid Exposure at Term Increases Adult Adiposity: An Experimental Study in

Sheep. Acta Obsetetricia et Gynecologica Scandinavica, 92(7), 862 – 865. https://doi.org/10.1111/aogas.12149

Bennett, K.A., Fedak, M.A., Moss, S.E.W., Pomeroy, P.P., Speakman, J.R., Hall., A.J. 2013. The role of glucocorticoids in naturally fasting grey seal (*Halichoerus grypus*) pups: dexamethasone stimulates mass loss and protein utilisation, but not depature from the colony. *Journal of Experimental Biology*, 216(6), 985 – 991. https://doi.org/10.1242/jeb.077438

British Small Animal Veterinary Association (BSAVA). 2017. BSAVA Manual of Wildlife Causalities, second ed. eBook.

Brommer, J.E., Class, B. 2015. The Importance of Genotype-By-Age Interactions for the Development of Repeatable Behavior and Correlated Behaviors Over Lifetime. *Frontiers in Zoology*, 12(S2). <u>https://doi.org/10.1186/1742-994-12-S1-S2</u>

Brooks, M.J., Melnik, G. 1995. The Refeeding Syndrome: An Approach to Understanding its Complications and Preventing its Occurrence. *Pharamacotherapy*, 15(6), 713 – 726. https://doi.org/10.1002/j.1875-9114.1995.tb02887.x

Burroughes, N.D., Dowler, J., Burroughes, G. 2021. Admission and Survival Trends in Hedgehogs Admitted to RSPCA Wildlife Rehabilitation Centres. *Proceedings of the Zoological Society*, 74, 198 – 204.

Bull, J.C., Janes, O.R., Börger, C., Franconi, N., Banga, R., Lock, K., Stringell, T.B. 2021. Climate Causes Shifts in Grey Seal Phenology by Modifying Age Structure. *Proceedings of the Royal Society of Biology: Biological Sciences*, 288(1964). <u>https://doi.org/10.1098/rspb.2021.2284</u>

Chaput, J., Tremblay, A. 2012. Insufficient Sleep as a Contributor to Weight Gain: An Update. *Current Obesity Reports*, 1, 245 – 256.

Chan, D.L. 2015. Refeeding syndrome in small mammals, in: Nutritional Management of Hospitalized Small Animals. John Wiley & Sons Ltd, Chichester, UK, pp. 159 – 164.

Chilvers, R., Colebourne, M., Grant, B., Oliver, R., Lea, S.E.G. 1999. Vigilance in Grey Seals as a Function of Time Since Haul Out. *Journal of the Lundy Field Society*, 50.

Crespi, E.J., Denver, R.J. 2004. Ontogeny of Corticotrophin-Releasing Factor Effects on Locomotion and Foraging in the Western Spadefoot Toad (*Spea hammondii*). *Hormones and Behavior*, 46(4), 399 – 410. <u>https://doi.org/10.1016/j.yhbeh.2004.03.011</u>

Cope, H.R., McArthur, C., Dickman, C.R., Newsome, T.M., Gray, R., Herbert, C.A. 2022. A Systematic Review of Factors Affecting Wildlife Survival During Rehabilitation and Release. *PLoS One*, 17(3), e0265514. <u>https://doi.org/10.1371/journal.pone.0265514</u>

Dickens, M.J., Delehanty, D.J., Romero, L.M. 2009. Stress and Translocation: Alterations in Stress Physiology of Translocated Birds. *Proceedings of the Royal Society B: Biological Sciences*, 276(1664). <u>https://doi.org/10.1098/rspb.2008.1778</u> Dmitriew, C.M. 2011. The Evolution of Growth Trajectories: What Limits Growth Rate. *Biological Reviews*, 86(1), 97 – 116. <u>https://doi.org/10.1111/j.1469-185x.2010.00136.x</u>

Đukanović, N., la Spada, F., Emmenegger, Y., Niederhäuser, G., Preitner, F., Franken, P. 2022. Depriving Mice of Sleep also Deprives of Food. *Clocks & Sleep*, 4(1), 37 – 51. <u>https://doi.org/10.3390/clockssleep4010006</u>

Dunlap, K.D. 1995. Hormonal and Behavioral Responses to food and Water Deprivation in a Lizard (*Sceloporus occidentalis*): Implications for Assessing Stress in a Natural Population. *Journal of Herpetology*, 29(3), 345 – 351. <u>https://doi.org/10.2307/1564983</u>

Eagan, T. 2018. Evaluation of Enrichment for Reptiles in Zoos. *Journal of Applied Animal Welfare Science*, 22(1), 69 – 77.

Gaspari, S. 1994. Haul-Out Behaviour, Site Fidelity and Vigilance of Common Seals (*Phoca vitulina*) and Grey Seals (*Halichoerus grypus*) in the Tees Estuary. *PhD. Thesis. Durham University, Durham, United Kingdom.* 

Guy, A.J., Curnoe, D., Banks, P.B. 2013. A Survey of Current Mammal Rehabilitation and Release Practices. *Biodiversity and Conservation*, 22, 825 – 837.

Hall, A.J., McConnell, B.J., Barker, R.J. 2002. The Effect of Total Immunoglobulin Levels, Mass, and Condition on the First-Year Survival of Grey Seal Pups. *Functional Ecology*, 16(4), 462 – 474. <u>https://doi.org/10.1046/j.1365-2435.2002.00649.x</u>

Hall, A.J., McConnell, B.J., Barker, R.J. 2008. Factors Affecting First-Year Survival in Grey Seals and Their Implications for Life History Strategy. *Journal of Animal Ecology*, 70(1), 138 – 149. <u>https://doi.org/10.1111/j.1365-2656.2001.00468.x</u>

Harding, K.C., Fujiwara, M., Axberg, Y., Härkönen, T. 2005. Mass-Dependent Energetics and Survival in Harbour Seal Pups. *Functional Ecology*, 19(1), 129 – 135. <u>https://doi.org/10.1111/j.0269-8463.2005.00945.x</u>

Harding, K.C., Härkönen, T., Helander, B., Karlsson, O. 2007. Status of Baltic Grey Seals: Population Assessment and Extinction Risk. *NAMMCO Scientific Publications*, 6, 33 – 56. <u>https://doi.org/10.7557/3.2720</u>

Härkönen, T., Dietz, R., Reijnders, P., Teilmann, J., Harding, K., Hall, A., Brasseur, S., Siebert, U., Goodman, S.J., Jepson, P.D, Dau Rasmussen, T., Thompson, P. 2006. The 1998 and 2002 Phcoine Distemper Virus Epidemics in European Harbour Seals. *Disease of Aquatic Organisms*, 68, 115 – 130.

Harris, R.B.S., Zhou, J., Youngblood, B.D., Rybkin, I.I., Smahin, G.N., Ryan, D.H. 1998. Effect of Repeated Stress on Body Weight and Body Composition of Rats Fed Low- and High-Fat Diets. *Regulatory and Integrative Physiology*, 275(6), R1928 – R1938. <u>https://doi.org/10.1152/ajpregu.1998.275.6.R1928</u>

Harper, D.G.C. 1994. Some Comments on the Repeatability of Measurements. *Ringing and Migration*, 51, 84 – 90.

Hector, K.L., Nakagawa, S. 2012. Quantitative Analysis of Compensatory and Catch-Up Growth in Diverse Taxa. *Journal of Animal Ecology*, 81(3), 583 – 593. https://doi.org/10.1111/j.1365-2656.2011.01942.x

Herborn, K.A., Heidinger, B.J., Boner, W., Noguera, J.C., Adam, A., Daunt, F., Monaghan, P. 2014. Stress Exposure in Early Post-Natal Life Reduced Telomere Length: an Experimental Demonstration in a Long-Lived Seabird. *Proceedings of the Royal Society Biology: Biological Sciences*, 281(1782), 20133151. https://doi.org/10.1098/rspb/2013.3151

Hoy, R.A., Brereton, J.E. 2022. Does Observer Presence Modify the Behavior and Enclosure Use of Captive Edwards' Pheasants? *Journal of Zoological and Botanical Gardens*, 3(2), 147 – 157. <u>https://doi.org/10.3390/jzbg3020012</u>

Inness, C.L.W., Metcalfe, N.B. 2008. The Impact of Dietary Restriction, Intermittent Feeding, and Compensatory Growth on Reproductive Investment and Lifespan in a Short-Lived Fish. *Proceedings of the Royal Society Biology: Biological Sciences*, 275(1644). <u>https://doi.org/10.1098/rspb.2008.0357</u>

Ivy, A.S., Brunson, K.L., Sandman, C., Baram, T.Z. 2008. Dysfunctional Nurturing Behavior in Rat Dams with Limited Access to Nesting Material: A Clinically Relevant Model for Early-Life Stress. *Neuroscience*, 154(3), 1132 – 1142.
<u>https://doi.org/10.1016/j.neuroscience.2008.04.019</u>

Jacquier, M., Crauste, F., Soulage, C.O., Soula, H.A. 2014. A Predictive Model of the Dynamics of Body Weight and Food Intake in Rats Submitted to Caloric Restrictions. *PLOS One*, 9(7), e103664. <u>https://doi.org/10.1371/journal.pone.0100073</u>

Jarvis, d., Marsh, S. 2017. An Overview of Current Issues in Seal Rescue and Rehabilitation in the U.K. *British Divers Marine Life Rescue*, Presentation for European Cetacean Society 2017 Conference. <u>https://bdmlr.org.uk/wp-</u> content/uploads/2020/05/ECS-2018-poster.pdf

Jayakody, S., Sibbald, A.M., Gordon, I.J., Lambin, X. 2008. Red Deer *Cervis elaphus* vigilance behaviour differs with habitat and type of human disturbance. *Wildlife Biology*, 14(1), 81 – 91. <u>https://doi.org/10.2981/0909-</u> <u>6396(2008)14[81:RDCEVB]2.0.CO;2</u>

Jenssen, B.M., Åsmul, J.I., Ekker, M., Vongraven, D. 2010. To Go for a Swim or Not? Consequences of Neonatal Aquatic Dispersal Behaviour for Growth in Grey Seal Pups. *Animal Behaviour*, 80(4), 667 – 673. <u>https://doi.org/10.1016/j.anbehav.2010.06.028</u>

Kannan, G., Terrill, T.H., Kouakou, B., Gazal, O.S., Gelaye, S., Amoah, E.A., Samaké, S. 2000. Transportation of Goats: Effects on Physiological Stress Responses and Live Weight Loss. *Journal of Animal Science*, 78(6), 1450 – 1547. <u>https://doi.org/10.2527/2000.7861456.x</u>

Kershaw, J.L., Hall., A.J. 2016. Seasonal Variation in Harbour Seal (*Phoca vitulina*) Blubber Cortisol – A Novel Indicator of Physiological State? *Scientific Reports*, 6, 21889. Krieber, M., Barrette, C. 1984. Aggregation Behaviour of Harbour Seals at Forillon National Park, Canada. *Journal of Animal Ecology*, 53(3), 913 – 928. <u>https://doi.org/10.2307/4667</u>

Leger, D., Bayon, V., de Sanctis, A. 2015. The Role of Sleep in the Regulation of Body Weight. *Molecular and Cellular Endocrinology*, 418(2), 101 – 107. <u>https://doi.org/10.1016/j.mce.2015.06.030</u>

Lidgard, D. 2003. Intraspecific Variation in Male Mating Behaviour in the Grey Seal: Linking Phenotype, Behaviour and More. *PhD Thesis, University of La Rochelle, La Rochelle, France.* 

MacRae, A.M., Haulena, M., Fraser, D. 2010. The Effect of Diet and Feeding Level on Survival and Weight Gain of Hand-Raised Harbour Seal Pups (*Phoca vitulina*). *Zoo Biology*, 30(5), 532 – 541. <u>https://doi.org/10.1002/zoo.20356</u>

Malik, S., Spencer, S.J. 2019. Early Life Stress and Metabolism. *Current Opinion in Behavioral Sciences*, 28, 25 – 30. <u>https://doi.org/10.1016/j.coheba.2019.01.016</u>

Mangel, M., Stamps, J. 2001. Trade-Offs Between Growth and Mortality and the Maintenance of Individual Variation in Growth. *Evolutionary Ecology Research*, 3, 583 – 593.

Maniam, J., Morris, M.J. 2012. The Link Between Stress and Feeding Behaviour. *Neuropharmacology*, 63(1), 97 – 110. <u>https://doi.org/10.1016/j.neuropharm.2012.04.017</u>

Matisz, C.E., Bandenhorst, C.A., Gruber, A.J. 2021. Chronic Unpredictable Stress Shifts Rat Behavior from Exploration to Exploitation. *The International Journal on the Biology of Stress*, 24(5), 635 – 644. <u>https://doi.org/10.1080/10253890.2021.1947235</u>

Merrick, M.J., Koprowski, J.L. 2017. Should We Consider Individual Behavior Differences in Applied Wildlife Conservation Studies? *Biological Conservation*, 209, 34 – 44. <u>https://doi.org/10.1016/j.biocon.2017.01.021</u>

McLaren, G.W., Mathews, F., Fell, R., Gelling, M., Macdonald, D.W. 2023. Body Weight Change as a Measure of Stress: A Practical Test. *Animal Welfare*, 13(3).

Mellish, J.A., Iverson, S.J., Don Bowen, W. 1999. Variation in Milk Production and Lactation Performance in Grey Seals and Consequences for Pup Growth and Weaning Characteristics. *Physiological and Biochemical Zoology*, 72(6). <u>https://doi.org/10.1086/316708</u>

Moe, B., Brunvoll, S., Mork, D., Brobakk, T.E., Bech, C. 2004. Developmental Plasticity of Physiology and Morphology in Diet-Restricted European Shag Nestlings (*Phalacrocorax aristotelis*). *Journal of Experimental Biology*, 207(23), 4067 – 4076. https://doi.org/10.1242/jeb.01226

Molinari-Jobin, A., Zimmermann, F., Borel, S., Le Grand, L., Iannino, E., Andres, O., Belotti, E., Bufka, L., Ćirović, D., Drouet-Hoguet, N., Engleder, T., Figura, M., Fuxjäger, C., Gregorova, E., Heuricj, M., Idelberger, S., Kubala, J., Kusak, J., Melovski, D., Middlehof, T.L., Mináriková, T., Molinari, P., Mouzon-Moyne, L., Moyne, G., Mysłajek, R.W., Nowak, S., Ozolins, J., Ryser, A., Sanaja, B., Shkvyria, M., Sin, T., Sindičić, M., Slijepčević, V., Stauffer, C., Tám, B., Trajce, A., Volfová, J., Wölfl, S., Zlatanova, D., Vogy, K. 2024. Rehabilitation and Release of Orphaned Eurasian Lynx (*Lynx lynx*) in Europe: Implications for Management and Conservation. *PLoS One*, 19(3), e0297789. <u>https://doi.org/10.1371/journal.pone.0297789</u>

Nakagawa, S., Schielzeth, H. 2010. Repeatability for Gaussian and Non-Gaussian Data: A Practical Guide for Biologists. *Biological Reviews*, 85(4). <u>https://doi.org/10.1111/j.1469-185X.2010.00141.x</u>

Noren, S.R. 2020. Postnatal Development of Diving Physiology: Implications of Anthropogenic Disturbance for Immature Marine Mammals. *Journal of Experimental Biology*, 223(7), jeb227736. <u>https://doi.org/10.1242/jeb/227736</u>

Noren, S.R., Boness, D.J., Iverson, S.J., McMillan, J., Don Bowen, W. 2008. Body Condition at Weaning Affects the Duration of the Postweaning Fast in Gray Seal Pups (*Halichoerus grypus*). *Physiological and Biochemical Zoology*, 81(3). <u>https://doi.org/10.1086/528777</u>

O'Hara, A.D. 2019. Does Rehabilitation Work? Investigating Sex and Species-Specific Differences in Daily Weight Gain of Harbour and Grey Seal Pups in rehabilitation. *Bachelor of Science Dissertation, BSc Dissertation, University of Bangor, Bangor, United Kingdom.* 

Osinga, N., Ferdous, M.M.S., Morick, D., Garćia Hartmann, M., Ulloa, J.A., Vedder, L., Udo de Haes, H.A., Brakefield, P.M., Osterhaus, A.D.M.E., Kuiken, T. 2012. Patterns of Stranding and Mortality in Common Seals (*Phoca vitulina*) and Grey Seals (*Halichoerus grypus*) in The Netherlands between 1979 and 2008. *Journal of Comparative Pathology*, 147(4), 550 – 565. <u>https://doi.org/10.1016/j.jcpa.2012.04.001</u>

Paterson, W.D., Russell, D.J.F., Wu, G-M., McConnell, B., Currie, J.I., McCafferty, D.J., Thompson, D. 2019. Post-Disturbance Haulout Behaviour of Harbour Seals. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(1), 144 – 156. <u>https://doi.org/10.1002/aqc.3092</u>

Reeder, D.M., Kramer, K.M. 2005. Stress in Free-Ranging Mammals: Integrating Physiology, Ecology, and Natural History. *Journal of Mammalogy*, 86(2). 225 – 235. <u>https://doi.org/10.1644/BHE-003.1</u>

Reidman, M. 1990a. Maternal Care and Lactation Strategies, in: Reidman, M. The Pinnipeds: Seals, Sea Lions, and Walruses. University of California Press, Oxford, England, pp. 277.

Reidman, M. 1990b. The Pinnipeds: Seals, Sea Lions, and Walruses. University of California Press, Oxford, England, pp. 142.

Romero, L.M., Beattie, U.K. 2022. Common Myths of Glucocorticoid Function in Ecology and Conservation. *JEZ-A Ecological and Integrative Physiology*, 377(1), 1 – 106. <u>https://doi.org/10.1002/jez.2459</u>

Rudin, S.F., Tomkins, J.L., Simmons, L.W. 2018. The Effects of the Social Environment and Physical Disturbance on Personality Traits. *Animal Behaviour*, 138, 109 – 121. <u>https://doi.org/10.1016/j.anbehv.2018.02.013</u>

Ruppert, N. 2017. Effects of environmental enrichment on activities of captive orangutans at Taiping Zoo, Malaysia. *Mayalan Nature Journal*, 69(4), 327 – 335.

Salazar-Casals, A., de Reus, K., Grsekewitz, N., Havermans, J., Guet, M., Villanueva, S., Rubio-Garcia, A. 2022. Increased Incidence of Entanglements and Ingested Marine Debris in Dutch Seals from 2010 to 2020. *Oceans*, 3(3), 289 – 400. <u>https://doi.org/10.3390/oceans3030026</u>

Sayer, S., Allen, R., Bellman, K., Beaulieu, M., Cooper, T., Dyer, N., Hockin, K., Hockley, K., Jarvis, D., Jones, G., Oaten, P., Waddington, N., Witt, M.J., Hawkes, L. 2021. Post Release Monitoring of Rehabilitated Gray Seal Pups Over Large Temporal and Spatial Scales. *Marine Mammal Science*. <u>https://doi.org/10.1111/mms.12885</u>

Schumann, K., Guenther, A., Jewgenow, K., Trillmich, F. 2014. Animal Housing and Welfare: Effects of Housing Conditions on Body Weight and Cortisol in a Medium-Sized Rodent (*Cavia aperea*). *Journal of Applied Animal Welfare Science*, 17(2), 111 – 124. <u>https://doi.org/10.1080/10888705.2014.884407</u>

Shaughnessy, P.D., Semmelink, A., Cooper, J., Frost, P.G.H. 1981. Attempts to Develop Acoustic Methods of Keeping Cape Fur Seals *Arctocephalus pusillus* form Fishing Nets. *Biological Conservation*, 21(2), 141 – 158. <u>https://doi.org/10.1016/0006-</u> 3207(81)90076-8

Short, M.A., Banks, S. 2013. The Function Impact of Sleep Deprivation, Sleep Restriction, and Sleep Fragmentation. In: Bianchi, M. (eds) *Sleep Deprivation and Disease*. Springer, New York, NY. <u>https://doi.org/10.1007/978-1-4614-9087-6\_2</u>

Silpa, M.A.C., Thornton, S.M., Cooper, T., Hedley, J. 2015. Prevalence of Presenting Conditions in Grey Seal Pups (*Halichoerus grypus*) Admitted for Rehabilitation. *Veterinary Sciences*, 2(1), 1 – 11. <u>https://doi.org/10.3390/vetsci.2010001</u>

Silva, W.T.A.F., Harding, K.C., Marques, G.M., Bäcklin, B.M., Sonne, C., Dietz, R., Kauhala, K., Desforges, J. 2020. Life Cycle Bioenergetics of the Gray Seal (*Halichoerus grypus*) in the Baltic Sea: Population Response to Environmental Stress. *Environmental International*, 145, 106145. <u>https://doi.org/10.1016/j.envint.2020.106145</u>

Spotte, S., Stake, P.E. 1982. Hand-Rearing of Twin Grey Seals (*Halichoerus Grypus*) from Birth to Weaning. *Marine Ecology Progress Series*, 9(2), 181 – 189. <u>https://doi.org/10.3354/meps009181</u> Stead, S.M., Bådescu, I., Boonstra, R. 2021. Of Mammals and Milk: How Maternal Stress Affects Nursing Offspring. *Mammal Review*, 52(1), 129 – 147. <u>https://doi.org/10.1111/mam.12267</u>

Surviliene, V., Ruksenas, O., Pomeroy, P. 2016. Play Behavior of Wild Grey Seals (*Halichoerus grypus*): Effect of Haulout Group Size and Composition. *Aquatic Mammals*, 42(2), 144 – 161. <u>https://doi.org/10.1578/AM.42.2.2016.144</u>

Taylor, A.A., Davis, H., Boyle, G.J. 1998. Increased Vigilance Toward Unfamiliar Humans by Harbour (*Phoca vitulina*). *Marine Mammal Science*, 14(3), 575 – 583. <u>https://doi.org/10.1111/j.1748-7692.1998.tb00743.x</u>

Twiss, S., Bishop, A., Culloch, R. 2022. The Gray Seal: 80 Years of Insight into Intrinsic and Extrinsic Drivers of Phocid Behaviour. In. Costa, D.P., McHuron, E.A. (eds) Ethology and Behavioural Ecology of Phocids. Springer, Cham. <u>https://doi.org/10.1007/978-3-030-88923-4\_9</u>

Voellmy, I.K., Goncalves, I.G., Barrette, M., Monfort, S.L., Manser, M.B. 2014. Mean Feval Glucocorticoid Metabolites are Associated with Vigilance, Whereas Immediate Cortisol Levels Better Reflect Acute Anti-Predator Responses in Meerkats. *Hormones and Behavior*, 66(5), 759 – 765. <u>https://doi.org/10.1016/j.yhbeh.2014.08.008</u>

Yam, K.Y., Naninck, E.F.G., Abbink, M.R., la Fleur, S.E., Schipper, L., van den Beukel, J.C., Grefhorst, A., Oosting, A., van der Beek, E.M., Lucassen, P.J., Korosi, A. 2017. Exposure to Chronic Early-Life Stress Lastingly Alters the Adipose Tissue, the Leptin System and Changes the Vulnerability to Western-Style Diet Later in Life in Mice. *Psychoneuroendocrinology*, 77, 186-195.

https://doi.org/10.1016/j.psyneuen.2016.12.012

Waples, K.A., Gales., N.J. 2002. Evaluating and Minimising Social Stress in the Care of Captive Bottlenose Dolphins (*Tursiops aduncus*). *Zoo Biology*, 21(1), 5 – 26. <u>https://doi.org/10.1002/zoo.10004</u>

Wells, R.S., Fauquier, D.A., Gulland, F.M.D., Townsend, F.I., DiGiovanni Jr, R.A. 2012. Evaluating PostIntervention Survival of Free-Ranging Odontocete Cetaceans. *Marine Mammal Science*, 29(4), E463 – E483. <u>https://doi.org/10.1111/mms.12007</u>

Zatrak, M., Brittain, S., Himmelreich, L., Lovick-Earle, S., Pizzi, R., Shaw, K.J., Grant, R.A., Geary, M. 2022. Factors Affecting the Survival of Harbor (*Phoca vitulina*) and Gray Seal (*Halichoerus grypus*) Juvenilles Admitted for Rehabilitation in the UK and Ireland. *Marine Mammal Science*, 39(2), 462 – 480. <u>https://doi.org/10.1111/mms.12983</u>

# Software:

R Core Team. 2023. R: A Language and Environment for Statistical Computing (Version 2023.12.0+369). R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/ Stoffel, M.A., Nakagawa, S., Schielzeth, H. 2017. rptR: Repeatability Estimation and Variance Decomposition by Generalized Linear Mixed-Effects Models. *Methods in Ecology and Evolution*, 8(11), 1639 – 1644. <u>https://doi.org/10.1111/2041-210X.12797</u>

Bates, D., Mäechler, M., Bolker, B., Walker, S. 2015. Fitting Linear Mixed-Effects Models Using Ime4. *Journal of Statistical Software*, 67(1), 1 – 48. <u>https://doi.org/10.18637/jss.v067.i01</u>

Fox, J., Weisberg, S. 2019. An R Companion to Applied Regression, third ed. Thousand Oaks, California. <u>https://doi.org/10.18637/jss.v067.i01</u>

Lüdecke, D. 2023. sjPlot: Data Visualisation for Statistics in Social Science (Version 2.8.15). <u>https://CRAN.R-project.org/package=sjPlot</u>

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H. 2019. Welcome to the tidyverse. *Journal of Open Source Software*, 4(43), 1686. <u>https://doi.org/10.21105/joss.01686</u>

Websites:

British Veterinary Zoological Society. 2016. Good Practice Guidelines for Wildlife Rehabilitation Centres. <u>https://www.rcvs.org.uk/document-library/bvzs-good-practice-guidelines-for-wildlife-rehabilitation/?&&type=rfst&set=true#cookie-widget</u> (accessed 4 March 2024).

Cornish Seal Sanctuary. 2020. Into the Wild. <u>https://sealsanctuary.sealifetrust.org/en/about-us/blog/into-the-wild/</u> (accessed 5 February 2024)

Cornish Seal Sanctuary. 2024. Grey Seals.

https://sealsanctuary.sealifetrust.org/en/about-us/grey-seals/ (accessed 2 February 2024)

Cumbria Wildlife Trust. 2023. Grey Seal Pup Season. <u>https://www.cumbriawildlifetrust.org.uk/blog/grey-seal-pup-season</u> (accessed 30 January 2024)

North Wales Wildlife Trust. 2021a. Grey Seal – Mating. <u>https://www.northwaleswildlifetrust.org.uk/blog/living-seas/grey-seal-</u> <u>mating#:~:text=Grey%20seals%20(Halichoerus%20grypus)%20will,mate%20with%20a</u> <u>%20dominant%20male</u> (accessed 30 January 2024)

North Wales Wildlife Trust. 2021b. Grey Seal – Moulting Season. <u>https://www.northwaleswildlifetrust.org.uk/blog/living-seas/grey-seal-moulting-</u> <u>season#:~:text=Grey%20seals%20(Halichoerus%20grypus)%20begin,a%20year%20duri</u> <u>ng%20this%20season</u> (accessed 30 January 2024) North Wales Wildlife Trust. 2021c. Grey Seal BLOG – Pupping Season.

https://www.northwaleswildlifetrust.org.uk/blog/living-seas/grey-seal-blog-puppingseason (accessed 30 January 2024)

Seal Rescue Ireland. 2019. Rehabilitation and Release.

https://www.sealrescueireland.org/our-work/rehab-release/ (accessed 30 January 2024)

NOAA, 2022. Grey Seal. <u>https://www.fisheries.noaa.gov/species/gray-seal</u> (accessed 27 March 2024)

### Appendix:

### A.1: Behavioural Statistical Methods:

Behaviour was analysed using generalised mixed effects models (GLMM's) on Night (n = 90 observations, N = 27 seals), and Day (n = 334 observations, N = 30 seals), using package "lme4" (Bates et al., 2015), for three behaviours (head elevated, movement, and rest). Aggression was not analysed due to rarity of behaviour (Night: n = 2 observations; Day: n = 28 observations).

Fixed independents variables for each dataset (See Table A1 for reference levels); Day: pool, hour, water level, staff feed, staff intensity, and in proximity of another seal; Night: pool, time period, weather, and in proximity of another seal. Individual ID was included in each analysis as a random effect as it was determined to be an important variable through comparing AIC numbers (see Supplementary 4 for further details). All unidentifiable individuals were removed from the analysis. Confounding effects were tested for using package "car" (Fox and Weisberg, 2019).

Fixed Independent Variable	Reference Level	Data Analysis Used In
Pool	Pre-release Pool.	Both.
In Proximity of Another Seal	Not in proximity of another seal.	Both.
Hour	12:00 – 13:00.	Day.
Water Level	No water in the pool (0).	Day.
Staff feed	No staff feed occurring.	Day.
Staff intensity	No staff intensity (level 0).	Day.
Time Period	00:00 – 01:00 (Time period 1).	Night.
Weather	Dry	Night.

Table A1: Reference levels for fixed independent variables for behavioural generalised linear mixed effects models (GLMER's).

Consistency repeatability estimates, following Nakagawa and Schielzeth, 2010, determined if the behaviours were behavioural syndromes, using package "rptR" (Stoffel et al., 2017). Estimated adjusted repeatability (an estimate adjusting for confounding effects) was used as there were confounding effects within the GLMM's. To be classed as a behavioural syndrome, the estimates needed to be higher than 0.2 (Harper 1994).

See results in Supplementary 5.

# A.2: Justification for removing certain admittance reasons from the model.

Admittance Reasons	Reasons for exclusion
Premature	Rarity of occurrence (n = $29$ ) and potential inconsistent reporting.
Harassed	Rarity of occurrence (n = 40) and potential inconsistent reporting.
Dehydrated	Correlated with injured (Pearson's chi-square test: X-squared = 0.22, p-value = 0.64) and illness (Pearson's chi-square test: X-squared = 0, p-value = 1).
Emaciated	Combined with underweight due to and potential inconsistent reporting.
Bad location	Rarity of occurrence (n = 19), and colinear with age (Pearson's chi-square test: X-squared = $2.57$ , df = 2, p-value = $0.2767$ ).
Entanglement	Combined with injured due to close similarity in definitions.
Orphaned	Potential inconsistent reporting: individuals in non- dependent age class (already abandoned) have this admittance reason (n = 3).

Table A2: Justification for excluding admittance reasons from analyses:

A.3: Six Consecutive Pupping Seasons Main Weight Results Figures.



Figure A3: Key significant variables influencing weight gain over six consecutive pupping seasons (N = 292 seals, n = 3433 weight measurements; full model reported at Table 5). (A) The actual weight (kgs) of an individual is positively correlated with weight gain between days. (B) The duration of rehabilitation is negatively correlated with weight gain between days. (C) Compared to individuals without an illness, individuals admitted with an illness had lower weight gain between days. (D) Compared to individuals not underweight, individuals that were underweight had a greater weight gain between days. In A and B, points represent individual weight measurements (regression line is illustrative). In C and D, the line represents the standard error, the box represents main data and the points represents any extraneous individuals.

A.4: Case Study Pupping Season Main Weight Results Figures.



Figure A4: Main significant variables influencing weight gain within the case study pupping seasons (N = 67 seals, n = 378 weight measurements; full model reported at Table 6). (A) The actual weight (kgs) of an individual is positively correlated with weight gain between days. (B) The duration of rehabilitation is negatively correlated with weight change between days. (C) Compared to an individual that is not injured, individuals admitted with an injury had higher gain between days. (D) Although marginally significant, compared to Pre-Release Pool (shielded pool), individuals within Rock Pool gain more weight between days. In A and B, points represent individual weight measurements (regression line is illustrative). In C and D, the line represents the standard error, the box represents main data, and the points represents any extraneous individual.

## **Supplementary Information:**

# S.1: Definitions of Reason for Admittance:

Table S1: Definitions of the Different Admittance Reasons for Grey Seals (*Halichoerus grypus*) Admitted to Seal Rescue Ireland, Ireland. These definitions have been provided through personal communication with Seal Rescue Ireland. These can vary slightly between people admitting the seal.

Admittance Reason	Definition				
Emaciated	Seal is extremely underweight for their age, with very visible bones and rolls of skin. Young pups severely under normal birth weight around 15kgs are also classed as emaciated.				
Underweight	Seal is underweight, has some visible bones and rolls of skin, closer to their birth weight for young pups. If weaned, greys coming in around 18-20kg are underweight. It depends on the body condition relative to the seal's size (length) and rough age.				
Illness	Seals with evident parasites e.g., lungworms, roundworms or if the seal has alopecia, or with virus e.g., phocine distemper virus, seal pox				
Injured	This ranges from small scratches/grazes anywhere on the body to large open wounds.				
Dehydrated	Seals who have no eye rings and very small-looking eyes.				
Orphaned	Seals who are dependent on their mother's milk to survive and have been separated prematurely before being weaned.				
Premature	Grey seals born small and without teeth are classed as premature.				
Entanglement	Seals with active entanglement or evidence that there was an ex-entanglement (scarring).				
Bad Location	Seal that seems relatively healthy and can survive without rehab will need to be relocated, for example seal is on a busy beach and being harassed, or the seal has found itself far from the sea etc.				
Harassed	Seals healthy, injured, sick or orphaned that are approached continually by members of the public, lifted by members of the public etc, not allowing the seal to rest.				
Unknown	Admittance reason has not been recorded or it does not fall into one of the above categories.				

### S.2: Camera locations



Figure S1 (left): Reolink Go PT Plus (Reolink, China) in Nursery pool (camera 2). Camera is attached to the pole using supplied strap. Solar panel charges the camera and is placed in a place that will get direct sunlight to be able to charge the cameras.

Figure S2 (Right): Reolink Go PT Plus (Reolink, China) in Nursery pool (camera 3). Camera is attached to the pole using supplied strap. Solar panel charges the camera and is placed in a place that will get direct sunlight to be able to charge the cameras.





Figure S3 (left): Reolink Go PT Plus (Reolink, China) in Nursery pool (camera 4). Camera is attached to the pole using supplied strap. Solar panel charges the camera and is placed in a place that will get direct sunlight to be able to charge the cameras.

### S.3: AIC Numbers for Comparison of Behavioural Models

An anova test was used to compare the two models and look at the importance of the

random effect within the models: GLM and GLMER. The random effect in the GLMER

model was Field ID.

### S.3.1: Combined Behavioural Dataset:

Table S3.1.1: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Movement Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	7	1306.1	1334.3	-646.04	1292.1			
GLMER	8	1263.5	1295.8	-623.77	1274.5	44.534	1	2.5e-11

Table S3.1.2: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Resting Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	7	1343.8	1372	-664.91	1329.8			
GLMER	8	1286.7	1318.9	-635.35	1270.7	59.13	1	1.476e-14

Table S3.1.3: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Head Elevated Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	7	1259.6	1287.8	-622.80	1245.6			
GLMER	8	1247.9	1280.1	-615.94	1231.9	13.719	1	0.0002123

S.3.2: Day Behavioural Dataset:

Table S3.2.1: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Movement behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	14	1220.0	1273.4	-596.01	1192.0			
GLMER	15	1178.6	1235.7	-574.28	1148.6	43.455	1	4.338e-11

Table S3.2.2: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Resting Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	14	1283.9	1337.2	-627.95	1255.9			
GLMER	15	1229.9	1287.1	-599.96	1199.9	55.97	1	7.358e-14

Table S3.2.3: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Head Elevated Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	14	1021.6	1074.9	-496.79	993.57			
GLMER	15	1012.9	1070.1	-491.44	982.89	10.683	1	0.001081

### S.3.3: Night Behavioural Dataset:

Table S3.3.1: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Movement behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	8	296.69	316.68	-140.34	280.69			
GLMER	9	269.50	292.00	-125.75	251.50	29.184	1	6.582e-08

Table S3.3.2: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Resting Behaviour:

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	6	209.01	224.01	-98.505	197.01			
GLMER	7	193.84	211.34	-89.920	179.84	17.17	1	3.418e-05

Table S3.3.3: Anova Test for AIC Numbers for Comparison of GLM to GLMER Models for Head Elevated Behaviour.

Model	npar	AIC	BIC	logLik	Deviance	Chisq	DF	Pr(>Chisq)
GLM	8	661.10	681.10	-322.55	645.10			
GLMER	9	352.16	374.66	-167.08	334.16	310.95	1	< 2.2e-16
# S.4: Behaviour Model Results (Including Repeatability).

## S4.1: Day Data:

## S4.1.1: Movement Behaviour:

Table S4: Variables Influencing the likelihood of observing movement behaviour in pupping year 2022-2023 (n = 30 seals, N = 334 observations). Reference level: Pre-Release Pool, 12:00 – 13:00, little to no water present in the pool, no staff feeds, no intensity of staff in the pool, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.18, ICC = 0.05, Marginal R<sup>2</sup> = 0.094, Conditional R<sup>2</sup> = 0.140.

Predictors	Estimate	Confidence interval	p-value
(Intercept)	0.26	0.14 - 0.50	<0.001
Nursery Pool	0.85	0.52 – 1.38	0.509
Rock Pool	1.67	1.15 – 2.41	0.006
Physio Pool	2.22	1.44 - 3.42	<0.001
13:00 - 14:00	1.38	1.01 - 1.88	0.044
14:00 - 15:00	1.52	1.11 - 2.06	0.008
15:00 – 16:00	1.57	1.15 – 2.15	0.004
Shoulder Depth Water Level	0.89	0.52 – 1.53	0.673
Swimmable Water Level	1.66	0.98 – 2.79	0.059
Staff Feed (Yes)	1.22	0.72 – 2.07	0.454
Low Intensity of Staff	1.28	0.82 - 2.01	0.276
Medium Intensity of Staff	0.63	0.36 - 1.09	0.099
High Intensity of Staff	1.87	0.89 – 3.96	0.100
In Proximity of Another Seal	0.65	0.53 - 0.81	<0.001

Table S5: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Movement Behaviour.

R	0.024
Standard Error	0.012
Confidence Interval	[0, 0.047]
p-value	0.000926 [LRT]
	NA [Permutation]

### S4.1.2: Rest Behaviour:

Table S6: Variables influencing the likelihood of observing resting behaviour in pupping year 2022-2023 (n = 30 seals, N = 334 observations). Reference level: Pre-Release Pool, 12:00 – 13:00, little to no water present in the pool, no staff feeds, no intensity of staff in the pool, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.22, ICC = 0.06, Marginal R<sup>2</sup> = 0.082, Conditional R<sup>2</sup> = 0.139.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	2.28	1.24 - 4.20	0.008
Nursery Pool	0.70	0.43 - 1.14	0.152
Rock Pool	0.34	0.23 – 0.49	<0.001
Physio Pool	0.52	0.33 – 0.82	0.005
13:00 - 14:00	0.72	0.53 – 0.96	0.028
14:00 – 15:00	0.71	0.53 – 0.95	0.022
15:00 – 16:00	0.53	0.39 – 0.72	<0.001
Shoulder Depth Water Level	1.04	0.63 – 1.72	0.868
Swimmable Water Level	0.96	0.60 - 1.56	0.879
Staff Feed (Yes)	0.88	0.52 - 1.50	0.647
Low Intensity of Staff	1.14	0.73 – 1.77	0.577
Medium Intensity of Staff	1.63	0.99 – 2.69	0.056
High Intensity of Staff	0.93	0.45 - 1.91	0.844
In Proximity of Another Seal	1.51	1.22 - 1.86	<0.001

Table S7: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Rest Behaviour.

Link-Scale Approximation

,,	
R	0.031
Standard Error	0.014
Confidence Interval	[0.002, 0.059]
p-value	0.000236 [LRT]
	NA [Permutation]

#### S4.1.3: Head Elevated Behaviour:

Table S8: Variables influencing the likelihood of observing head elevated behaviour in pupping year 2022-2023 (n = 30 seals, N = 334 observations). Reference level: Pre-Release Pool, 12:00 – 13:00, little to no water present in the pool, no staff feeds, no intensity of staff in the pool, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.12, ICC = 0.03, Marginal R<sup>2</sup> = 0.079, Conditional R<sup>2</sup> = 0.111.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	0.45	0.23 – 0.89	0.022
Nursery Pool	0.65	0.38 - 1.12	0.118
Rock Pool	1.62	1.11 – 2.38	0.013
Physio Pool	0.53	0.34 - 0.82	0.005
13:00 – 14:00	0.94	0.66 - 1.33	0.711
14:00 – 15:00	0.82	0.58 - 1.16	0.264
15:00 – 16:00	1.45	1.03 - 2.06	0.033
Shoulder Depth Water Level	0.96	0.55 - 1.68	0.881
Swimmable Water Level	0.62	0.35 - 1.08	0.090
Staff Feed (Yes)	0.99	0.55 – 1.80	0.984
Low Intensity of Staff	1.10	0.68 - 1.76	0.702
Medium Intensity of Staff	2.13	1.23 - 3.66	0.007
High Intensity of Staff	0.52	0.23 - 1.19	0.122
In Proximity of Another Seal	0.93	0.73 - 1.18	0.560

Table S9: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Alertness (Head Up) Behaviour.

R	0.012
Standard Error	0.008
Confidence Interval	[0, 0.027]
p-value	0.0741 [LRT]
	NA [Permutation]

### S4.2: Night Data:

S4.2.1: Movement Behaviour:

Table S10: Variables influencing the likelihood of observing movement behaviour in pupping year 2022-2023 (n = 26 seals, N = 90 observations). Reference level: Pre-Release Pool, 00:00 – 01:00, dry, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 1.60, ICC = 0.33, Marginal R<sup>2</sup> = 0.121, Conditional R<sup>2</sup> = 0.408.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	0.03	0.01 - 0.10	<0.001
Nursery Pool	1.03	0.21 - 5.03	0.975
Rock Pool	0.67	0.17 – 2.58	0.561
Physio Pool	0.25	0.06 - 1.08	0.063
02:00 - 04:00	0.40	0.07 – 2.27	0.298
05:00 – 07:00	1.17	0.60 - 2.25	0.647
Rain	0.75	0.27 – 2.10	0.581
In Proximity of Another Seal	0.56	0.23 - 1.35	0.195

Table S11: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Movement Behaviour.

Link-Scale Approximation



S4.2.2: Rest Behaviour:

Table S12: Variables influencing the likelihood of observing resting behaviour in pupping year 2022-2023 (n = 26 seals, N = 90 observations). Reference level: Pre-Release Pool, dry, and not in proximity of another seal. Time period was removed from this analysis due to insufficient data within time period 02:00 – 04:00 (N = 600). Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 2.11, ICC = 0.39, Marginal R<sup>2</sup> = 0.176, Conditional R<sup>2</sup> = 0.497.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	42.61	10.13 - 179.31	<0.001
Nursery Pool	1.62	0.17 – 15.72	0.680
Rock Pool	6.87	1.10 - 43.03	0.040
Physio Pool	7.38	1.10 - 49.56	0.040
Rain	0.74	0.19 - 2.83	0.661
In Proximity of Another Seal	2.88	0.76 – 10.97	0.120

Table S13: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Rest Behaviour.

Link-Scale Approximation



S4.2.3: Head Elevated Behaviour:

Table S14: Variables influencing the likelihood of observing head elevated behaviour in pupping year 2022-2023 (n = 26 seals, N = 90 observations). Reference level: Pre-Release Pool, 00:00 – 01:00, dry, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 5.26, ICC = 0.62, Marginal R<sup>2</sup> = 0.340, Conditional R<sup>2</sup> = 0.746.

Predictors	Estimates	Confidence Interval	p-value
(Intercept)	0.16	0.04 - 0.71	0.016
Nursery Pool	0.93	0.17 - 5.11	0.932
Rock Pool	0.21	0.04 - 1.06	0.059
Physio Pool	0.03	0.00 - 0.25	0.001
02:00 - 04:00	0.27	0.06 - 1.24	0.092
05:00 – 07:00	4.49	2.92 - 6.90	<0.001
Rain	1.08	0.37 – 3.16	0.889
In Proximity of Another Seal	0.17	0.09 - 0.32	<0.001

Table S15: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Alertness (Head Up) Behaviour.

R	0.095
Standard Error	0.058
Confidence Interval	[0, 0.214]
p-value	0.00601 [LRT]
	NA [Permutation]

## S4.3: Combined Night and Day

### S4.3.1: Movement Behaviour:

Table S16: Variables influencing the likelihood of observing movement behaviour in pupping year 2022-2023 (n = 31 seals, N = 415 observations). Reference level: Pre-Release Pool, day, heron absent, and not in proximity of another seal. Random effects are individual ID. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.16, ICC = 0.05, Marginal R<sup>2</sup> = 0.477, Conditional R<sup>2</sup> = 0.502.

Predictors	Estimate	Confidence interval	p-value
(Intercept)	0.41	0.29 – 0.58	<0.001
Nursery Pool	1.05	0.66 - 1.66	0.835
Rock Pool	2.35	1.74 - 3.17	<0.001
Physio Pool	2.69	1.79 - 4.03	<0.001
Night	0.02	0.01 - 0.03	<0.001
Heron Present	1.12	0.88 - 1.42	0.346
In Proximity of Another Seal	0.67	0.55 – 0.81	<0.001

Table S17: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Movement Behaviour.

Link-Scale Approximation



S4.3.2: Rest Behaviour:

Table S18: Variables influencing the likelihood of observing resting behaviour in pupping year 2022-2023 (n = 31 seals, N = 415 observations). Reference level: Pre-Release Pool, day, heron absent, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.21, ICC = 0.06, Marginal R<sup>2</sup> = 0.508, Conditional R<sup>2</sup> = 0.537.

Predictors	Estimate	Confidence interval	p-value
(Intercept)	1.86	1.29 – 2.66	0.001
Nursery Pool	0.62	0.40 - 0.98	0.042
Rock Pool	0.33	0.24 - 0.44	<0.001
Physio Pool	0.48	0.32 - 0.74	0.001
Night	76.89	42.79 – 138.16	<0.001
Heron Present	1.12	0.64 - 1.03	0.088
In Proximity of Another Seal	1.45	1.20 - 1.76	<0.001

Table S19: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Rest Behaviour.

Link-Scale Approximation

R	0.017
Standard Error	0.009
Confidence Interval	[0, 0.033]
p-value	0.000122 [LRT]
	NA [Permutation]

S4.3.3: Head Elevated Behaviour:

Table S20: Variables influencing the likelihood of observing head elevated behaviour in pupping year 2022-2023 (n = 31 seals, N = 415 observations). Reference level: Pre-Release Pool, day, heron absent, and not in proximity of another seal. Random effects are individual ID, residual variance ( $\sigma^2$ ) = 3.29, Between subject variance ( $\tau_{00}$ ) = 0.15, ICC = 0.04, Marginal R<sup>2</sup> = 0.135, Conditional R<sup>2</sup> = 0.172.

Predictors	Estimate	Confidence interval	p-value
(Intercept)	0.29	0.21 - 0.40	<0.001
Nursery Pool	1.30	0.79 – 2.14	0.297
Rock Pool	1.48	1.11 – 1.97	0.008
Physio Pool	0.58	0.39 - 0.88	0.010
Night	0.29	0.22 – 0.39	<0.001
Heron Present	1.54	1.21 – 1.95	<0.001
In Proximity of Another Seal	0.84	0.69 - 1.03	0.097

Table S21: Link-Scale Approximation and Original-Scale Approximation for Repeatability of Alertness (Head Up) Behaviour.

R	0.008
Standard Error	0.005
Confidence Interval	[0, 0.018]
p-value	0.193 [LRT]
	NA [Permutation]